

Thu. Sep 12, 2024

Oral presentation | S1: Dynamics of igneous processes (Special Session)

10:00 AM - 12:00 PM JST | 1:00 AM - 3:00 AM UTC | ES Hall Higashiyama Campus

S1: Dynamics of igneous processes (Special Session)

Chairperson: Shumpei Yoshimura (Hokkaido University), Yuuki Hagiwara (Japan Agency for Marine-Earth Science and Technology), Morihisa Hamada (JAMSTEC)

10:00 AM - 10:20 AM JST | 1:00 AM - 1:20 AM UTC

[S1-01] Tungsten (VI) speciation in subduction-zone aqueous fluids and its significance for W stable isotope fractionation during slab dehydration

「招待講演」

*Naoko TAKAHASHI¹, Michihiko Nakamura², Shigeru Yamashita³, Hiroyuki Kagi¹ (1. UTokyo Sci., 2. Tohoku Univ. Sci., 3. Okayama Univ. IPM)

10:20 AM - 10:40 AM JST | 1:20 AM - 1:40 AM UTC

[S1-02] Reconstruction of mantle structures using mantle xenoliths from Nushan, SE-edge of Sino-Korean craton: How far does the subduction fluid reach?

「招待講演」

*Yuto SATO¹, Eiichi Takahashi² (1. JAMSTEC Kochi, 2. GIG)

10:40 AM - 10:55 AM JST | 1:40 AM - 1:55 AM UTC

[S1-03] Elucidating Deep processes leading to maar eruptions using mantle xenoliths in the West Eifel volcanic field

「発表賞エントリー」

*Masanari Arai¹, Michihiko Nakamura¹, Mayumi Mujin¹, Naoki Araya¹, Sando Sawa¹, Takayuki Nakatani², Mari Sumita³, Hans-Ulrich Schmincke³ (1. Tohoku Univ. Sci., 2. AIST, 3. GEOMAR)

10:55 AM - 11:10 AM JST | 1:55 AM - 2:10 AM UTC

[S1-04] Water contents and pressures of melts in unerupted felsic magma constrained by SEM-EDS analysis of homogenized melt inclusions in zircon

「発表賞エントリー」

*Taichi Kawashima¹, Kazuya Shimooka², Toko Fukui¹, Satoshi Saito¹ (1. Ehime Univ., 2. Kwansei Gakuin Univ.)

11:10 AM - 11:25 AM JST | 2:10 AM - 2:25 AM UTC

[S1-05] Permeability and pore microstructures on the diktytaxitic texture of Ogurayama lava dome, Towada volcano

*Natsuko Sekiya¹, Michihiko Nakamura², Ryosuke Sakurai³, Mayumi Mujin², Naoki Araya², Shingo Takeuchi⁴, Yukiko Suwa⁵ (1. Tohoku Univ. Sci., 2. Grad. Sch. of Sci., Tohoku Univ., 3. Univ. Tokyo, 4. CRIEPI, 5. CERES)

11:25 AM - 11:40 AM JST | 2:25 AM - 2:40 AM UTC

[S1-06] MD simulation of silicate melt and glass under tension deformation

*Ryota Mamizuka¹, Satoshi Okumura¹, Hiroshi Sakuma² (1. Tohoku Univ. Sci., 2. NIMS)

11:40 AM - 11:55 AM JST | 2:40 AM - 2:55 AM UTC

[S1-07] The magma plumbing system of Izu-Oshima Volcano: Constraints from the H₂O-saturated plagioclase liquidus*Hidemi ISHIBASHI¹ (1. Shizuoka University)

Oral presentation | R1: Characterization and description of minerals (Joint Session with The Gemmological Society of Japan)

10:00 AM - 12:00 PM JST | 1:00 AM - 3:00 AM UTC | ES024 Higashiyama Campus

R1: Characterization and description of minerals (Joint Session with The Gemmological Society of Japan)

Chairperson: Masanori Kurosawa, Hiroshi Kitawaki

10:00 AM - 10:20 AM JST | 1:00 AM - 1:20 AM UTC

[R1-01] Determination of analytical conditions in quantitative electron probe microanalysis of ultra-trace elements: probe current and sampling time

「招待講演」

*Takenori KATO¹ (1. ISEE, Nagoya University)

10:20 AM - 10:35 AM JST | 1:20 AM - 1:35 AM UTC

[R1-02] Mössbauer Spectra and Intensity Tensors of Quadrupole Doublets for Fe³⁺ and Fe²⁺ of Bridgmanite

*Keiji SHINODA¹, Yasuhiro Kobayashi², Takuo Okuchi² (1. Osaka Met. Univ. Sci, 2. KURNS)

10:35 AM - 10:50 AM JST | 1:35 AM - 1:50 AM UTC

[R1-03] Clay mineral characterization by Optical Photothermal Induced Resonance (O-PTIR)

「発表賞エントリー」

*Taro Kido¹, Yohey Suzuki¹ (1. The University of Tokyo)

10:50 AM - 11:05 AM JST | 1:50 AM - 2:05 AM UTC

[R1-04] Characterization and Identification of Synthetic Colored Diamonds for Jewelry Use

*Hiroshi KITAWAKI¹, Kentaro Emori¹, Mio Hisanaga¹, Masahiro Yamamoto¹ (1. Central Gem Laboratory)

11:05 AM - 11:15 AM JST | 2:05 AM - 2:15 AM UTC

[1Lecture-201-07-5add] 休憩

11:15 AM - 11:30 AM JST | 2:15 AM - 2:30 AM UTC

[R1-05] Blue Sapphire from Australia and its origin

*Kentaro Emori¹, Hiroshi Kitawaki¹ (1. Central Gem Laboratory)

11:30 AM - 11:45 AM JST | 2:30 AM - 2:45 AM UTC

[R1-06] Emerald origin characteristics and problems in origin determination

*Zhenghao Zhao¹, Hiroshi Kitawaki¹, Kentaro Emori¹ (1. Central Gem Laboratory)

11:45 AM - 12:00 PM JST | 2:45 AM - 3:00 AM UTC

[R1-07] Gem zircon and sapphire age dating and application of origin determination; A study from New England sapphire fields, New South Wales, Australia

*Ahmadjan ABDURIYIM¹ (1. Tokyo Gem Science LLC)

Oral presentation R3: High-pressure science and deep Earth's material

10:00 AM - 12:00 PM JST | 1:00 AM - 3:00 AM UTC | ES025 Higashiyama Campus

R3: High-pressure science and deep Earth's material

Chairperson: Takeshi Sakai (Ehime University), Ryosuke Sinmyo (Meiji University), Takayuki Ishii (Okayama University), Takaaki Kawazoe (Hiroshima University)

10:00 AM - 10:15 AM JST | 1:00 AM - 1:15 AM UTC

[R3-01] Pressure-induced polyamorphic transition in CaAl_2O_4 glass revealed by elastic wave velocity and X-ray diffraction measurements and molecular dynamics simulations

「招待講演」

*Itaru Ohira¹, Yoshio Kono^{2,3}, Steeve Gréaux³, James W E Drewitt⁴, Sandro Jahn⁵, Fumiya Noritake⁶, Koji Ohara^{7,8}, Satoshi Hiroi^{7,8}, Nozomi M Kondo⁹, Rostislav Hrubia¹⁰, Yuji Higo⁸, Noriyoshi Tsujino⁸, Sho Kakizawa⁸, Kiyofumi Nitta⁸, Oki Sekizawa⁸ (1. Gakushuin Univ. Sci., 2. Kwansai Gakuin Univ. Sci., 3. Ehime Univ. GRC, 4. U. Bristol Phys., 5. U. Cologne IGM, 6. U. Yamanashi Interdisciplinary Research, 7. Shimane Univ. Materials for Energy, 8. JASRI, 9. Okayama Univ. IPM, 10. ANL HPCAT)

10:15 AM - 10:30 AM JST | 1:15 AM - 1:30 AM UTC

[R3-02] Structural analysis of CO_2 -bearing sodium silicate melts using molecular dynamics simulations and synchrotron radiation X-ray diffraction experiments

「発表賞エントリー」

*Shino HAYAFUNE¹, Yohei Onodera², Shinji Kohara², Haruki Ichikawa³, Tatsuya Sakamaki¹, Kenichi Funakoshi⁴, Akio Suzuki¹ (1. Tohoku Univ. Graduate School of Science., 2. NIMS, 3. Tohoku Univ. School of Science., 4. CROSS)

10:30 AM - 10:45 AM JST | 1:30 AM - 1:45 AM UTC

[R3-03] Possible presence of poirierite, a high-pressure phase of $(\text{Mg,Fe})_2\text{SiO}_4$, in the deep Earth's mantle

*Naotaka TOMIOKA¹, Takuo Okuchi², Narangoo Purevjav³, Masaaki Miyahara⁴ (1. KOCHI, JAMSTEC, 2. KRUNS, Kyoto Univ., 3. Seoul National Univ., 4. Earth Planet Sci., Hiroshima Univ.)

10:45 AM - 11:00 AM JST | 1:45 AM - 2:00 AM UTC

[R3-04] Temperature dependence of water content in wadsleyite and ringwoodite

*Toru INOUE^{1,2}, Yuji HAMADA², Hanae KAYA², Yusuke EGI¹, Daichi MAEDA¹, Kazutaka YAMAGUCHI¹, Konosuke YAMADA¹, Takaaki KAWAZOE^{1,2} (1. Hiroshima Univ. Adv. Sci. Eng., 2. Hiroshima Univ. Sci.)

11:00 AM - 11:15 AM JST | 2:00 AM - 2:15 AM UTC

[R3-05] Temperature dependence of the $\text{Fe}^{3+}/\Sigma\text{Fe}$ ratio in wadsleyite by electron energy loss spectroscopy (EELS)

「発表賞エントリー」

*Kazutaka YAMAGUCHI¹, Takaaki Kawazoe¹, Toru Inoue¹, Naotaka Tomioka² (1. Hiroshima University, 2. JAMSTEC, Kochi institute)

11:15 AM - 11:30 AM JST | 2:15 AM - 2:30 AM UTC

[R3-06] Advantage on the analysis of cation site occupancies by neutron diffraction and some examples of application for high-pressure minerals

*Takuo OKUCHI¹ (1. Kyoto Univ.)

11:30 AM - 11:45 AM JST | 2:30 AM - 2:45 AM UTC

[R3-07] In situ lattice volume observation of davemaite in the $\text{CaSiO}_3\text{-H}_2\text{O}$ system up to uppermost lower mantle conditions

*Goru Takaichi¹, Takayuki Ishii², Yu Nishihara¹, Kyoko Matsukage³, Yuji Higo⁴, Noiyoshi Tsujino⁴, Sho Kakizawa⁴ (1. GRC, Ehime Univ., 2. IPM, Okayama Univ., 3. Teikyo University of Science, 4. JASRI)

11:45 AM - 12:00 PM JST | 2:45 AM - 3:00 AM UTC

[R3-08] Elasticity Measurements of Hydrated SiO_2 post-stishovite and its implication to the seismic anomalies in the lower mantle

「発表賞エントリー」

*Yuichiro MORI^{1,2}, Motohiko MURAKAMI², Takashi YOSHINO³, Hiroyuki KAGI¹ (1. UTokyo. Sci., 2. ETHZ. D-ERDW., 3. Okayama Univ. IPM)

Oral presentation | R7: Petrology, Mineralogy and Economic geology (Joint Session with Society of Resource Geology)

2:00 PM - 5:30 PM JST | 5:00 AM - 8:30 AM UTC | ES Hall Higashiyama Campus

R7: Petrology, Mineralogy and Economic geology (Joint Session with Society of Resource Geology)

Chairperson: Norikatsu Akizawa (Atmosphere and Ocean Research Institute, University of Tokyo), Takuya Echigo (Akita University)

2:00 PM - 2:25 PM JST | 5:00 AM - 5:25 AM UTC

[R7-01] The Frontier of Ilmenite Geochronology Using the Uranium-Lead Decay System

「招待講演」

*Kengo Ito¹, Sota Niki², Tsuyoshi Iizuka³, Takafumi Hirata¹ (1. The Univ. of Tokyo. GcRC, 2. Nagoya Univ. ISEE, 3. The Univ. of Tokyo. Sci. EPS)

2:25 PM - 2:40 PM JST | 5:25 AM - 5:40 AM UTC

[R7-02] Distribution of rhenium in hydrothermal deposits in Japan, based on LA-ICP-MS analysis

「発表賞エントリー」

*Mitsuki Ogasawara¹, Junichiro Ohta^{1,2}, Kazutaka Yasukawa¹, Kentaro Nakamura^{2,1}, Yasuhiro Kato^{1,2} (1. Univ. Tokyo. Eng, 2. CIT. ORCeNG)

2:40 PM - 2:55 PM JST | 5:40 AM - 5:55 AM UTC

[R7-03] Occurrences of the PGE bearing (Fe,Ni,Co)AsS solid solution in the Bushveld Complex, South Africa

「発表賞エントリー」

*Sakuya Kubota¹, Takuya Echigo¹, Yasushi Watanabe¹ (1. Akita Univ. Irs.)

2:55 PM - 3:10 PM JST | 5:55 AM - 6:10 AM UTC

[R7-04] Shiranuiite and placer deposit of platinum group minerals, Haraigawa, Misato machi, Kumamoto Prefecture, Japan

*Daisuke HAMANE¹, Takahiro Tanaka, Tadashi Shinmachi (1. The University of Tokyo)

3:10 PM - 3:25 PM JST | 6:10 AM - 6:25 AM UTC

[R7-05] Geochemical and Sr-Nd isotopic characteristics of the Aitutaki peridotite xenoliths from the Cook islands

*Masako YOSHIKAWA¹, Norikatsu Akizawa² (1. Hiroshima Univ., 2. Tokyo Univ. AORI)

3:25 PM - 3:40 PM JST | 6:25 AM - 6:40 AM UTC

[R7-06] Hydration of the mantle and magma genesis in the Izu-Bonin-Mariana fore-arc

*Rion Yamaoka¹, Norikatsu Akizawa¹, Yuji Ichiyama², ToTomoaki Morishita³, Akihiro Tamura³, Hiroyuki Yamashita⁴, Yumiko Harigane⁵, Yasuhiko Ohara^{6,7,8} (1. University of Tokyo, 2. Chiba Univ., 3. Kanazawa Univ., 4. Kanagawa Prefectural Museum of Natural History, 5. AIST, 6. Japan Coast Guard, Hydrographic and Oceanographic Department, 7. JAMSTEC, 8. Nagoya Univ.)

3:40 PM - 3:50 PM JST | 6:40 AM - 6:50 AM UTC

[1Lecture-108-19-7add] 休憩

3:50 PM - 4:15 PM JST | 6:50 AM - 7:15 AM UTC

[R7-07] Fault-rock mineralogy and weakening mechanisms at deep- to shallow-crustal depths of Main Himalayan Thrust, NW India

「招待講演」

*Dyuti Prakash SARKAR¹, Takehiro Hirose¹, Jun-ichi Ando², Kaushik Das², Gautam Ghosh³ (1. Japan Agency for Marine-Earth Science and Technology (JAMSTEC), 2. Hiroshima University, 3. Presidency University)

4:15 PM - 4:30 PM JST | 7:15 AM - 7:30 AM UTC

[R7-08] On the Relationship between Phosphorus Concentration and Microstructure in Goethite in Australian Iron Ore

*Takashi MIKOUCHI¹, Reiko Murao², Toru Takayama² (1. Univ. Museum, Univ. of Tokyo, 2. Nippon Steel)

4:30 PM - 4:45 PM JST | 7:30 AM - 7:45 AM UTC

[R7-09] Mineral zoning and formation temperatures around the Myokenzan lithium pegmatites

「発表賞エントリー」

*Takumi Ishizaki¹, Takuya Echigo¹, Yasushi Watanabe¹ (1. Akita Univ.)

4:45 PM - 5:00 PM JST | 7:45 AM - 8:00 AM UTC

[R7-10] Copper as a Coloring Factor of Pale Blue Sepiolite in amygdales in Higashimatsuura basalt from Kabeshima, Saga Prefecture, SW Japan

「発表賞エントリー」

*Satsuki KIO¹, Terumi Ezima², Yoshiaki Kon³, Hikari Minamisawa⁴, Masaomi Horita⁴ (1. Shinshu Univ. Sci., 2. Shinshu Univ. Sci., 3. GSJ, AIST, 4. Shinshu Univ. Eng.)

5:00 PM - 5:15 PM JST | 8:00 AM - 8:15 AM UTC

[R7-11] Quantification of Mineral Modal and Zoning Ratios by Powder X-ray Diffraction: Advancing Towards Big Data Era

「発表賞エントリー」

*Satoshi MATSUNO¹, Shuhou Maitani², Kenta Yoshida³, Nobuo Hirano¹, Ryosuke Sinmyo², Keiichi Osaka⁴, Shin-ichi Yamasaki¹, Masaaki Uno¹, Atsushi Okamoto¹ (1. Tohoku University, 2. Meiji University, 3. JAMSTEC, 4. Japan Synchrotron Radiation Research Institute)

5:15 PM - 5:30 PM JST | 8:15 AM - 8:30 AM UTC

[R7-12] Searching for Dark Matter Using Geological Samples: Techniques and Preliminary Results

*Natsue ABE^{1,2}, Shigenobu Hirose¹, Qing CHANG¹, Takeshi Hanyu¹, Noriko Hasebe², Yasushi Hoshino³, Takashi Kamiyama⁴, Yoji Kawamura¹, Kohta Murase⁵, Tatsuhiko Naka⁶, Kenji Oguni¹, Katsuhiko Suzuki¹, Seiko Yamazaki⁷ (1. JAMSTEC, 2. Kanazawa University, 3. Kanagawa University, 4. Hokkaido University, 5. Penn State University, 6. Toho University, 7. AIST)

Oral presentation | R1: Characterization and description of minerals (Joint Session with The Gemmological Society of Japan)

2:00 PM - 3:15 PM JST | 5:00 AM - 6:15 AM UTC | ES024 Higashiyama Campus

R1: Characterization and description of minerals (Joint Session with The Gemmological Society of Japan)

Chairperson: Koichi Momma, Yohei Shirose

2:00 PM - 2:15 PM JST | 5:00 AM - 5:15 AM UTC

[R1-08] Ultra nano inclusion in blue sapphire from Diego, Madagascar

*Akira MIYAKE¹, Seika Oto¹, Yohei Igami¹, Kentaro Emori² (1. Kyoto University, 2. CGL)

2:15 PM - 2:30 PM JST | 5:15 AM - 5:30 AM UTC

[R1-09] Reexamination of the symmetry of itoigawaite

*Chigako KIYOHARA, Yohei Igami¹, Akira Miyake¹, Norimasa Shimobayashi¹ (1. Kyoto Univ. Sci.)

2:30 PM - 2:45 PM JST | 5:30 AM - 5:45 AM UTC

[R1-10] Zeolite and altered boninite from Miyanojima, Ogasawara, Japan

*Yuki INOUE¹, Mana Yasui², Jun-ichiro Ishibashi³, Hagiya Hiroshi⁴, Kazuhiko Shimada¹, Athushi Yamazaki² (1. Kyushu Univ. Sci., 2. Waseda Univ., 3. Kobe Univ., 4. Tokyo City Univ.)

2:45 PM - 3:00 PM JST | 5:45 AM - 6:00 AM UTC

[R1-11] Willemseite from the Hamayokawa mine, Nagano Prefecture, Japan

*Satoshi Matsubara¹, Koichi Momma¹, Norio Yanagisawa², Yoko Kusaba¹, Akiko Tokumoto¹, Ritsuro Miyawaki¹ (1. National Museum of Nature and Science, 2. Geological Survey of Japan, AIST)

3:00 PM - 3:15 PM JST | 6:00 AM - 6:15 AM UTC

[R1-12] On the Fe²⁺-analogue of zemannite from Kawazu mine, Shizuoka Prefecture, Japan

*Koichi MOMMA¹, Owen Missen², Stuart Mills³, Ritsuro Miyawaki¹, Satoshi Matsubara¹, Eiji Ohtani⁴, Seiji Kamada⁵, Shin Ozawa⁴ (1. National Museum of Nature and Science, 2. University of Tasmania, 3. The Arkenstone, 4. Tohoku Univ., 5. AD Science Inc.)

Oral presentation | R3: High-pressure science and deep Earth's material

2:00 PM - 3:15 PM JST | 5:00 AM - 6:15 AM UTC | ES025 Higashiyama Campus

R3: High-pressure science and deep Earth's material

Chairperson: Takeshi Sakai (Ehime University), Ryosuke Sinmyo (Meiji University), Takayuki Ishii (Okayama University), Takaaki Kawazoe (Hiroshima University)

2:00 PM - 2:15 PM JST | 5:00 AM - 5:15 AM UTC

[R3-09] Electrical conductivity of the lower mantle materials under high pressure with implication for valence and spin states of iron

「招待講演」

*Izumi MASHINO¹ (1. Okayama Univ.)

2:15 PM - 2:30 PM JST | 5:15 AM - 5:30 AM UTC

[R3-10] GHz-DAC ultrasonics to measure elastic anomaly with the spin transition of ferrous ion in ferropericlae

*Akira Yoneda¹, Izumi Mashino², Ryosuke Matsui¹, Ryoma Ishida¹, Tadashi Kondo¹ (1. Osaka Univ. Sci, 2. Okayama Univ. IPM)

2:30 PM - 2:45 PM JST | 5:30 AM - 5:45 AM UTC

[R3-11] Reaction between water and iron in the early magma ocean and the present core-mantle boundary

*Yongjae Lee¹ (1. Yonsei University)

2:45 PM - 3:00 PM JST | 5:45 AM - 6:00 AM UTC

[R3-12] Sound velocity of B2-FeNiSi alloy at high pressure and high temperature and constitution of the Earth's inner core

*Eiji OHTANI¹, Daijo IKUTA², Hiroshi FUKUI^{3,4}, Tatsuya SAKAMAKI¹, Daisuke Ishikawa^{3,4}, Alfred Q. R. BARON^{3,4} (1. Tohoku University, 2. Institute for Planetary Materials, Okayama University, 3. JASRI, 4. RIKEN)

3:00 PM - 3:15 PM JST | 6:00 AM - 6:15 AM UTC

[R3-13] On the phase boundary between FeS IV and V

*Satoru URAKAWA¹ (1. Okayama University)

Oral presentation | R8: Metamorphic rocks and tectonics

3:30 PM - 6:00 PM JST | 6:30 AM - 9:00 AM UTC | ES024 Higashiyama Campus

R8: Metamorphic rocks and tectonics

Chairperson: Yui Kouketsu (Nagoya University), Shunsuke Endo

3:30 PM - 3:45 PM JST | 6:30 AM - 6:45 AM UTC

[R8-01] Toward the establishment of "EBSD method for the detection of quartz with phase transition to coesite": Incorporation of the Gongen area, Sanbagawa metamorphic belt, Shikoku, SW-Japan

「発表賞エントリー」

*Momoko Minowa¹, Takayoshi Nagaya¹, Taisuke Ito², Simon Wallis² (1. Tokyo Gakugei University, 2. The University of Tokyo)

3:45 PM - 4:00 PM JST | 6:45 AM - 7:00 AM UTC

[R8-02] Pressure-Temperature-Time Conditions of The Basement Rocks in The Banggai-Sula Microcontinent, Indonesia

「発表賞エントリー」

*Fransiska Ayuni Catur Wahyuandari¹, Nobuhiko Nakano¹, Tatsuro Adachi¹, Nugroho Imam Setiawan² (1. Kyushu Univ., 2. Gadjah Mada Univ.)

4:00 PM - 4:15 PM JST | 7:00 AM - 7:15 AM UTC

[R8-03] Amphibolites from the Mineoka belt revisited

*Yuji ICHIYAMA¹, Shun Takamizawa¹, Hisatoshi Ito², Akihiro Tamura³, Tomoaki Morishita³ (1. Chiba University, 2. Central Research Institute of Electric Power Industry, 3. Kanazawa University)

4:15 PM - 4:30 PM JST | 7:15 AM - 7:30 AM UTC

[R8-04] Compositional heterogeneity of garnet in garnet-biotite felsic granulite, Czech Republic

*Daisuke NAKAMURA¹, Yuuka FUJIMURA¹, Yuuki KODERA¹ (1. Okayama University)

4:30 PM - 4:45 PM JST | 7:30 AM - 7:45 AM UTC

[R8-05] Formation and alteration of a zoned calcsilicate vein from the contact aureole of Kasuga area, central Japan: insights from chemical ages of U-Th oxides

*Shunsuke ENDO¹ (1. Shimane University)

4:45 PM - 5:00 PM JST | 7:45 AM - 8:00 AM UTC

[R8-06] Observations of geological structures and microstructures associated with strain release at a continental plate boundary fault

「発表賞エントリー」

*Haruki Yoshiasa¹, Jun-ichi Ando^{1,2}, Kaushik Das^{1,2}, Dyuti Prakash Sarkar³ (1. Hiroshima University, 2. HiPer, 3. JAMSTEC)

5:00 PM - 5:15 PM JST | 8:00 AM - 8:15 AM UTC

[R8-07] Comparison and verification study of new sample forms for use in Raman carbonaceous material geothermometer

「発表賞エントリー」

*Shunsuke Ogino¹, Yui Kouketsu¹, Satoshi Takahashi¹ (1. Nagoya Univ. Env.)

5:15 PM - 5:30 PM JST | 8:15 AM - 8:30 AM UTC

[R8-08] Pressure and temperature conditions for the occurrence of index minerals in pelitic schists in the Sanbagawa belt, Kanto Mountains, Japan, inferred from Raman carbonaceous material geothermometry

*Yui KOUKETSU¹, Ichiko Shimizu² (1. Nagoya Univ. Env., 2. Kyoto Univ. Sci.)

5:30 PM - 5:45 PM JST | 8:30 AM - 8:45 AM UTC

[R8-09] Microstructural and petrological characteristics of the Ryoke mylonite occurred along the Median Tectonic Line in the Shinshiro area, Aichi Prefecture

*Miharu Niwa^{1,2}, Katsuyoshi Michibayashi^{1,3}, Kenichiro Tani⁴, Takuma Nishimura⁵ (1. Nagoya Univ. Env., 2. TMNH, 3. JAMSTEC, 4. NMNS, 5. HMNH)

5:45 PM - 6:00 PM JST | 8:45 AM - 9:00 AM UTC

[R8-10] Re-proposal of the metamorphic zone division in the Horokanai area of the Kamuikotan Belt based on the mineral paragenesis and phengite K-Ar Age.

*Takao HIRAJIMA¹, Kousuke Naemura², Kenta Yoshida³ (1. Kyoto University, 2. Iwate University, 3. JAMSTEC)

Oral presentation | S3: Rheology and Material Transfer in Mantle and Crust (Special Session)

3:30 PM - 6:00 PM JST | 6:30 AM - 9:00 AM UTC | ES025 Higashiyama Campus

S3: Rheology and Material Transfer in Mantle and Crust (Special Session)

Chairperson: Ikuo Katayama (Hiroshima University), Katsuyoshi Michibayashi (Nagoya University)

3:30 PM - 3:50 PM JST | 6:30 AM - 6:50 AM UTC

[S3-01] High-speed time-resolved in-situ stress-strain measurements under high pressure and high temperature using synchrotron radiation X-rays

「招待講演」

*Noriyosi TSUJINO¹ (1. JASRI)

3:50 PM - 4:05 PM JST | 6:50 AM - 7:05 AM UTC

[S3-02] Deformation-induced crystallographic-preferred orientation of ϵ -FeOOH

*Yu NISHIHARA¹, Yui MORI¹, Wentian WU¹, Noriyoshi TSUJINO² (1. GRC, Ehime Univ., 2. JASRI)

4:05 PM - 4:20 PM JST | 7:05 AM - 7:20 AM UTC

[S3-03] Strength of metastable olivine and the rheology of subducting cold slab at mantle transition zone

「発表賞エントリー」

*Rikuto HONDA¹, Tomoaki Kubo¹, Noriyoshi Tsujino², Yuji Higo², Sho Kakizawa², Yuki Shibazaki³, Yu Nishihara⁴ (1. Kyushu University, 2. JASRI, 3. KEK, 4. Ehime Univ. GRC)

4:20 PM - 4:40 PM JST | 7:20 AM - 7:40 AM UTC

[S3-04] High-resolution measurement of ice anelasticity over a broad frequency range with a new cryogenic forced oscillation apparatus

「招待講演」

*Hatsuki Yamauchi¹, Christine McCarthy¹, Benjamin Holtzman¹, David Goldsby², Travis Hager² (1. Columbia Univ. LDEO, 2. U. Penn)

4:40 PM - 4:55 PM JST | 7:40 AM - 7:55 AM UTC

[S3-05] Relationship between creep and grain growth in rock with bicontinuous structure

「発表賞エントリー」

*Shenghao Jiang¹, Takehiko Hiraga¹ (1. The University of Tokyo)

4:55 PM - 5:10 PM JST | 7:55 AM - 8:10 AM UTC

[S3-06] Microstructures of plagioclase in gabbroic ultramylonites and quadruple point analysis

*Itsuki Natsume¹, Katsuyoshi Michibayashi^{2,3}, Yohei Igami⁴ (1. Kanagawa Prefectural Museum of Natural History, 2. Nagoya Univ., 3. JAMSTEC, 4. Kyoto Univ.)

5:10 PM - 5:25 PM JST | 8:10 AM - 8:25 AM UTC

[S3-07] Reconstruction of the uppermost mantle continuous structure of the Oman ophiolite

*Takeo Okuwaki¹, Natsume Itsuki², Katsuyoshi Michibayashi¹ (1. Nagoya Univ. Env, 2. Kanagawa Pref. Mus. Nat. Hist.)

5:25 PM - 5:40 PM JST | 8:25 AM - 8:40 AM UTC

[S3-08] Deformation and melt-rock interaction in the Horoman peridotite: Petrological and structural study of the MHL suite and BDH suite rocks

「発表賞エントリー」

*Aya Hihara¹, Miki Tasaka¹, Keisuke Kurihara¹, Hajime Taniuchi², Tastuhiko Kawamoto¹ (1. Shizuoka Univ., 2. AIST)

5:40 PM - 5:55 PM JST | 8:40 AM - 8:55 AM UTC

[S3-09] Multicomponent measurements of seismic velocity and electrical resistivity using foliated serpentinite and peridotite

「発表賞エントリー」

*Tomohiro Ito¹, Ikuo Katayama¹, Katsuyoshi Michibayashi², Kazuki Matsuyama² (1. Hiroshima University, 2. Nagoya University)

Poster presentation | S1: Dynamics of igneous processes (Special Session)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC | Entrance Hall Higashiyama Campus

S1: Dynamics of igneous processes (Special Session)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S1-P-01] Oxidation states of HIMU-type ocean island basalts: Insights from μ -XANES analysis of quenched glasses and melt inclusions

*Yuuki HAGIWARA¹, Hidemi Ishibashi², Takeshi Hanyu¹ (1. Japan Agency for Marine-Earth Science and Technology, 2. Shizuoka Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S1-P-02] The composition and pressure of the fluid in crystal mush estimated from cordierite in tonalitic polycrystalline volcanic ejecta

*Shumpei YOSHIMURA¹ (1. Hokkaido University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S1-P-03] Geochemical evolution of Izu-Oshima volcano: Constraints from analysis of melt inclusions in a submarine core

*Morihisa HAMADA¹, Erika TANAKA², Takeshi HANYU¹, Kenji SHIMIZU³, Takayuki USHIKUBO², Qing CHANG¹, Yoshihiko TAMURA¹ (1. IMG, JAMSTEC, 2. Marine Core Research Institute, Kochi Univ., 3. Kochi Institute for Core Sample Research, JAMSTEC)

Poster presentation | S3: Rheology and Material Transfer in Mantle and Crust (Special Session)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC | Entrance Hall Higashiyama Campus

S3: Rheology and Material Transfer in Mantle and Crust (Special Session)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S3-P-01] Preliminary results of deformation experiments on hydrous stishovite using a rotational DAC

*Shintaro AZUMA¹, Keishi Okazaki², Kentaro Uesugi³, Masahiro Yasutake³, Steeve Gréaux⁴, Yoshiyuki Okuda^{1,5}, Bunrin Natsui¹, Eranga Jayawickrama², Kenji Ohta¹ (1. Tokyo Tech., 2. Hiroshima Univ., 3. JASRI, 4. Ehime Univ., 5. University of Hawai'i)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S3-P-02] Water weakening of Mg₂SiO₄ ringwoodite

「発表賞エントリー」

*Yuta Goto¹, Tomoaki Kubo¹, Rikuto Honda¹, Yuki Shibazaki² (1. Kyushu Univ., 2. KEK-PF)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S3-P-03] Toward an understanding of dehydration process of partially serpentinized slab peridotite under conditions where deep earthquakes occur

*Tomoaki KUBO¹, Musashi Ezaki¹, Nobumasa Fujiwara¹, Rikuto Honda¹, Goto Yuta¹, Noriyoshi Tsujino² (1. Kyushu University, 2. JASRI)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S3-P-04] Viscous anisotropy of olivine aggregates using micro Vickers indentation tests

「発表賞エントリー」

*Namu Fujii¹, Miki Tasaka¹ (1. Shizuoka University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S3-P-05] Crystal-fabric analysis using principal component analysis method for the Horoman peridotite

「発表賞エントリー」

*Kazuki Matsuyama¹, Katsuyoshi Michibayashi¹ (1. Nagoya Univ. Env.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S3-P-06] Traces of mantle fluid/melt within olivine phenocrysts from Ohima-Oshima picritic basalts

*Ryo Tsukawaki¹, Terumi Ejima², Atusi Ninomiya³, Shoji Arai⁴ (1. Shinshu Univ. Sci., 2. Shinshu Univ. Sci., 3. Sumiko Res. Exp. & Dev. Co., 4. Kanazawa Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S3-P-07] Microstructural characteristics of ultramafic rocks in the Tosa Megamullion, the Shikoku Basin.

「発表賞エントリー」

*So Inoue¹, Katsuyoshi Michibayashi^{1,2}, Yumiko Harigane³, Yasuhiko Ohara^{1,2,4} (1. GSES, Nagoya Univ., 2. JAMSTEC, 3. GSJ/AIST, 4. JCG)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S3-P-08] Deformation microstructures of granitic mylonite in Hida Metamorphic Belt

「発表賞エントリー」

*Masaaki Horie¹, Katsuyoshi Michibayashi¹ (1. GSES, Nagoya Univ.)

Poster presentation | R1: Characterization and description of minerals (Joint Session with The Gemmological Society of Japan)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC | Entrance Hall Higashiyama Campus

R1: Characterization and description of minerals (Joint Session with The Gemmological Society of Japan)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-01] Chemical quantitative analysis of heulandite using SEM-EDS - How can we accurately estimate the chemical composition of zeolite?

「発表賞エントリー」

*Atsushi ISHIHARA¹, Hiroaki Ohfujii¹ (1. Tohoku university)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-02] Quantitative electron microprobe analysis of xenotime

*Yasuyuki BANNO¹ (1. AIST)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-03] Deformation and compositional changes of plagioclase: A combined electron backscattered diffraction and energy dispersive X-ray spectroscopy approach

「発表賞エントリー」

*Kohei Nimura¹, Katsuyoshi Michibayashi^{1,2} (1. Nagoya University, 2. JAMSTEC)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-04] Mineralogical features of smelting slags from the Early Iron Age Yashin Tepe site, northeastern Iraq

*Masanori KUROSAWA¹, Shin'ichi Nishiyama² (1. Univ. Tsukuba, 2. Chubu Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-05] The origin of abundant graphite in quartz veins in Ishidera area, Wazuka Town, Kyoto Prefecture, Japan

*Masaki Nishio¹, Itaru Mitsukawa¹, Yohei Igami¹, Akira Miyake¹, Norimasa Shimobayashi¹ (1. Kyoto Univ. Sci.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-06] Michitoshiite-(Cu), a new Ge-containing platinum group mineral from Haraigawa, Misato machi, Kumamoto Prefecture, Japan

*Takahiro TANAKA¹, Daisuke Nishio Hamane², Tadashi Shinmachi (1. Nittetsu Mining Co., Ltd., 2. ISSP, Univ. of Tokyo)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-07] Fibrous inclusions in rose quartz

*Yohei SHIROSE¹, Hayato Fudamoto¹, Sayako Inoue² (1. Ehime Univ. Sci., 2. Ehime Univ. GRC)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-08] Rose quartz in gneisses from Uoshima Island, Ehime Prefecture

*Yohei SHIROSE¹, Shoma Sakai¹ (1. Ehime Univ. Sci.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-09] Secondary arsenate minerals from the Takumi Mine, Hyogo Prefecture, Japan

*Yohei SHIROSE¹, Riakako Kamise¹, Katsuichi Nishida, Yoshiteru Fujiwara (1. Ehime Univ. Sci.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-10] Mineralogical properties of lautenthalite and wroewolfeite from the Nii mine, Hyogo Prefecture, Japan

*Masayuki Ohnishi, Norimasa Shimobayashi¹, Daisuke Nishio-Hamane², Keiji Shinoda³, Takeshi Hisano (1. Sci., Kyoto Univ., 2. ISSP, Univ. of Tokyo, 3. Sci., Osaka Metro. Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-11] A re-examination of Sr-rich apatite from Itoigawa, Niigata Prefecture, Japan

*Seiichiro UEHARA¹, Koichi MONMA², Masayuki OHNISHI, Shunsuke OHSUMI, Yoshiya OHKI, Hiroki OKA³ (1. Kyushu Univ. Museum, 2. Nat'l. Mus. Nat. Sci., 3. OYO Corp.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-12] Hydroxylchondrodite from Ogouchi, Hinokage, Nisiusuki, Miyazaki Prefecture, Japan

*Toshiro Okada¹, Seiichiro Uehara², Isao Yukinori³, Yohei Shirose⁴ (1. Kashii 2 JHS, 2. Kyushu Univ, 3. Fukuoka Stc, 4. Ehime Univ)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-13] Arsenmedaite from the Yamato mine, Kagoshima Prefecture, SW Japan

*Shunsuke Ohsumi, Daisuke Nishio-Hamane¹, Hiroki Oka², Masashi Tamura³, Kosuke Takagi⁴ (1. ISSP, Univ. of Tokyo, 2. OYO Corp., 3. Fac. Eng. Tech. Div., Mie Univ., 4. Grad. Sch. of Eng., Mie Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-14] "Common Hornblende" from Mt. Tawarayama (Goou-toge), the outer-rim of Mt. Aso, Kumamoto Prefecture

*Haruki Inoue¹, Seiichiro Uehara² (1. Enecom Co., Ltd., 2. Kyushu Univ. Museum)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-15] Chemical composition of tourmaline and amphibole associated with gabbro from Kajishima, Ehime Prefecture, Japan

「発表賞エントリー」

*Itsuki Ota¹, Kazuya Shimooka², satoshi saitou¹, youhei shirose¹ (1. Ehime Univ. Sci and Eng, 2. Kwansei Gakuin Univ. Sci)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-16] Constituent minerals of clay associated with the pegmatite dike in Nagatare, Fukuoka Prefecture, Japan

*Yuya TAKEDA¹, Seiichiro Uehara², Yoshihiro Kuwahara³ (1. Kyushu Univ. ISGS, 2. Kyushu Univ. Museum, 3. Kyushu Univ. SCS)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-17] Microstructure of serpentine veins in peridotite in Ooshika Village, Nagano Prefecture, Japan

*Yuya TAKEDA¹, Yoshihiro Kuwahara³, Seiichiro Uehara² (1. Kyushu Univ. ISGS, 2. Kyushu Univ. Museum, 3. Kyushu Univ. SCS)

Poster presentation | R3: High-pressure science and deep Earth's material

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC | Entrance Hall Higashiyama Campus

R3: High-pressure science and deep Earth's material

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R3-P-01] Influence of co-existing inorganic minerals on chemical reaction of *n*-alkane under high-pressure and high-temperature conditions of subduction zone.*Ayako SHINOZAKI¹, Kina Takimoto¹, Takaya Nagai¹, Koichi Mimura² (1. Hokkaido University, 2. Nagoya University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R3-P-02] Differential Scanning Calorimetry of Mn₂SiO₄ tephroite*Yuta Asami¹, Itaru Ohira², Hiroshi Kojitani² (1. Gakushuin Univ. Sci, 2. Gakushuin Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R3-P-03] Ab initio calculation of the polarized IR spectra and hydrogen positions of hydrous Bridgmanite

*Kikuyo Inagaki¹, Jun Tsuchiya¹, Yanyao Zhang³, Jung-Fu Lin², Shun-ichiro Karato⁴, Jennifer Kung⁵, ChingChien Li⁵ (1. GRC Ehime Univ., 2. Univ. Texas Austin, 3. Stanford Univ., 4. Yale Univ., 5. National Cheng Kung Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R3-P-04] Determination of the stability of silica phases under high pressure by ultra-fast X-ray diffraction measurements

*Ryosuke SINMYO¹, Saori Kawaguchi-Imada², Takayuki Ishii³, Hiroshi Sakuma⁴, Ayase Ogawa¹, Kenta Kobayashi¹, Shuhou Maitani¹ (1. Meiji Univ. Sci. Tech., 2. JASRI, 3. Okayama Univ. IPM, 4. NIMS)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R3-P-05] Crystallographic preferred orientation properties of Ferropericlasite polycrystals obtained from large strain deformation experiments under lower mantle pressures

「発表賞エントリー」

*Bunrin Natsui¹, Shintaro Azuma¹, Keishi Okazaki^{2,5}, Kentaro Uesugi³, Masahiro Yasutake³, Saori Kawaguchi³, Ryuichi Nomura⁴, Kenji Ohta¹ (1. Tokyo Tech, 2. Hiroshima Univ., 3. JASRI, 4. Kyoto Univ., 5. JAMSTEC)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R3-P-06] Investigation of hydrogen sealing materials at high temperature and high pressure using neutron imaging

*Sho KAKIZAWA¹, Hiroyuki Kagi², Masahiro Takano², Asami Sano-Furukawa³, Takanori Hattori³, Abe Jun⁴, Kenichi Funakoshi⁴ (1. JASRI, 2. UTokyo Sci., 3. JAEA J-PARC Center, 4. CROSS, Neutron Science and Technology Center)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R3-P-07] Reactions of FeS with hydrogen at high pressure and high temperature revisited

「発表賞エントリー」

*Masahiro Takano¹, Hiroyuki Kagi¹, Yuichiro Mori¹, Katsutoshi Aoki¹, Sho Kakizawa², Noriyoshi Tsujino², Yuji Higo², Asami Sano-Furukawa³ (1. UTokyo, 2. JASRI, 3. J-PARC center, JAEA)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R3-P-08] Extreme pressure generation using toroidal diamond anvil cell

*Takeshi SAKAI¹, Yuki Nakamoto², Satoru Nakamura¹, Sotaro Iwatsu², Shuto Fukuda², Yuki Kato², Katsuya Shimizu², Hirokazu Kadobayashi³, Saori Kawaguchi-Imada³ (1. GRC, Ehime University, 2. KYOKUGEN, Osaka University, 3. JASRI)

Fri. Sep 13, 2024

Oral presentation | T1: Comprehensive understanding of the crustal evolution and resource exploration in Asia (Symposium)

9:00 AM - 12:00 PM JST | 12:00 AM - 3:00 AM UTC | ES Hall Higashiyama Campus

T1: Comprehensive understanding of the crustal evolution and resource exploration in Asia (Symposium)

Chairperson: Yasuhito Osanai, Masaaki Owada

9:00 AM - 9:25 AM JST | 12:00 AM - 12:25 AM UTC

[T1-01] Decarbonized Society and Essential Metal Resources

「招待講演」

*YOSHITAKA HOSO¹ (1. JICA)

9:25 AM - 9:50 AM JST | 12:25 AM - 12:50 AM UTC

[T1-02] Japan's current approach to securing mineral resources

「招待講演」

*Kazuhiro YONEMURA¹ (1. JOGMEC)

9:50 AM - 10:15 AM JST | 12:50 AM - 1:15 AM UTC

[T1-03] Critical metal potentiality of Mongolia

「招待講演」

*Sereenen Jargalan¹, M. Arvinzun² (1. Mongolian University of Science and Technology, 2. Mongolian Society of Economic Geologists)

10:15 AM - 10:30 AM JST | 1:15 AM - 1:30 AM UTC

[2Lecture-101-06-4add] 休憩

10:30 AM - 10:55 AM JST | 1:30 AM - 1:55 AM UTC

[T1-04] Geology, sedimentation environment of the Ovoot khural coal bearing depression, in South Mongolia

「招待講演」

*Magsarjav Ochirbat², Sereenen Jargalan¹ (1. Mongolian University of Science and Technology, 2. Mongolian Society of Economic Geologists)

10:55 AM - 11:20 AM JST | 1:55 AM - 2:20 AM UTC

[T1-05] Insights into the mineralogical characteristics of Li-enriched metasomatic albitite from the Iwagi islet, SW Japan

「招待講演」

*Mariko NAGASHIMA¹, Teruyoshi IMAOKA¹ (1. Yamaguchi Univ. Sci.)

11:20 AM - 11:45 AM JST | 2:20 AM - 2:45 AM UTC

[T1-06] Ion adsorption-type REE deposits: the source of HREE

「招待講演」

*Yasushi Watanabe¹ (1. Akita Univ. Int. Res. Sci.)

11:45 AM - 12:00 PM JST | 2:45 AM - 3:00 AM UTC

[2Lecture-101-06-8add] 休憩

Oral presentation | S2: Water Rock Interaction (Special Session)

9:00 AM - 12:00 PM JST | 12:00 AM - 3:00 AM UTC | ES024 Higashiyama Campus

S2: Water Rock Interaction (Special Session)

Chairperson: Noriyoshi Tsuchiya

9:00 AM - 9:05 AM JST | 12:00 AM - 12:05 AM UTC

[2Lecture-201-10-1add] コンビーナ挨拶

9:05 AM - 9:35 AM JST | 12:05 AM - 12:35 AM UTC

[S2-01] Slab-derived fluid infiltrating back-arc mantle

「招待講演」

*Junji YAMAMOTO¹ (1. Kyushu University)

9:35 AM - 9:50 AM JST | 12:35 AM - 12:50 AM UTC

[S2-02] Geochemical diversity and significance of orthopyroxene pseudomorphs in ultramafic rocks derived from mantle wedges

「発表賞エントリー」

*Takumi Wani¹, Yuji Ichiyama¹, Akihiro Tamura², Tomoaki Morishita² (1. Chiba University, 2. Kanazawa University)

9:50 AM - 10:05 AM JST | 12:50 AM - 1:05 AM UTC

[S2-03] Petrogenesis and significance of ophicarbonates in the Kanasaki serpentinite body (Kanto Mountains, Central Japan)

*Ryosuke OYANAGI^{1,2}, Hikaru Sawada^{3,2}, Qing Chang², Madhusoodhan Satish-Kumar⁴ (1. Kokushikan Univ., 2. JAMSTEC, 3. Toyama Univ., 4. Niigata Univ.)

10:05 AM - 10:20 AM JST | 1:05 AM - 1:20 AM UTC

[S2-04] CO₂ mineralization in andesitic rocks revealed by hydrothermal experiments and thermal analyses*Otgonbayar DANDAR¹, Atsushi Okamoto¹, Masaaki Uno¹, Miku Takeya² (1. Tohoku University, 2. INPEX)

10:20 AM - 10:30 AM JST | 1:20 AM - 1:30 AM UTC

[2Lecture-201-10-6add] 休憩

10:30 AM - 10:45 AM JST | 1:30 AM - 1:45 AM UTC

[S2-05] Massive and foliated serpentinites from the Udonohana ultramafic body, Western Ehime Prefecture, Japan.

「発表賞エントリー」

*Yui Joguchi¹, Satomi Enju¹ (1. Ehime Univ. Sci. &Egn.)

10:45 AM - 11:00 AM JST | 1:45 AM - 2:00 AM UTC

[S2-06] "Element transport and magnetite decomposition during alteration of the gabbroic vein in serpentinite body from the Bayankhongor ophiolite, Mongolia"

「発表賞エントリー」

*Nomin Tumurkhuu¹, Otgonbayar Dandar¹, Masaaki Uno¹, Manzshir Bayarbold¹, Atsushi Okamoto¹ (1. Tohoku University)

11:00 AM - 11:15 AM JST | 2:00 AM - 2:15 AM UTC

[S2-07] Dissolution reprecipitation - re-equilibration process of feldspar in heat source granite and supercritical geothermal reservoir using borehole samples from Kakkonda granite

「発表賞エントリー」

*Masayoshi Hoshida¹, Masaaki Uno¹, Satoshi Matsuno¹, Astin Nurdiana¹, Noriyoshi Tsuchiya^{2,1} (1. Tohoku University, 2. National Institute of Technology, Hachinohe College)

11:15 AM - 11:30 AM JST | 2:15 AM - 2:30 AM UTC

[S2-08] Formation mechanism of "cleavable olivine"

*Jun-ichi ANDO^{1,2}, Naotaka Tomioka^{3,2}, Hirokazu Maekawa⁴ (1. Hiroshima Univ., 2. Hiroshima Univ., HiPeR, 3. JAMSTEC, 4. Osaka Metropolitan Univ.)

11:30 AM - 11:45 AM JST | 2:30 AM - 2:45 AM UTC

[S2-09] Experimental study to elucidate sulfide chimney development process and power generation characteristics in submarine hydrothermal systems

「発表賞エントリー」

*Kentarō Toda¹, Atsushi Okamoto¹, Dandar Otgonbayar¹, Misaki Takahashi¹, Yoshinori Sato¹ (1. Tohoku Univ. Environmental Sci)

11:45 AM - 12:00 PM JST | 2:45 AM - 3:00 AM UTC

[S2-10] Carbonation of Mantle Peridotite: An Approach From Fluid Inclusion Analysis and Hydrothermal Experiments

*Tatsuhiko KAWAMOTO¹ (1. Shizuoka University)

Oral presentation | S3: Rheology and Material Transfer in Mantle and Crust (Special Session)

9:00 AM - 10:00 AM JST | 12:00 AM - 1:00 AM UTC | ES025 Higashiyama Campus

S3: Rheology and Material Transfer in Mantle and Crust (Special Session)

Chairperson: Miki Tasaka (Shizuoka University)

9:00 AM - 9:15 AM JST | 12:00 AM - 12:15 AM UTC

[S3-10] Rheological evolution of olivine during formation of the mantle lithosphere

*Katsuyoshi MICHIBAYASHI^{1,2}, Takeo Okuwaki¹, Itsuki Natsume³ (1. Nagoya University, 2. JAMSTEC, 3. Kanagawa Prefectural Museum of Natural History)

9:15 AM - 9:30 AM JST | 12:15 AM - 12:30 AM UTC

[S3-11] Three-dimensional location analysis on acoustic emissions and faults in olivine under pressure-temperature conditions of subducting slabs

*Tomohiro OHUCHI¹, Masato Hoshino², Kentaro Uesugi², Satoshi Okumura³, Yuji Higo², Noriyoshi Tsujino², Sho Kakizawa² (1. GRC, Ehime Univ., 2. JASRI, 3. Tohoku Univ Sci.)

9:30 AM - 9:45 AM JST | 12:30 AM - 12:45 AM UTC

[S3-12] In-situ observation of grain growth and fluid movement using camphor as a rock analogue

*Junichi Fukuda¹ (1. Dept. Geos. Osaka Metropol. Univ.)

9:45 AM - 10:00 AM JST | 12:45 AM - 1:00 AM UTC

[S3-13] Mantle carbonation through seawater penetration along the outer-rise faults

*Ikuo KATAYAMA¹, Keishi Okazaki¹, Atsushi Okamoto² (1. Hiroshima University, 2. Tohoku University)

Oral presentation | R4: Mineral sciences of the Earth surface

10:15 AM - 12:00 PM JST | 1:15 AM - 3:00 AM UTC | ES025 Higashiyama Campus

R4: Mineral sciences of the Earth surface

Chairperson: Hiroshi Sakuma(NIMS), Satoko Motai(Yamagata Univ.), Jun Kawano(Hokkaido University)

10:15 AM - 10:35 AM JST | 1:15 AM - 1:35 AM UTC

[R4-01] Understanding and application of spherical concretions: A new durable sealing material learnt from nature

「招待講演」

*Hidekazu Yoshida¹ (1. Nagoya University)

10:35 AM - 10:50 AM JST | 1:35 AM - 1:50 AM UTC

[R4-02] Color change process of Hiroshima granite due to weathering

*Tadashi YOKOYAMA¹, Yuka Inkyo, Masahiro Kaibori¹ (1. Hiroshima University)

10:50 AM - 11:05 AM JST | 1:50 AM - 2:05 AM UTC

[R4-03] Crystallographic preferred orientation and grain size of apatite in terrestrial mammalian bones

*Kyoko N. MATSUKAGE¹, Momoka Ide², Masaya Kurata², Yu Nishihara³ (1. Teikyo Univ. of Sci. Natural and Environmental Sci., 2. Teikyo Univ. of Sci. Aminal Sci., 3. Ehime Univ.)

11:05 AM - 11:20 AM JST | 2:05 AM - 2:20 AM UTC

[R4-04] Microscopic distribution of sodium in biogenic aragonite

*Taiga Okumura¹, Michio Suzuki², Alberto Perez-Huerta³, Eshita Samajpati³, Toshihiro Kogure¹ (1. UTokyo Sci., 2. UTokyo Agri. Life Sci., 3. Univ. Alabama Geol. Sci.)

11:20 AM - 11:40 AM JST | 2:20 AM - 2:40 AM UTC

[R4-05] Structural and functional analyses of organic matrices regulating the formation of minerals in biomineralization.

「招待講演」

*Michio Suzuki¹ (1. UTokyo)

11:40 AM - 11:55 AM JST | 2:40 AM - 2:55 AM UTC

[R4-06] Aragonite formation from amorphous calcium carbonate (ACC) with addition of *n*-butylamine*Hiroyuki KAGI¹, Kensuke Muraoka¹ (1. The University of Tokyo)

Poster presentation | T1: Comprehensive understanding of the crustal evolution and resource exploration in Asia (Symposium)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC | Entrance Hall Higashiyama Campus

T1: Comprehensive understanding of the crustal evolution and resource exploration in Asia (Symposium)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[T1-P-01] Syenitic veining at the northern Eastern Ghats Belt, India: Formation mechanism, fluid-rock interaction and a review of its economic mineral potential

*Kaushik DAS^{1,5}, Proloy Ganguly², Aparupa Banerjee³, Sankar Bose^{4,5} (1. Hiroshima University, 2. Kazi Nazrul University, 3. Shahid Matangini Hazra Government General Degree College, 4. Presidency University, 5. HiPeR, Hiroshima University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[T1-P-02] Amphibole-bearing granitic rocks in the Ereendavaa block, NE Mongolia: Insights into multi-stage magmatic sources and crystallization conditions

*Munkhdelger Bold¹, Tatsuki Tsujimori¹, Yasuhito Osanai², Nobuhiko Nakano², Tatsuro Adachi², Otgonbayar Dandar¹, Fransiska Ayuni Catur Wahyuandari² (1. Tohoku Univ., 2. Kyushu Univ)

Poster presentation | S2: Water Rock Interaction (Special Session)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC | Entrance Hall Higashiyama Campus

S2: Water Rock Interaction (Special Session)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S2-P-01] Scales of extensional-shear fracturing and magnitudes of seismicity induced by magma intrusions into lower crust: Scale comparisons of dike swarm in the high-grade metamorphic rocks and deep low-frequency earthquakes

「発表賞エントリー」

*Takumi Nara¹, Masaoki Uno¹, Tetsuo Kawakami², Fumiko Higashino², Tatsuro Adachi³, Noriyoshi Tsuchiya^{1,4} (1. TOHOKU Univ. Env., 2. Kyoto Univ. Sci, 3. Kyushu Univ. Soc., 4. Hachinohe Kosen)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S2-P-02] Carbonation of serpentinite and formation process of listvenite from Urayama River, Shikokuchuo City, Ehime Prefecture, Japan

「発表賞エントリー」

*Hikaru Takagaki¹, Yohei Shirose¹ (1. Ehime Univ. Sci.&Egn.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S2-P-03] Texture transition during serpentinization in Hodono, Ehime Prefecture

「発表賞エントリー」

*Hinano Wada¹, Enju Satomi¹ (1. Ehime Univ. S/E)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S2-P-04] Water-rock interaction recorded in episyenites from Hakatajima Island, Ehime Prefecture

「発表賞エントリー」

*Toko FUKUI¹, Kazuya SHIMOOKA², Toshiro TAKAHASHI³, Satoshi SAITO¹ (1. Ehime Univ., 2. Kwansai Gakuin Univ., 3. Niigata Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S2-P-05] Mechanisms of Reaction-Induced Fracturing in Serpentinite Carbonation; Insights from Hydrothermal Experiments and Geochemical Modeling

「発表賞エントリー」

*Taiki Taiki¹, Masaoki Uno¹, Atsushi Okamoto¹ (1. Tohoku University)

Poster presentation | R4: Mineral sciences of the Earth surface

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC | Entrance Hall Higashiyama Campus

R4: Mineral sciences of the Earth surface

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R4-P-01] Design and synthesis of aragonite particles as a reinforcement of plastic materials

*Hiroshi SAKUMA¹, Shigeru SUEHARA¹, Masumi KAMON¹, Kenji TAMURA¹ (1. NIMS)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R4-P-03] Experiments on the Inhibitory Effect of Polysaccharides on Cation Ordering of Dolomite During Dolomitization Reaction at 200°C: Preliminary Results

Hiromi KONISHI¹, *Yao Chen¹ (1. Niigata Univ. Sci.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R4-P-04] Formation Process of Carbonate Minerals in Non-aqueous Solvents: Consideration of the Effects of Different Hydration States of Cations

Naoki IWANE¹, *Jun KAWANO¹, Hiroyuki KAGI², Ayako SHINOZAKI¹, Takaya NAGAI¹ (1. Hokkaido Univ. Sci., 2. UTokyo Sci.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R4-P-05] Impact of evaporation on CO₂ mineralization during enhanced rock weathering

*Naoki NISHIYAMA¹, Masao SORAI¹, Keisuke FUKUSHI², Yuto NISHIKI¹ (1. National Institute of Advanced Industrial Science and Technology (AIST), 2. Kanazawa University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R4-P-06] Framboidal pyrite in modern stromatolite from Fukiage-Jigoku, Onikobe Spring, Miyagi, Japan

「発表賞エントリー」

*Tatsuya Kamada¹, Hiroaki Ohfuji¹ (1. Tohoku Univ. Sci.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R4-P-07] Observations of nano-texture for amosite asbestos by using high-resolution STEM imaging

*Hayato Miura¹, Ichiro Ohnishi¹ (1. JEOL Ltd.)

Poster presentation | R7: Petrology, Mineralogy and Economic geology (Joint Session with Society of Resource Geology)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC | Entrance Hall Higashiyama Campus

R7: Petrology, Mineralogy and Economic geology (Joint Session with Society of Resource Geology)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R7-P-01] Ezochiite and placer deposit of platinum group minerals in northwestern Hokkaido, Japan

*Daisuke HAMANE¹, Katsuyuki Saito (1. The University of Tokyo)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R7-P-02] Review for Mineralogical Science: Mineral Resources, Heritage Stone, and SDGs

*Yuhei Takahashi¹ (1. NUE)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R7-P-03] Formation process of olivine with remarkable parting and apparently oblique extinction in the Iherzolite of Ochiai-Hokudo peridotite complex, Okayama Prefecture, Japan

*Terumi EJIMA¹, Takashima Chihiro², Arai Shoji³ (1. Shinshu University, 2. DAIYA SEIKI Co., Ltd., 3. Kanazawa University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R7-P-04] The effect of Na ion on carbonation reaction of forsterite

*Hiroki Hasegawa¹, Atsushi Kyono², Satoru Okada¹, Kosuke Yamaguchi¹ (1. Univ of Tsukuba, Grad. sch. of Life and Environmental. Sci, 2. Univ of Tsukuba, Life and Environmental Science)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R7-P-05] Mineralogical characteristics of Pothole Reef and Pseudo Merensky Reef at the western limb of the Bushveld Complex, South Africa

「発表賞エントリー」

*Amu Umesato¹, Takuya Echigo¹, Yasushi Watanabe¹ (1. Akita Univ. Int.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R7-P-06] High-Ti biotite in the alkali volcanic rock from the Akiyoshi Belt and its significance

*Kosuke Kimura¹, Kaushik Das², Yasutaka Hayasaka³ (1. Osaka Metro. Univ. Sci., 2. Hiroshima Univ., 3. Amakusa Mus. Goshoura Dinosaur Isl.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R7-P-07] Estimation of the origin of SDW in the Horoman peridotite complex by analysis of micro-inclusions in the olivine

「発表賞エントリー」

*Masaharu Aketa¹, Akira Miyake¹, Norikatsu Akizawa², Megumi Matsumoto³, Yohei Igami¹, Itaru Mitsukawa¹ (1. Kyoto University, 2. University of Tokyo, 3. Tohoku University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R7-P-08] Fe-Ni-S-As minerals in the Imono peridotite body, Besshi area, Niihama city, Ehime prefecture.

「発表賞エントリー」

*Masato Kuniyoshi¹, Satomi Enju¹ (1. Ehime Univ.Sci and Eng.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R7-P-09] Fluorite mineralization associated with alkaline metasomatism in the Jinmu-Mihara deposit, Hiroshima, Japan.

「発表賞エントリー」

*Masahiro SUNADA¹, Takuya Echigo¹, Yasushi Watanabe¹ (1. Akita Univ. IRS.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R7-P-10] Petrological study of ultramafic rocks from the Kiyama area, eastern Kumamoto City

Narumichi Nishio¹, *Satoko ISHIMARU² (1. Kumamoto Univ. Sci., 2. Kumamoto Univ. FAST)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R7-P-11] FLUID INCLUSION STUDIES IN QUARTZ VEINS WITH TIN MINERALIZATION IN THE KIBARAN INTRUSIVE ROCKS IN KALEHE (SOUTH KIVU, DR CONGO)

「発表賞エントリー」

*MUSA Moise-Kam's SAIDI¹, MADDHUSOODHAN Satish Kumar¹ (1. Niigata Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R7-P-12] Depleted peridotite and melt reaction as recorded by layered dunite-harzburgite in the Horoman peridotite, Hidaka Metamorphic Belt, Hokkaido, Japan.

「発表賞エントリー」

*Keisuke Kurihara¹, Tatsuhiko Kawamoto¹, Aya Hihara¹, Miki Tasaka¹, Hajime Taniuchi², Takeshi Kuritani³, Akiko Matsumoto³ (1. Shizuoka Univ., 2. AIST, 3. Hokkaido Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R7-P-13] Research and development for the exploration of unknown cosmic ray events using Paleo-Detector

*Yuki Ido¹, Tatsuhiko Naka², Shota Futamura³, Tohma Ori⁴, Takenori Kato⁵ (1. Nagoya Univ. Env., 2. Toho Univ. Sci, 3. Nagoya Univ. Sci., 4. N.I.T. Suzuka, 5. Nagoya Univ. ISEE)

Poster presentation | R8: Metamorphic rocks and tectonics

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC | Entrance Hall Higashiyama Campus

R8: Metamorphic rocks and tectonics

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R8-P-01] Mineralogical characterization of serpentinite varieties in Sangun-Renge Belt exposed at Sasaguri, Fukuoka Prefecture, and their geological implications

「発表賞エントリー」

*Swarna ANNADURAI MUNUSAMY¹, Jun-ichi ANDO^{1,2}, Yuki IWASAKI³, Kaushik DAS^{1,2}, Dyuti Prakash SARKAR⁴, Seiichiro UEHARA⁵ (1. Hiroshima Univ., 2. HiPeR, Hiroshima, 3. NIPPON STEEL CORP., 4. JAMSTEC, 5. The Kyushu Univ. Museum)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R8-P-02] Metamorphic temperature structure of Sanbagawa Metamorphic Belt in the southern part of Shinshiro City, Aichi Prefecture, Japan

「発表賞エントリー」

*Akane Matsuzaki¹, Yui Kouketsu¹, Katsuyoshi Michibayashi¹ (1. Nagoya Univ. Env.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R8-P-03] Origin and pyrometamorphism of gneissose granitoid xenoliths from Mt. Daisen, Tottori Prefecture, SW Japan

「発表賞エントリー」

*Mizuki TAKAHASHI¹, Shunsuke Endo¹ (1. Shimane University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R8-P-04] Petrography of monazite in a metapelite in the eastern Nepal Himalaya and Development of Th-Pb dating method for monazite

「発表賞エントリー」

*Shumpei KUDO¹, Tetsuo Kawakami¹, Sota Niki², Toru Nakajima³, Takafumi Hirata⁴, Takeshi Imayama⁵ (1. Kyoto Univ. Sci., 2. Nagoya Univ. ISEE., 3. JAEA, 4. UTokyo. Sci., 5. Okayama Univ. of Sci.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R8-P-05] Petrological description of ultramafic rocks in the low-grade metamorphic zone of the Sanbagawa belt: A case study of the Ina area, Nagano Prefecture, central Japan

「発表賞エントリー」

*Kaho Nobuhara¹, Hiroshi Mori¹, Takayoshi Nagaya² (1. Shinshu Univ. , 2. Tokyo Gakugei Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R8-P-06] Detrital zircon U–Pb dating and Raman spectral analysis of carbonaceous material in the boundary area of the Sanbagawa–Chichibu belts, central Kii Peninsula

*Hiroshi MORI¹, Kojiro USUI^{1,2}, Tetsuya Tokiwa¹, Kazuhiro Ozawa³ (1. Shinshu University, 2. Nippon Koei Co., Ltd., 3. Precision Forestry Measurement Ltd.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R8-P-07] New finding of wakefieldite from an amphibolite in the Horokanai area, Kamuikotan HP metamorphic belt, Hokkaido, Japan

*Taro Kato¹, Kosuke NAEMURA¹, Toru Takeshita² (1. Iwate University, 2. Pacific Consultants Co., Ltd.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R8-P-08] Thermal history and protolithic detritus provenance of a sillimanite–chrysoberyl-bearing gneiss from the Ashio mountains in the western part of Tochigi prefecture

*Ippei KITANO¹ (1. Hokkaido Univ. Mus.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R8-P-09] Petrological characterization and geochronology of metamorphic rocks from the Northern Subzone of the Maizuru Terrane

*Sota Muroi¹, Kaushik Das¹, Kenta Kawaguchi¹, Yasutaka Hayasaka² (1. Hiroshima University, 2. Amakusa Museum of Goshoura Dinosaur Island)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R8-P-10] The fate of organic carbon during subduction: Raman micro-spectroscopy and C-isotope geochemistry of carbonaceous materials in Sambagawa pelitic schists, central Shikoku, Japan

*Hironobu Harada¹, Tatsuki Tsujimori¹, Akizumi Ishida¹, Takeshi Kakegawa¹, Tetsumaru Itaya² (1. Tohoku Univ., 2. jGnet)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R8-P-11] Fluid inclusions of ophicarbonates in the Apennine Mountains, Italy

Hiroyuki Kaneko¹, *Tatsuhiko KAWAMOTO¹, Francesca Meneghini², Yosuke Osawa¹ (1. Shizuoka University, 2. Università di Pisa | UNIPI · Department of Earth Sciences)

JAMS General Meeting, Award ceremony

2:00 PM - 3:15 PM JST | 5:00 AM - 6:15 AM UTC | ES Hall Higashiyama Campus

JAMS General Meeting, Award ceremony

2:00 PM - 3:15 PM JST | 5:00 AM - 6:15 AM UTC

[2Lecture-1-1400-1add] 定時総会・授賞式

Lectures of JAMS Awardees

3:30 PM - 5:00 PM JST | 6:30 AM - 8:00 AM UTC | ES Hall Higashiyama Campus

Lectures of JAMS Awardees

Sat. Sep 14, 2024

Oral presentation | R6: Plutonic rocks, volcanic rocks and subduction factory

9:00 AM - 12:00 PM JST | 12:00 AM - 3:00 AM UTC | ES Hall Higashiyama Campus

R6: Plutonic rocks, volcanic rocks and subduction factory

Chairperson: Tatsuhiko Kawamoto (Shizuoka University), Takashi Yuguchi, Atsushi Kamei

9:00 AM - 9:15 AM JST | 12:00 AM - 12:15 AM UTC

[R6-01] Compositional evolution of slab-derived fluids during ascent and origin of subduction-zone primary magmas: Implications from trace-element partition between hydrous melts and Cl-free or Cl-rich aqueous fluids

*Hajime TANIUCHI¹, Tatsuhiko KAWAMOTO², Takayuki NAKATANI¹, Osamu ISHIZUKA¹, Toshihiro SUZUKI¹, Akihiko Tomiya¹ (1. GSJ, AIST, 2. Shizuoka Univ.)

9:15 AM - 9:30 AM JST | 12:15 AM - 12:30 AM UTC

[R6-02] New crust and mantle physical properties research from the development of rock synthesis methods

「招待講演」

*Sanae Koizumi¹ (1. The University of Tokyo)

9:30 AM - 9:45 AM JST | 12:30 AM - 12:45 AM UTC

[R6-03] The use of contact metamorphism to identify plutons associated with major contemporaneous eruption and estimate magma flux

*Simon R WALLIS¹, Ken Yamaoka², Akira Miyake³, Catherine Annen⁴ (1. The University of Tokyo, 2. AIST, 3. Aichi University of Education, 4. Institute of Geophysics, Czech Academy of Sciences)

9:45 AM - 10:00 AM JST | 12:45 AM - 1:00 AM UTC

[R6-04] Petrography and geochronology of the Kuki granite, Kitakami Mountains, northeastern Japan: Shallow crustal intrusion and emplacement processes of granitic magma

*Satoshi SUZUKI¹, Takashi YUGUCHI², Keito ISHIGURO¹, Kyoka ENDO¹, Asuka KATO¹, Kosuke YOKOYAMA¹, Yasuhiro OGITA³, Tatsunori YOKOYAMA³, Shuhei SAKATA⁴, Takeshi OHNO⁵, Eiji SASAO³ (1. Yamagata University, 2. Kumamoto University, 3. Japan Atomic Energy Agency, 4. University of Tokyo, 5. Gakushuin University)

10:00 AM - 10:15 AM JST | 1:00 AM - 1:15 AM UTC

[R6-05] Zircon U-Pb dating of granitoids in the Hobenzan area, western Yamaguchi Prefecture, southwest Japan

*Yukiyasu TSUTSUMI¹, Kenichiro TANI¹ (1. National Museum of Nature and Science)

10:15 AM - 10:30 AM JST | 1:15 AM - 1:30 AM UTC

[R6-06] Estimation of temperature and pressure conditions of granitoids in the eastern part of Shimane Prefecture, San'in Belt.

*Kai Nakahashi¹, Satoshi SAITO¹ (1. Ehime University Graduate School of Science and Engineering, Earth Science Division)

10:30 AM - 10:45 AM JST | 1:30 AM - 1:45 AM UTC

[R6-07] Primary magma of Cretaceous Northern Kyushu batholith: New approach from Cumulus rocks

*Keisuke ESHIMA¹, Masaaki OWADA¹, Atsushi KAMEI² (1. Yamaguchi Univ. Sci., 2. Shimane Univ.)

10:45 AM - 11:00 AM JST | 1:45 AM - 2:00 AM UTC

[R6-08] Origin of the Cretaceous ignimbrite flare-up in SW Japan

*Ken Yamaoka¹, Tokiyuki Morohoshi², Daisuke Sato¹, Kazuto Mikuni¹ (1. AIST, 2. Univ. Tokyo. Sci.)

11:00 AM - 11:15 AM JST | 2:00 AM - 2:15 AM UTC

[3Lecture-101-11-9add] 休憩

11:15 AM - 11:30 AM JST | 2:15 AM - 2:30 AM UTC

[R6-09] The genesis of garnet in the Quaternary rhyolite of Hime-shima volcanic group, EEast Kyushu, Southwest Japan Arc.

*Takehiro HIRAYAMA^{1,2}, Tomoyuki SHIBATA², Masako YOSHIKAWA² (1. Tokyo University, 2. Hiroshima University)

11:30 AM - 11:45 AM JST | 2:30 AM - 2:45 AM UTC

[R6-10] K–Ar ages and geochemical compositions of Early Pleistocene andesites from southern Kyushu, Japan

*Raiki YAMADA¹, Masakazu Niwa¹, Koji Umeda², Koshi Yagi³, Taisei Fujiwara³, Ikuo Okada³, Fukuka Kida¹ (1. Japan Atomic Energy Agency, 2. Hirosaki University, 3. Hiruzen Institute for Geology and Geochronology)

11:45 AM - 12:00 PM JST | 2:45 AM - 3:00 AM UTC

[R6-11] Thermal history of pluton formation from Sr diffusion in plagioclase: implications for magma flux estimation and identification of plutons associated with major eruptions

「発表賞エントリー」

*Tokiyuki Morohoshi¹, Simon Richard Wallis¹ (1. The University of Tokyo)

Oral presentation | R2: Crystal structure, crystal chemistry, physical properties of minerals, crystal growth and applied mineralogy

9:00 AM - 12:00 PM JST | 12:00 AM - 3:00 AM UTC | ES024 Higashiyama Campus

R2: Crystal structure, crystal chemistry, physical properties of minerals, crystal growth and applied mineralogy

Chairperson: Fumiya Noritake, Mariko Nagashima, Makoto Tokuda

9:00 AM - 9:20 AM JST | 12:00 AM - 12:20 AM UTC

[R2-01] Changes of physical and structural properties of hydrous minerals induced by dehydration of coordinated water molecules

「招待講演」

*Ryo YAMANE¹ (1. Tohoku University)

9:20 AM - 9:35 AM JST | 12:20 AM - 12:35 AM UTC

[R2-02] Phase change of priceite $\text{Ca}_2\text{B}_5\text{O}_7(\text{OH})_5 \cdot \text{H}_2\text{O}$ during thermal decomposition

*Atsushi KYONO¹, Kosuke Yamaguchi¹, Satoru Okada¹, Hiroki Hasegawa¹ (1. Life & Environmental Sciences, University of Tsukuba)

9:35 AM - 9:50 AM JST | 12:35 AM - 12:50 AM UTC

[R2-03] The incorporation mechanisms of water in aluminous orthoenstatite clarified via comprehensive NMR measurements and first-principles calculations

*Xianyu XUE¹, Masami Kanzaki¹, Rongzhang Yin² (1. Okayama University, 2. Peking University)

9:50 AM - 10:05 AM JST | 12:50 AM - 1:05 AM UTC

[R2-04] Evaluation of the influence of polishing condition on hydrogen diffusion in fluorapatite and estimation of the diffusion mechanism

「発表賞エントリー」

*Chikashi Yoshimoto¹, Isao Sakaguchi², Shoichi Itoh¹ (1. Kyoto University, 2. NIMS)

10:05 AM - 10:20 AM JST | 1:05 AM - 1:20 AM UTC

[R2-05] Evaporation kinetics of forsterite in low-pressure H_2 - H_2O atmosphere

「発表賞エントリー」

*Shiori Inada¹, Shogo Tachibana¹ (1. The University of Tokyo)

10:20 AM - 10:30 AM JST | 1:20 AM - 1:30 AM UTC

[3Lecture-201-11-6add] 休憩

10:30 AM - 10:45 AM JST | 1:30 AM - 1:45 AM UTC

[R2-06] Origine of framboidal goethite in sandstone in the Saiki Subgroup of the Lower Shimanto Group

*Hiroaki OHFUJI¹, Seira Konishi¹, Hiroshi Yamamoto² (1. Tohoku Univ. Sci, 2. Kagoshima Univ. Sci)

10:45 AM - 11:00 AM JST | 1:45 AM - 2:00 AM UTC

[R2-07] Precursor phenomenon of ultrananolite crystallization: liquid immiscibility in the volcanic glass

*Mayumi Mujin¹, Keiko Matsumoto², Akira Miyake³, Yohei Igami³, Michihiko Nakamura¹, Akihiko Yokoo³, Mitsuhiro Yoshimoto⁴ (1. Tohoku University, 2. National Institute of Advanced Industrial Science and Technology (AIST), 3. Kyoto University, 4. Mount Fuji Research Institute, Yamanashi Prefectural Government)

11:00 AM - 11:15 AM JST | 2:00 AM - 2:15 AM UTC

[R2-08] Calorimetric constraint of cation disordering enthalpy in MgAl_2O_4 spinel

*Hiroshi KOJITANI¹, Ryuji Matsuki¹, Yuta Nishimura¹, Itaru Ohira¹ (1. Gakushuin University)

11:15 AM - 11:30 AM JST | 2:15 AM - 2:30 AM UTC

[R2-09] Solid solubilities of Na^+ and K^+ in perovskite type SrTiO_3

Ryogo KITAI², *Takaya NAGAI¹, Jun KAWANO¹, Ayako Shinozaki¹ (1. Hokkaido Univ. Faculty Sci., 2. Hokkaido Univ. School Sci.)

11:30 AM - 11:45 AM JST | 2:30 AM - 2:45 AM UTC

[R2-10] Study on the consistency of Li conductivity measurements of (Li,La)TiO₃ with perovskite structure and Li migration during SIMS measurements

*Isao SAKAGUCHI¹, Tsuyoshi Onishi¹ (1. National Institute for Materials Science)

11:45 AM - 12:00 PM JST | 2:45 AM - 3:00 AM UTC

[R2-11] Structure transition mechanism under high pressure of Fe-Ti-O system phases

*Takamitsu YAMANAKA^{1,2}, Takanori Hattori³, Ho-kwang Mao¹ (1. Center for High Pressure Science and Technology Advanced Research, 2. Graduate School of Science Osaka University, 3. J-PARC)

Oral presentation | R5: Extraterrestrial materials

9:00 AM - 12:00 PM JST | 12:00 AM - 3:00 AM UTC | ES025 Higashiyama Campus

R5: Extraterrestrial materials

Chairperson: Megumi Matsumoto, Daiki Yamamoto, Yusuke Seto, Toru Matsumoto (Kyoto University)

9:00 AM - 9:15 AM JST | 12:00 AM - 12:15 AM UTC

[R5-01] Revealing Chemical Compositional Variety of Amorphous Silicate Dust around AGB Stars by Condensation Experiment and Spectroscopic Analysis

「発表賞エントリー」

*Hanako Enomoto¹, Aki Takigawa¹, Hiroki Chihara², Chiyoeko Koike² (1. UTokyo, 2. Osaka Sangyo univ.)

9:15 AM - 9:30 AM JST | 12:15 AM - 12:30 AM UTC

[R5-02] Formation process of circumstellar dust studied from presolar oxides in carbonaceous chondrite DOM 08006.

「発表賞エントリー」

*Hiroyuki Hashizume¹, Aki Takigawa¹ (1. UTokyo EPSS)

9:30 AM - 9:45 AM JST | 12:30 AM - 12:45 AM UTC

[R5-03] Progress of irreversible chemical reactions on dust moving in a protoplanetary disk

*Lily Ishizaki¹, Shogo Tachibana¹ (1. UT EPS)

9:45 AM - 10:00 AM JST | 12:45 AM - 1:00 AM UTC

[R5-04] Experimental study of the effect of evaporation during temperature rise on type B CAI formation

「発表賞エントリー」

*Yasuaki Tsuruoka¹, Shogo Tachibana¹ (1. UTokyo Sci.)

10:00 AM - 10:20 AM JST | 1:00 AM - 1:20 AM UTC

[R5-05] Chondrule formation by collisions of planetesimals containing volatiles

「招待講演」

*Sin-iti Sirono¹ (1. Navoya Univ.)

10:20 AM - 10:35 AM JST | 1:20 AM - 1:35 AM UTC

[3Lecture-301-10-6add] 休憩

10:35 AM - 10:55 AM JST | 1:35 AM - 1:55 AM UTC

[R5-06] Constraining the true timescale of chondrule formation through chondrule mineral chemistry

「招待講演」

*Kohei Fukuda¹, Noriko T Kita², Makoto Kimura³ (1. Osaka Univ. Sci., 2. UW-Madison, 3. NIPR)

10:55 AM - 11:10 AM JST | 1:55 AM - 2:10 AM UTC

[R5-07] Reproduction Experiments of Radial Pyroxene Chondrules Using a Gas-jet Levitation System under Reducing Conditions

「発表賞エントリー」

*Kana Watanabe¹, Tomoki Nakamura¹, Tomoyo Morita¹ (1. Tohoku Univ. Sci.)

11:10 AM - 11:25 AM JST | 2:10 AM - 2:25 AM UTC

[R5-08] Stratigraphy of poikilitic shergottite parent body(ies) as deduced from their textural and mineral compositional diversities

「発表賞エントリー」

*Sojiro Yamazaki¹, Akira Yamaguchi², Takashi Mikouchi³ (1. Fac. Sci., Univ. Tokyo, 2. Natl. Inst. Polar Res., 3. Univ. Museum, Univ. Tokyo)

11:25 AM - 11:40 AM JST | 2:25 AM - 2:40 AM UTC

[R5-09] High pressure phase transition of Ti-Cr-rich spinel in Asuka 12325 Martian meteorite

*Atsushi TAKENOUCHI¹, Yohei Igami², Takashi Mikouchi³, Akira Miyake², Akira Yamaguchi⁴ (1. The Kyoto Univ. Museum, 2. Kyoto Univ. Sci., 3. The Univ. Museum, The Univ. of Tokyo, 4. NIPR)

11:40 AM - 11:55 AM JST | 2:40 AM - 2:55 AM UTC

[R5-10] Estimation of thermal history using silica minerals in mesosiderites

*Haruka OONO¹, Atsushi Takenouchi², Takashi Mikouchi³, Akira Yamaguchi⁴, Naoji Sugiura⁵, Akira Miyake¹ (1. Kyoto Univ., 2. Museum of Kyoto Univ., 3. Museum of Univ. Tokyo, 4. NiPR, 5. Chiba Int. Tech.)

Oral presentation | R6: Plutonic rocks, volcanic rocks and subduction factory

2:00 PM - 3:00 PM JST | 5:00 AM - 6:00 AM UTC | ES Hall Higashiyama Campus

R6: Plutonic rocks, volcanic rocks and subduction factory

Chairperson: Takashi Hoshide (Akita University)

2:00 PM - 2:20 PM JST | 5:00 AM - 5:20 AM UTC

[R6-12] Zirconium minerals in ultramafic rocks

「招待講演」

*Hikaru Sawada^{1,2}, Ryosuke Oyanagi^{3,2} (1. University of Toyama, 2. JAMSTEC, 3. Kokushikan University)

2:20 PM - 2:35 PM JST | 5:20 AM - 5:35 AM UTC

[R6-13] Estimation of physical conditions during granitic magma crystallization process using melt inclusions in zircon

*Yuka TANIWAKI¹, Satoshi SAITO² (1. Ciatec., Co LTD, 2. Ehime Univ.)

2:35 PM - 2:50 PM JST | 5:35 AM - 5:50 AM UTC

[R6-14] Estimation of TiO₂ activity for accurate determination of quartz crystallization temperature

*Yasuhiro OGITA¹, Takenori Kato², Takashi Yuguchi³ (1. Yamagata Univ., 2. Nagoya Univ., 3. Kumamoto Univ.)

Oral presentation | R2: Crystal structure, crystal chemistry, physical properties of minerals, crystal growth and applied mineralogy

2:00 PM - 3:30 PM JST | 5:00 AM - 6:30 AM UTC | ES024 Higashiyama Campus

R2: Crystal structure, crystal chemistry, physical properties of minerals, crystal growth and applied mineralogy

Chairperson: Kazuki 小松 Komatsu

2:00 PM - 2:30 PM JST | 5:00 AM - 5:30 AM UTC

[R2-12] Structural stabilization of pyrite-type compounds and chemical bonds of anions.

「招待講演」

*Akira YOSHIASA¹, Makoto Tokuda¹, Ginga Kitahara¹, Akihiko Nakatsuka², Kazumasa Sugiyama³ (1. Kumamoto University, 2. Yomaguchi University, 3. Tohoku University)

2:30 PM - 2:45 PM JST | 5:30 AM - 5:45 AM UTC

[R2-13] Formation of Fundamental Building Blocks Composed of Calcium Borate Minerals

「発表賞エントリー」

*Kosuke Yamaguchi¹, Atsushi Kyono², Satoru Okada¹, Hiroki Hasegawa¹ (1. Univ of Tsukuba. Grad. Sch. of Life and Env., Sci, 2. Univ of Tsukuba. Life and Environmental Science)

2:45 PM - 3:00 PM JST | 5:45 AM - 6:00 AM UTC

[R2-14] Crystal structures of kimuraite-(Y) and lokkaite-(Y)

*Ritsuro Miyawaki¹, Koichi Momma¹ (1. National Museum of Nature and Science)

3:00 PM - 3:15 PM JST | 6:00 AM - 6:15 AM UTC

[R2-15] Thermal Vibration of Oxygen atoms in Keatite Crystal

*Fumiya NORITAKE¹ (1. University of Yamanashi)

3:15 PM - 3:30 PM JST | 6:15 AM - 6:30 AM UTC

[R2-16] Development of a simulation method for electron back-scattered diffraction based on dynamical theory and its application to mineralogy

*Yusuke SETO¹, Masahiro Ohtsuka² (1. Osaka Metropolitan Univ. Sci., 2. Nagoya Univ. Sci. IMaSS.)

Oral presentation | R5: Extraterrestrial materials

2:00 PM - 3:30 PM JST | 5:00 AM - 6:30 AM UTC | ES025 Higashiyama Campus

R5: Extraterrestrial materials

Chairperson: Megumi Matsumoto, Daiki Yamamoto, Yusuke Seto, Toru Matsumoto (Kyoto University)

2:00 PM - 2:15 PM JST | 5:00 AM - 5:15 AM UTC

[R5-11] Mineralogy of micrometeoroids captured in silica aerogel aboard the ISS by the Dandelion Project

*Takaaki NOGUCHI¹, Akira MIYAKE¹, Hikaru Yabuta², Yoko Kebukawa³, Daiki Suga⁴, Makoto Tabata⁵, Kyoko Okudaira⁶, Akihiko Yamagishi⁷, Hajime Yano⁸ (1. Kyoto University, 2. Hiroshima University, 3. Tokyo Institute of Technology, 4. JASRI, 5. Chiba University, 6. University of Aizu, 7. Tokyo University of Pharmacy and Life Sciences, 8. JAXA)

2:15 PM - 2:30 PM JST | 5:15 AM - 5:30 AM UTC

[R5-12] STEM analysis of sodium-rich grains recovered from asteroid Ryugu

*Toru MATSUMOTO¹, Takaaki Noguchi¹, Toru Araki², Hayato Yuzawa², Akira Miyake¹ (1. Kyoto University, 2. UVSOR)

2:30 PM - 2:45 PM JST | 5:30 AM - 5:45 AM UTC

[R5-13] Important occurrence of amorphous hydrated Na-bearing Mg phosphate in Asteroid Ryugu samples and CI chondrites

*Takashi MIKOUCHI¹, Hideto Yoshida², Satoru Hayashi², Minami Masuda², Hiroyuki Kagi², Kazumasa Sugiyama³, Tomoki Nakamura⁴, Michael Zolensky⁵ (1. Univ. Museum, Univ. of Tokyo, 2. Fac. of Sci., Univ. of Tokyo, 3. Inst. Mat. Res., Tohoku Univ., 4. Fac. of Sci., Tohoku Univ., 5. NASA-JSC)

2:45 PM - 3:00 PM JST | 5:45 AM - 6:00 AM UTC

[R5-14] On the relationship between lithological classification and degree of aqueous alteration in Asteroid Ryugu samples

*Minami Masuda¹, Takashi Mikouchi², Hideto Yoshida¹, Tomoki Nakamura³, Michael Zolensky⁴ (1. Univ. Tokyo, Sci., 2. Univ. Tokyo, Univ. Museum, 3. Tohoku Univ., Sci., 4. NASA-JSC)

3:00 PM - 3:15 PM JST | 6:00 AM - 6:15 AM UTC

[R5-15] Highly primitive clasts with GEMS-like grains in Ryugu C0002 sample

*Megumi MATSUMOTO¹, Akira Tsuchiyama^{2,3}, Masahiro Yasutake⁴, Tomoki Nakamura¹ (1. Tohoku Univ., 2. Ritsumeikan Univ., 3. GIG, 4. JASRI/SPRing-8)

3:15 PM - 3:30 PM JST | 6:15 AM - 6:30 AM UTC

[R5-16] Aqueous alteration experiments using GEMS simulant particles

*Akira TSUCHIYAMA^{1,2}, Satomi Enju³, Haiyang Xian¹, Mingqi Sun¹, Ding Xing¹, Yuanyuan Wang⁴, Zhaolu He¹, Jing Du¹, Shan Li¹, Jiaxin Xi¹, Xiaojun Lin¹, Jian Di¹, Yuto Imura⁵ (1. Guangzhou Inst. Geochem., 2. Ritsumeikan Univ., 3. Ehime Univ. Sci., 4. Tokyo Univ. Sci., 5. Kyoto Univ. Sci.)

Poster presentation | R2: Crystal structure, crystal chemistry, physical properties of minerals, crystal growth and applied mineralogy

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC | Entrance Hall Higashiyama Campus

R2: Crystal structure, crystal chemistry, physical properties of minerals, crystal growth and applied mineralogy

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R2-P-01] Advanced four-circle goniometer system for single-crystal diffraction measurements at BL-10A, PF: anomalous X-ray scattering method & in situ high-pressure experiments

*Takahiro KURIBAYASHI¹, Ryo Yamane², Keita Harada¹, Toshiro Nagase³, Kazumasa Sugiyama² (1. Tohoku University (Sci), 2. Tohoku University (IMR), 3. Tohoku University (Mus))

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R2-P-02] Fine Structural Analysis using Single-Crystal Anomalous X-ray Scattering in KEK Photon Factory BL-6C

*Makoto TOKUDA¹, Takumi Ichimura², Koichi Momma⁶, Ritsuro Miyawaki⁶, Takashi Mikouchi⁴, Akira Yoshiasa³, Kazumasa Sugiyama⁵ (1. IINa, Kumamoto Univ., 2. Tohoku Univ. Eng, 3. Kumamoto Univ. Sci, 4. Univ. of Tokyo UMUT, 5. IMR, Tohoku Univ., 6. Nat'l. Mus. Nat. Sci.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R2-P-03] Evaluation of Fe K-edge XANES anisotropy and accurate determination of micro-area Fe²⁺/Fe³⁺ in clinopyroxene using multivariate analysis

「発表賞エントリー」

*Taisuke Ito¹, Simon Richard Wallis¹, Yoshio Takahashi¹ (1. The Univ. of Tokyo, EPS)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R2-P-04] Fine structural analysis of Diopside (Violan) containing Mn and Fe

「発表賞エントリー」

*Hayato Chiba¹, Kazumasa Sugiyama², Makoto Tokuda³, Takashi Mikouchi⁴ (1. Tohoku Univ. Eng., 2. Tohoku Univ. IMR, 3. Kumamoto Univ. IINa, 4. UMUT)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R2-P-05] Single-crystal X-ray diffraction study of a largely Cs-exchanged natural Ca-chabazite: crystal-chemical factors for its excellent Cs-exchange ability

Naomi KAWATA¹, Ryo FURUHASHI¹, Keiko FUJIWARA¹, Makio OHKAWA², *Akihiko NAKATSUKA¹ (1. Yamaguchi Univ. Sci. Tech. Innov., 2. Hiroshima Univ. Adv. Sci. Eng.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R2-P-06] Distribution of Mn³⁺ and Mn⁴⁺ studied by X anomalous scattering: the details of 3×4 tunneled structure formed by MnO₆ octahedron in woodruffite

「発表賞エントリー」

*Christopher Miyazaki¹, Ryo Yamane², Kazumasa Sugiyama², Makoto Tokuda³, Takeshi Mikouchi⁴ (1. Tohoku Univ. Eng., 2. Tohoku Univ. IMR, 3. Kumamoto Univ. IINa, 4. UMUT)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R2-P-07] Mg²⁺ and Ba²⁺ ion-exchange of GTS-type Na titanosilicate and its structural characterization

*Keiko FUJIWARA¹, Naomi Kawata¹, Akihiko Nakatsuka¹ (1. Yamaguchi Univ. Sci. Tech. Innov.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R2-P-08] Distribution of Fe³⁺ among tetrahedral sites and crystal structural change of gehlenite-Fe³⁺ gehlenite

*Maki HAMADA¹, Mariko Nagashima² (1. Kanazawa University, 2. Yamaguchi University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R2-P-09] Crystallographic site preference of impurity elements in olivine by ALCHEMI method

*Yohei IGAMI¹, Akira MIYAKE¹, Ryo WAKABAYASHI¹, Norikatsu AKIZAWA² (1. Kyoto University, 2. The University of Tokyo)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R2-P-10] Precipitated structures of pyroxene lamellar with various directions

*Shoichi TOH¹ (1. Fukuoka University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R2-P-11] Crystal chemistry of bridgmanite with Al/Fe contents expected in MORB component

*Akihiko NAKATSUKA¹, Hiroshi FUKUI², Seiji KAMADA³, Naohisa HIRAO², Makio OHKAWA⁴, Kazumasa SUGIYAMA⁵, Takashi YOSHINO⁶ (1. Yamaguchi Univ. Sci. Tech. Innov., 2. JASRI, 3. AD Science Inc., 4. Hiroshima Univ. Adv. Sci. Eng., 5. Tohoku Univ. IMR, 6. Okayama Univ. IPM)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R2-P-12] Mechanism of the pressure-induced crystallization of amorphous calcium carbonate

*Chiho Morita¹, Hiroki Kobayashi¹, Hiroyuki Kagi¹ (1. UTokyo. Sci.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R2-P-13] Differences in the effects of Li and Cs on the calcitization of vaterite.

*Noboru FURUKAWA¹, Ami Kinoda² (1. Chiba University, 2. NIWAKA)

Poster presentation | R5: Extraterrestrial materials

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC | Entrance Hall Higashiyama Campus

R5: Extraterrestrial materials

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R5-P-01] Dehydration of Newberyite($\text{MgHPO}_4 \cdot 3\text{H}_2\text{O}$) under low pressures

*Yukako Matsumoto¹, Shogo Tachibana¹ (1. UTokyo)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R5-P-02] Experimental simulation of oxygen isotope exchange between chondrule melt and low-pressure water vapor

*Daiki YAMAMOTO¹, Noriyuki Kawasaki² (1. Kyushu University, 2. Hokkaido University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R5-P-03] Early impact events recorded in anomalous eucrites

*Akira YAMAGUCHI¹ (1. National Institute of Polar Research)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R5-P-04] Internal Structure Analysis of Carbonaceous Material in Yamato 8448 Ureilite

*Masahiro YASUTAKE¹, Megumi Matsumoto², Akira Tsuchiyama^{3,4}, Kentaro Uesugi¹, Akihisa Takeuchi¹, Akira Yamaguchi⁵ (1. JASRI, 2. Tohoku Univ., 3. Ritsumeikan Univ., 4. CAS/GIG, 5. NIPR)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R5-P-05] Effects of initial porosity on shock deformation microstructures in rutile

*Yuhei Umeda^{1,2}, Yuma Nagai^{1,2}, Naotaka Tomioka³, Toshimori Sekine⁴, Masashi Miyakawa⁵, Takamichi Kobayashi⁵, Hitoshi Yusa⁵, Takuo Okuchi^{1,2} (1. KURNS, 2. Kyoto Univ. Eng., 3. JAMSTEC, 4. HPSTAR, 5. NIMS)

Poster presentation | R6: Plutonic rocks, volcanic rocks and subduction factory

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC | Entrance Hall Higashiyama Campus

R6: Plutonic rocks, volcanic rocks and subduction factory

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R6-P-01] Orthopyroxene cumulates from a serpentinite mud-volcano in the Mariana forearc: constituent rocks of the forearc Moho transition zone

*Pia Ataka¹, Yuji Ichiyama¹, Shun Takamizawa¹, Akihiro Tamura², Tomoaki Motishita² (1. Chiba University, 2. Kanazawa University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R6-P-02] Formation processes of high-Mg andesite associated with ophiolitic rocks in the Setogawa belt

*Tsumugi Kato¹, Yuji Ichiyama¹, Akihiro Tamura², Tomoaki Morishita² (1. Chiba Univ., 2. Kanazawa Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R6-P-03] Petrological and geochemical study of volcanics from Kuju volcanic group, Kyushu Island, Southwest Japan Arc.

「発表賞エントリー」

*Soma Yamanaka¹, Tomoyuki Shibata¹, Ryotaro Fujihara¹, Masako Yoshikawa¹, Tomo Shibata² (1. Hiroshima Univ. Sci. & Tech, 2. Fukuoka Univ. Sci.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R6-P-04] Petrological study of intrusive rocks and their crustal xenoliths in the Suzu area, Atsumi district, Yamagata

*Kazuhiro Tojo¹, Takashi Hoshide¹ (1. Akita Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R6-P-05] Crystallization conditions of zircon and its crystallization relationship with surrounding minerals in the Tono plutonic complex, Kitakami mountains

*Yasuhiro OGITA^{1,2}, Shuhei Sakata³, Takeshi Ohno⁴, Tatsunori Yokoyama², Satoshi Suzuki¹, Kyoka Endo¹, Takashi Yuguchi⁵ (1. Yamagata Univ., 2. JAEA, Tono Geoscience Center, 3. The Univ. Tokyo, Earthquake Research Inst., 4. Gakushuin Univ., 5. Kumamoto Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R6-P-06] Petrography and magmatic processes of a diorite body distributed in the Kawakami region, Yamaguchi Prefecture, Japan

「発表賞エントリー」

*Ryosei Sugiura¹, Keisuke Eshima¹ (1. Yamaguchi University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R6-P-07] Sr-Nd-Hf isotopic systematics for the Habu granodiorite, eastern part of Yamaguchi Prefecture

*Masaaki OWADA¹, Yusaku Akimoto², Mariko Nagashima¹, Keisuke Eshima¹, Atsushi Kamei³, Nobuhiko Nakano⁴, Kenta Kawaguchi⁵, Yasuhito Osanai⁴ (1. Yamaguchi Univ., 2. YON-C, 3. Shimane Uni., 4. Kyushu Univ., 5. Hiroshima Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R6-P-08] Rare earth element compositions and U-Pb zircon age of granodiorite porphyrites in the Noko Island, western part of northern Kyushu

*Masaki YUHARA¹, Yoshinobu KAWANO², Kazuhiro TSUKADA³, Purevdulam SUKHBAATAR⁴ (1. Fukuoka Univ. Sci., 2. Rissyo Univ. Env. Sci., 3. Nagoya Univ. Mus., 4. Nagoya Univ. Env. St.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R6-P-09] Compositional changes and growth processes of the Asakura granodiorite, Northern Kyushu, Japan.

「発表賞エントリー」

*Yudai HAMANO¹, Keisuke ESHIMA¹ (1. Yamaguchi Uni. Sci.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R6-P-10] Intrusive relationship and lithofacies classification of Granite–Granodiorite in Eastern Fukuoka Prefecture

「発表賞エントリー」

*Ryo Nakamura¹, Keisuke Eshima¹ (1. Yamaguchi University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R6-P-11] Crystallization processes of quartz in a granitic magma: Implications for the magma chamber processes of Okueyama granite, Kyushu, Japan

*Takashi YUGUCHI¹, Takenori Kato⁴, Yasuhiro Ogita², Minori Watanabe⁶, Asuka Kato⁶, Daichi Itoh⁶, Tatsunori Yokoyama², Shuhei Sakata³, Takeshi Ohno⁵ (1. Kumamoto University, 2. JAEA, 3. Uni. Tokyo ERI, 4. Nagoya University, 5. Gakushuin University, 6. Yamagata University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R6-P-13] Differentiation Processes of Granitic Magma Chamber Recorded in 'Oshima-ishi' Distributed in Oshima, Geiyo islands, Ehime Prefecture

*Kazuya SHIMOOKA^{1,2}, Soichiro AONO², Takahito ONISHI², Toko FUKUI², Motohiro TSUBOI¹, Toshiro TAKAHASHI³, Satoshi SAITO² (1. Kwansei Gakuin Univ., 2. Ehime Univ., 3. Niigata Univ.)

"Mineralogy 2022" General Outreach lecture

3:30 PM - 5:00 PM JST | 6:30 AM - 8:00 AM UTC | ES Hall Higashiyama Campus

General Outreach lecture

3:30 PM - 5:00 PM JST | 6:30 AM - 8:00 AM UTC

[3Lecture-1-1530-1add] 一般普及講演

Oral presentation | S1: Dynamics of igneous processes (Special Session)

📅 Thu. Sep 12, 2024 10:00 AM - 12:00 PM JST | Thu. Sep 12, 2024 1:00 AM - 3:00 AM UTC | 🏢 ES Hall Higashiyama Campus

S1: Dynamics of igneous processes (Special Session)

Chairperson: Shumpei Yoshimura (Hokkaido University), Yuuki Hagiwara (Japan Agency for Marine-Earth Science and Technology), Morihisa Hamada (JAMSTEC)

10:00 AM - 10:20 AM JST | 1:00 AM - 1:20 AM UTC

[S1-01] Tungsten (VI) speciation in subduction-zone aqueous fluids and its significance for W stable isotope fractionation during slab dehydration

「招待講演」

*Naoko TAKAHASHI¹, Michihiko Nakamura², Shigeru Yamashita³, Hiroyuki Kagi¹ (1. UTokyo Sci., 2. Tohoku Univ. Sci., 3. Okayama Univ. IPM)

10:20 AM - 10:40 AM JST | 1:20 AM - 1:40 AM UTC

[S1-02] Reconstruction of mantle structures using mantle xenoliths from Nushan, SE-edge of Sino-Korean craton: How far does the subduction fluid reach?

「招待講演」

*Yuto SATO¹, Eiichi Takahashi² (1. JAMSTEC Kochi, 2. GIG)

10:40 AM - 10:55 AM JST | 1:40 AM - 1:55 AM UTC

[S1-03] Elucidating Deep processes leading to maar eruptions using mantle xenoliths in the West Eifel volcanic field

「発表賞エントリー」

*Masanari Arai¹, Michihiko Nakamura¹, Mayumi Mujin¹, Naoki Araya¹, Sando Sawa¹, Takayuki Nakatani², Mari Sumita³, Hans-Ulrich Schmincke³ (1. Tohoku Univ. Sci., 2. AIST, 3. GEOMAR)

10:55 AM - 11:10 AM JST | 1:55 AM - 2:10 AM UTC

[S1-04] Water contents and pressures of melts in unerupted felsic magma constrained by SEM-EDS analysis of homogenized melt inclusions in zircon

「発表賞エントリー」

*Taichi Kawashima¹, Kazuya Shimooka², Toko Fukui¹, Satoshi Saito¹ (1. Ehime Univ., 2. Kwansei Gakuin Univ.)

11:10 AM - 11:25 AM JST | 2:10 AM - 2:25 AM UTC

[S1-05] Permeability and pore microstructures on the diktytaxitic texture of Ogurayama lava dome, Towada volcano

*Natsuko Sekiya¹, Michihiko Nakamura², Ryosuke Sakurai³, Mayumi Mujin², Naoki Araya², Shingo Takeuchi⁴, Yukiko Suwa⁵ (1. Tohoku Univ. Sci., 2. Grad. Sch. of Sci., Tohoku Univ., 3. Univ. Tokyo, 4. CRIEPI, 5. CERES)

11:25 AM - 11:40 AM JST | 2:25 AM - 2:40 AM UTC

[S1-06] MD simulation of silicate melt and glass under tension deformation

*Ryota Mamizuka¹, Satoshi Okumura¹, Hiroshi Sakuma² (1. Tohoku Univ. Sci., 2. NIMS)

11:40 AM - 11:55 AM JST | 2:40 AM - 2:55 AM UTC

[S1-07] The magma plumbing system of Izu-Oshima Volcano: Constraints from the H₂O-saturated plagioclase liquidus

*Hidemi ISHIBASHI¹ (1. Shizuoka University)

11:55 AM - 12:00 PM JST | 2:55 AM - 3:00 AM UTC

調整

Tungsten (VI) speciation in subduction-zone aqueous fluids and its significance for W stable isotope fractionation during slab dehydration

*Naoko TAKAHASHI¹, Michihiko Nakamura², Shigeru Yamashita³, Hiroyuki Kagi¹

1. UTokyo Sci., 2. Tohoku Univ. Sci., 3. Okayama Univ. IPM



Keywords: Diamond anvil cell, In-situ Raman spectroscopy, Tungsten (VI) speciation, Subduction zone

Reconstruction of mantle structures using mantle xenoliths from Nushan, SE-edge of Sino-Korean craton: How far does the subduction fluid reach?

*Yuto SATO¹, Eiichi Takahashi²

1. JAMSTEC Kochi, 2. GIG

Keywords: Mantle xenolith, Big mantle wedge, NAMs, Mantle petrology, FTIR



Elucidating Deep processes leading to maar eruptions using mantle xenoliths in the West Eifel volcanic field

*Masanari Arao¹, Michihiko Nakamura¹, Mayumi Mujin¹, Naoki Araya¹, Sando Sawa¹, Takayuki Nakatani², Mari Sumita³, Hans-Ulrich Schmincke³

1. Tohoku Univ.Sci., 2. AIST, 3. GEOMAR

Keywords: West Eifel volcanic field, Mantle xenolith, Fluid inclusion, Microcrack, Hydraulic fracturing

Water contents and pressures of melts in unerupted felsic magma constrained by SEM-EDS analysis of homogenized melt inclusions in zircon

*Taichi Kawashima¹, Kazuya Shimooka², Toko Fukui¹, Satoshi Saito¹

1. Ehime Univ., 2. Kwansei Gakuin Univ.

Granitic rocks (*sensu lato*) represent unerupted felsic magmas crystallized in the crust. In this study, we estimate water contents of melts and crystallization pressures of zircons in granitoid magma using melt inclusions in zircon, a ubiquitous accessory mineral in granitoids. Homogenization experiments of polymineralic inclusions hosted in zircon have been conducted for a granitoid sample from the Cretaceous Gamano granodiorite in Yashiro-jima Island, southwest Japan, using a piston-cylinder high-pressure-high-temperature apparatus. SEM-EDS analysis reveals that the homogenized melt inclusions have high water contents (6.4–11.3 wt%) and high SiO₂ contents (76–78 wt% anhydrous basis) implying that they represent fractionated interstitial hydrous melts trapped in growing zircon crystals. A recently proposed machine learning-based melt-phase assemblage geobarometer yields pressures ranging from 563 to 266 MPa interpreted as crystallization pressures of the zircons. The results of this study suggest high water activity of the interstitial melts within the Gamano granodiorite magma at the time of zircon crystallization. The melt inclusions in zircons record a wide range of pressures, from intrusion of the magma into the deeper crustal levels (~563–500 MPa) to final solidification at shallower levels (~266 MPa).

Keywords: Unerupted felsic magma, Water contents, Crystallization pressures, Melt inclusion in zircon

Permeability and pore microstructures on the diktytaxitic texture of Ogurayama lava dome, Towada volcano

*Natsuko Sekiya¹, Michihiko Nakamura², Ryosuke Sakurai³, Mayumi Mujin², Naoki Araya², Shingo Takeuchi⁴, Yukiko Suwa⁵

1. Tohoku Univ. Sci., 2. Grad. Sch. of Sci., Tohoku Univ., 3. Univ. Tokyo, 4. CRIEPI, 5. CERES

Keywords: permeability, porosity, evaporation-condensation, degassing, Gas pycnometer

MD simulation of silicate melt and glass under tension deformation

*Ryota Mamizuka¹, Satoshi Okumura¹, Hiroshi Sakuma²

1. Tohoku Univ. Sci., 2. NIMS

Keywords: silicate melt, MD simulation, tensional deformation, ring structure

The magma plumbing system of Izu-Oshima Volcano: Constraints from the H₂O-saturated plagioclase liquidus

*Hidemi ISHIBASHI¹

1. Shizuoka University

Keywords: Izu-Oshima volcano, Magma plumbing system, plagioclase, liquidus, pre-eruptive condition

Oral presentation | R1: Characterization and description of minerals (Joint Session with The Gemmological Society of Japan)

📅 Thu. Sep 12, 2024 10:00 AM - 12:00 PM JST | Thu. Sep 12, 2024 1:00 AM - 3:00 AM UTC | 🏠 ES024 Higashiyama Campus

R1: Characterization and description of minerals (Joint Session with The Gemmological Society of Japan)

Chairperson: Masanori Kurosawa, Hiroshi Kitawaki

10:00 AM - 10:20 AM JST | 1:00 AM - 1:20 AM UTC

[R1-01] Determination of analytical conditions in quantitative electron probe microanalysis of ultra-trace elements: probe current and sampling time

「招待講演」

*Takenori KATO¹ (1. ISEE, Nagoya University)

10:20 AM - 10:35 AM JST | 1:20 AM - 1:35 AM UTC

[R1-02] Mössbauer Spectra and Intensity Tensors of Quadrupole Doublets for Fe³⁺ and Fe²⁺ of Bridgmanite

*Keiji SHINODA¹, Yasuhiro Kobayashi², Takuo Okuchi² (1. Osaka Met. Univ. Sci, 2. KURNS)

10:35 AM - 10:50 AM JST | 1:35 AM - 1:50 AM UTC

[R1-03] Clay mineral characterization by Optical Photothermal Induced Resonance (O-PTIR)

「発表賞エントリー」

*Taro Kido¹, Yohey Suzuki¹ (1. The University of Tokyo)

10:50 AM - 11:05 AM JST | 1:50 AM - 2:05 AM UTC

[R1-04] Characterization and Identification of Synthetic Colored Diamonds for Jewelry Use

*Hiroshi KITAWAKI¹, Kentaro Emori¹, Mio Hisanaga¹, Masahiro Yamamoto¹ (1. Central Gem Laboratory)

11:05 AM - 11:15 AM JST | 2:05 AM - 2:15 AM UTC

[1Lecture-201-07-5add] 休憩

11:15 AM - 11:30 AM JST | 2:15 AM - 2:30 AM UTC

[R1-05] Blue Sapphire from Australia and its origin

*Kentaro Emori¹, Hiroshi Kitawaki¹ (1. Central Gem Laboratory)

11:30 AM - 11:45 AM JST | 2:30 AM - 2:45 AM UTC

[R1-06] Emerald origin characteristics and problems in origin determination

*Zhenghao Zhao¹, Hiroshi Kitawaki¹, Kentaro Emori¹ (1. Central Gem Laboratory)

11:45 AM - 12:00 PM JST | 2:45 AM - 3:00 AM UTC

[R1-07] Gem zircon and sapphire age dating and application of origin determination; A study from New England sapphire fields, New South Wales, Australia

*Ahmadjan ABDURIYIM¹ (1. Tokyo Gem Science LLC)

Determination of analytical conditions in quantitative electron probe microanalysis of ultra-trace elements: probe current and sampling time

*Takenori KATO¹

1. ISEE, Nagoya University

Keywords: electron probe microanalysis (EPMA), trace elements, quantitative analysis, analytical conditions

Mössbauer Spectra and Intensity Tensors of Quadrupole Doublets for Fe³⁺ and Fe²⁺ of Bridgmanite

*Keiji SHINODA¹, Yasuhiro Kobayashi², Takuo Okuchi²

1. Osaka Met. Univ. Sci, 2. KURNS

Keywords: Bridgmanite, Mössbauer spectroscopy, Intensity tensor

Clay mineral characterization by Optical Photothermal Induced Resonance (O-PTIR)

*Taro Kido¹, Yohey Suzuki¹

1. The University of Tokyo


Keywords: smectite, IR spectroscopy, Raman spectroscopy, X-ray diffraction, SEM-EDS

Characterization and Identification of Synthetic Colored Diamonds for Jewelry Use

*Hiroshi KITAWAKI¹, Kentaro Emori¹, Mio Hisanaga¹, Masahiro Yamamoto¹

1. Central Gem Laboratory

Keywords: melee size, synthetic diamond, fancy color diamond, CVD, HPHT



Oral presentation

R1: Characterization and description of minerals (Joint Session with The Gemmological Society of Japan)

Chairperson: Masanori Kurosawa, Hiroshi Kitawaki

Thu. Sep 12, 2024 10:00 AM - 12:00 PM ES024 (Higashiyama Campus)

11:05 AM - 11:15 AM

[1Lecture-201-07-5add]休憩

Blue Sapphire from Australia and its origin

*Kentaro Emori¹, Hiroshi Kitawaki¹

1. Central Gem Laboratory


Keywords: LA-ICP-MS, corundum, origin determination, Australia

Emerald origin characteristics and problems in origin determination

*Zhenghao Zhao¹, Hiroshi Kitawaki¹, Kentaro Emori¹

1. Central Gem Laboratory

Keywords: Emerald, Origin determination, FTIR Spectrum, UV-Vis-NIR Spectrum, LA-ICP-MS



Gem zircon and sapphire age dating and application of origin determination; A study from New England sapphire fields, New South Wales, Australia

*Ahmadjan ABDURIYIM¹

1. Tokyo Gem Science LLC

Keywords: zircon, sapphire, U-Pb age dating, geographic origin, LA-ICP-MS

Oral presentation | R3: High-pressure science and deep Earth's material

📅 Thu. Sep 12, 2024 10:00 AM - 12:00 PM JST | Thu. Sep 12, 2024 1:00 AM - 3:00 AM UTC | 📍 ES025 Higashiyama Campus

R3: High-pressure science and deep Earth's material

Chairperson: Takeshi Sakai (Ehime University), Ryosuke Sinmyo (Meiji University), Takayuki Ishii (Okayama University), Takaaki Kawazoe (Hiroshima University)

10:00 AM - 10:15 AM JST | 1:00 AM - 1:15 AM UTC

[R3-01] Pressure-induced polyamorphic transition in CaAl_2O_4 glass revealed by elastic wave velocity and X-ray diffraction measurements and molecular dynamics simulations

「招待講演」

*Itaru Ohira¹, Yoshio Kono^{2,3}, Steeve Gréaux³, James W E Drewitt⁴, Sandro Jahn⁵, Fumiya Noritake⁶, Koji Ohara^{7,8}, Satoshi Hiroi^{7,8}, Nozomi M Kondo⁹, Rostislav Hrubíak¹⁰, Yuji Higo⁸, Noriyoshi Tsujino⁸, Sho Kakizawa⁸, Kiyofumi Nitta⁸, Oki Sekizawa⁸ (1. Gakushuin Univ. Sci., 2. Kwansai Gakuin Univ. Sci., 3. Ehime Univ. GRC, 4. U. Bristol Phys., 5. U. Cologne IGM, 6. U. Yamanashi Interdisciplinary Research, 7. Shimane Univ. Materials for Energy, 8. JASRI, 9. Okayama Univ. IPM, 10. ANL HPCAT)

10:15 AM - 10:30 AM JST | 1:15 AM - 1:30 AM UTC

[R3-02] Structural analysis of CO_2 -bearing sodium silicate melts using molecular dynamics simulations and synchrotron radiation X-ray diffraction experiments

「発表賞エントリー」

*Shino HAYAFUNE¹, Yohei Onodera², Shinji Kohara², Haruki Ichikawa³, Tatsuya Sakamaki¹, Kenichi Funakoshi⁴, Akio Suzuki¹ (1. Tohoku Univ. Graduate School of Science., 2. NIMS, 3. Tohoku Univ. School of Science., 4. CROSS)

10:30 AM - 10:45 AM JST | 1:30 AM - 1:45 AM UTC

[R3-03] Possible presence of poirierite, a high-pressure phase of $(\text{Mg,Fe})_2\text{SiO}_4$, in the deep Earth's mantle

*Naotaka TOMIOKA¹, Takuo Okuchi², Narangoo Purevjav³, Masaaki Miyahara⁴ (1. KOCHI, JAMSTEC, 2. KRUNS, Kyoto Univ., 3. Seoul National Univ., 4. Earth Planet Sci., Hiroshima Univ.)

10:45 AM - 11:00 AM JST | 1:45 AM - 2:00 AM UTC

[R3-04] Temperature dependence of water content in wadsleyite and ringwoodite

*Toru INOUE^{1,2}, Yuji HAMADA², Hanae KAYA², Yusuke EGI¹, Daichi MAEDA¹, Kazutaka YAMAGUCHI¹, Konosuke YAMADA¹, Takaaki KAWAZOE^{1,2} (1. Hiroshima Univ. Adv. Sci. Eng., 2. Hiroshima Univ. Sci.)

11:00 AM - 11:15 AM JST | 2:00 AM - 2:15 AM UTC

[R3-05] Temperature dependence of the $\text{Fe}^{3+}/\Sigma\text{Fe}$ ratio in wadsleyite by electron energy loss spectroscopy (EELS)

「発表賞エントリー」

*Kazutaka YAMAGUCHI¹, Takaaki Kawazoe¹, Toru Inoue¹, Naotaka Tomioka² (1. Hiroshima University, 2. JAMSTEC, Kochi institute)

11:15 AM - 11:30 AM JST | 2:15 AM - 2:30 AM UTC

[R3-06] Advantage on the analysis of cation site occupancies by neutron diffraction and some examples of application for high-pressure minerals

*Takuo OKUCHI¹ (1. Kyoto Univ.)

11:30 AM - 11:45 AM JST | 2:30 AM - 2:45 AM UTC

[R3-07] In situ lattice volume observation of davemaioite in the $\text{CaSiO}_3\text{-H}_2\text{O}$ system up to uppermost lower mantle conditions

*Goru Takaichi¹, Takayuki Ishii², Yu Nishihara¹, Kyoko Matsukage³, Yuji Higo⁴, Noiyoshi Tsujino⁴, Sho Kakizawa⁴ (1. GRC, Ehime Univ., 2. IPM, Okayama Univ., 3. Teikyo University of Science, 4. JASRI)

11:45 AM - 12:00 PM JST | 2:45 AM - 3:00 AM UTC

[R3-08] Elasticity Measurements of Hydrous SiO₂ post-stishovite and its implication to the seismic anomalies in the lower mantle

「発表賞エントリー」

*Yuichiro MORI^{1,2}, Motohiko MURAKAMI², Takashi YOSHINO³, Hiroyuki KAGI¹ (1. UTokyo. Sci., 2. ETHZ. D-ERDW., 3. Okayama Univ. IPM)

Pressure-induced polyamorphic transition in CaAl_2O_4 glass revealed by elastic wave velocity and X-ray diffraction measurements and molecular dynamics simulations

*Itaru Ohira¹, Yoshio Kono^{2,3}, Steeve Gréaux³, James W E Drewitt⁴, Sandro Jahn⁵, Fumiya Noritake⁶, Koji Ohara^{7,8}, Satoshi Hiroi^{7,8}, Nozomi M Kondo⁹, Rostislav Hrubyak¹⁰, Yuji Higo⁸, Noriyoshi Tsujino⁸, Sho Kakizawa⁸, Kiyofumi Nitta⁸, Oki Sekizawa⁸

1. Gakushuin Univ. Sci., 2. Kwansei Gakuin Univ. Sci., 3. Ehime Univ. GRC, 4. U. Bristol Phys., 5. U. Cologne IGM, 6. U. Yamanashi Interdisciplinary Research, 7. Shimane Univ. Materials for Energy, 8. JASRI, 9. Okayama Univ. IPM, 10. ANL HPCAT

In recent years, there has been increasing emphasis on the fabrication and characterization of non-conventional network glass formers such as $\text{CaO-Al}_2\text{O}_3$, $\text{BaO-Al}_2\text{O}_3$, and BaO-TiO_2 in material science [e.g., 1]. Although these systems are different from those of magmas traditionally studied in geoscience field, recent experiments have confirmed the formation of network-modifiers- and Al-rich partial melts at the conditions of the transition zone and the lower mantle [2,3]. Therefore, the high-pressure studies of non-conventional network glass formers, as well as typical network-forming oxide glasses such as SiO_2 glass, are also important in understanding the pressure-induced changes of structure and physical properties of the magmas in the deep earth.

In this study, we conducted elastic wave velocity and XRD measurements and molecular dynamics (MD) simulations on CaAl_2O_4 glass. Ultrasonic velocity measurements up to 24 GPa at BL04B1 at SPring-8 reveal abrupt and irreversible increases in the v_p and v_s at ~ 8 –10 GPa. Total structure factor and pair distribution functions measured by synchrotron XRD at 16-BM-B at APS and BL37XU at SPring-8 show a rapid change in the intermediate range structure, which is likely attributed to a rearrangement of Ca ions over this narrow pressure condition. Structure models obtained from MD simulations reveal that this intermediate range structure is explained by a transition of Ca–O void radius distribution from a bimodal distribution with peaks at ~ 2.1 Å and ~ 2.4 Å to a single distribution centered at ~ 2.1 Å. The abrupt structural changes involving the rapid increase in elastic wave velocity in CaAl_2O_4 glass are markedly different to the continuous transformations reported in SiO_2 glass. The polyamorphic transition observed in this study may be one of the key mechanisms in the densification and the changes in seismic wave velocity of the magma just above the 410 km discontinuity.

References

- [1] A. Masuno, *J. Phys. Soc. Jpn.* **91**, 091003 (2022).
- [2] G. K. Pradhan *et al.*, *Earth Planet. Sci. Lett.*, **431**, 247 (2015).
- [3] A. Nakajima *et al.* *Sci. Rep.* **9**, 7420 (2019).


Keywords: oxide glass, polyamorphism, elastic wave velocity, synchrotron X-ray diffraction, molecular dynamics simulations

Structural analysis of CO₂-bearing sodium silicate melts using molecular dynamics simulations and synchrotron radiation X-ray diffraction experiments

*Shino HAYAFUNE¹, Yohei Onodera², Shinji Kohara², Haruki Ichikawa³, Tatsuya Sakamaki¹, Kenichi Funakoshi⁴, Akio Suzuki¹

1. Tohoku Univ. Graduate School of Science., 2. NIMS, 3. Tohoku Univ. School of Science., 4. CROSS

Keywords: Sodium Silicate melt, Amorphous Structure Analysis, CO₂



Possible presence of poirierite, a high-pressure phase of $(\text{Mg,Fe})_2\text{SiO}_4$, in the deep Earth's mantle

*Naotaka TOMIOKA¹, Takuo Okuchi², Narangoo Purevjav³, Masaaki Miyahara⁴

1. KOCHI, JAMSTEC, 2. KRUNS, Kyoto Univ., 3. Seoul National Univ., 4. Earth Planet Sci., Hiroshima Univ.

Keywords: olivine, poirierite, high-pressure phase transformation, transmission electron microscope

Temperature dependence of water content in wadsleyite and ringwoodite

*Toru INOUE^{1,2}, Yuji HAMADA², Hanae KAYA², Yusuke EGI¹, Daichi MAEDA¹, Kazutaka YAMAGUCHI¹, Konosuke YAMADA¹, Takaaki KAWAZOE^{1,2}

1. Hiroshima Univ. Adv. Sci. Eng., 2. Hiroshima Univ. Sci.

Many researchers agree that the main constituent minerals of the mantle transition zone (MTZ: corresponding to 410-660 km depth) are wadsleyite (Wd) and ringwoodite (Rw), which are high-pressure polymorph of olivine. Although these phases are nominally anhydrous minerals (NAM), it has been shown that water can be present in the crystal structure at several wt% levels (e.g. Inoue et al., 1995; Kohlstedt et al., 1996) and that the MTZ can act as a water reservoir in the Earth interiors. In this context, Pearson et al. (2014) found ~1.4 wt% hydrous Rw in diamond inclusions, indicating that the MTZ is at least locally hydrous. Since the water content of minerals has a great influence on their physical properties, it is important to clarify the temperature dependence of water content in Wd and Rw in order to discuss the dynamics of the Earth's deep interior. Although previous studies have already shown the temperature dependence (Ohtani et al., 2001; Litasov and Ohtani, 2003), the results are inconsistent, and it is difficult to conclude that the temperature dependence has been clearly determined. Therefore, this experimental study was conducted to reexamine the temperature dependence of water content in Wd and Rw.

Keywords: wadsleyite, ringwoodite, maximum water solubility, mantle transition zone

Temperature dependence of the $\text{Fe}^{3+} / \Sigma \text{Fe}$ ratio in wadsleyite by electron energy loss spectroscopy (EELS)

*Kazutaka YAMAGUCHI¹, Takaaki Kawazoe¹, Toru Inoue¹, Naotaka Tomioka²

1. Hiroshima University, 2. JAMSTEC, Kochi institute

Keywords: Wadsleyite, Ferric iron, Oxygen fugacity, Electron energy loss spectroscopy

Advantage on the analysis of cation site occupancies by neutron diffraction and some examples of application for high-pressure minerals

*Takuo OKUCHI¹

1. Kyoto Univ.

Keywords: Neutron diffracton

In situ lattice volume observation of davemaoite in the $\text{CaSiO}_3\text{-H}_2\text{O}$ system up to uppermost lower mantle conditions

*Goru Takaichi¹, Takayuki Ishii², Yu Nishihara¹, Kyoko Matsukage³, Yuji Higo⁴, Noiyoshi Tsujino⁴, Sho Kakizawa⁴

1. GRC, Ehime Univ., 2. IPM, Okayama Univ., 3. Teikyo University of Science, 4. JASRI

Keywords: Water transport, Nominally anhydrous minerals, Davemaoite, Mantle transition zone, Lower mantle

Elasticity Measurements of Hydrous SiO₂ post-stishovite and its implication to the seismic anomalies in the lower mantle

*Yuichiro MORI^{1,2}, Motohiko MURAKAMI², Takashi YOSHINO³, Hiroyuki KAGI¹

1. UTokyo. Sci., 2. ETHZ. D-ERDW., 3. Okayama Univ. IPM

Keywords: High-pressure experiments, Lower mantle, Water, Sound velocity measurements

Oral presentation | R7: Petrology, Mineralogy and Economic geology (Joint Session with Society of Resource Geology)

📅 Thu. Sep 12, 2024 2:00 PM - 5:30 PM JST | Thu. Sep 12, 2024 5:00 AM - 8:30 AM UTC | 🏢 ES Hall Higashiyama Campus

R7: Petrology, Mineralogy and Economic geology (Joint Session with Society of Resource Geology)

Chairperson: Norikatsu Akizawa (Atmosphere and Ocean Research Institute, University of Tokyo), Takuya Echigo (Akita University)

岩石学, 鉱物学, 鉱床学, 地球化学などの分野をはじめとして, 地球・惑星物質科学全般にわたる岩石及び鉱物に関する研究発表を広く募集する。地球構成物質についての多様な研究成果の発表の場となることを期待する。

2:00 PM - 2:25 PM JST | 5:00 AM - 5:25 AM UTC

[R7-01] The Frontier of Ilmenite Geochronology Using the Uranium-Lead Decay System

「招待講演」

*Kengo Ito¹, Sota Niki², Tsuyoshi Iizuka³, Takafumi Hirata¹ (1. The Univ. of Tokyo. GcRC, 2. Nagoya Univ. ISEE, 3. The Univ. of Tokyo. Sci. EPS)

2:25 PM - 2:40 PM JST | 5:25 AM - 5:40 AM UTC

[R7-02] Distribution of rhenium in hydrothermal deposits in Japan, based on LA-ICP-MS analysis

「発表賞エントリー」

*Mitsuki Ogasawara¹, Junichiro Ohta^{1,2}, Kazutaka Yasukawa¹, Kentaro Nakamura^{2,1}, Yasuhiro Kato^{1,2} (1. Univ. Tokyo. Eng, 2. CIT. ORCeNG)

2:40 PM - 2:55 PM JST | 5:40 AM - 5:55 AM UTC

[R7-03] Occurrences of the PGE bearing (Fe,Ni,Co)AsS solid solution in the Bushveld Complex, South Africa

「発表賞エントリー」

*Sakuya Kubota¹, Takuya Echigo¹, Yasushi Watanabe¹ (1. Akita Univ. Irs.)

2:55 PM - 3:10 PM JST | 5:55 AM - 6:10 AM UTC

[R7-04] Shiranuiite and placer deposit of platinum group minerals, Haraigawa, Misato machi, Kumamoto Prefecture, Japan

*Daisuke HAMANE¹, Takahiro Tanaka, Tadashi Shinmachi (1. The University of Tokyo)

3:10 PM - 3:25 PM JST | 6:10 AM - 6:25 AM UTC

[R7-05] Geochemical and Sr-Nd isotopic characteristics of the Aitutaki peridotite xenoliths from the Cook islands

*Masako YOSHIKAWA¹, Norikatsu Akizawa² (1. Hiroshima Univ., 2. Tokyo Univ. AORI)

3:25 PM - 3:40 PM JST | 6:25 AM - 6:40 AM UTC

[R7-06] Hydration of the mantle and magma genesis in the Izu-Bonin-Mariana fore-arc

*Rion Yamaoka¹, Norikatsu Akizawa¹, Yuji Ichiyama², ToTomoaki Morishita³, Akihiro Tamura³, Hiroyuki Yamashita⁴, Yumiko Harigane⁵, Yasuhiko Ohara^{6,7,8} (1. University of Tokyo, 2. Chiba Univ., 3. Kanazawa Univ., 4. Kanagawa Prefectural Museum of Natural History, 5. AIST, 6. Japan Coast Guard, Hydrographic and Oceanographic Department, 7. JAMSTEC, 8. Nagoya Univ.)

3:40 PM - 3:50 PM JST | 6:40 AM - 6:50 AM UTC

[1Lecture-108-19-7add] 休憩

3:50 PM - 4:15 PM JST | 6:50 AM - 7:15 AM UTC

[R7-07] Fault-rock mineralogy and weakening mechanisms at deep- to shallow-crustal depths of Main Himalayan Thrust, NW India

「招待講演」

*Dyuti Prakash SARKAR¹, Takehiro Hirose¹, Jun-ichi Ando², Kaushik Das², Gautam Ghosh³ (1. Japan Agency for Marine-Earth Science and Technology (JAMSTEC), 2. Hiroshima University, 3. Presidency University)

4:15 PM - 4:30 PM JST | 7:15 AM - 7:30 AM UTC

[R7-08] On the Relationship between Phosphorus Concentration and Microstructure in Goethite in Australian Iron Ore

*Takashi MIKOUCHI¹, Reiko Murao², Toru Takayama² (1. Univ. Museum, Univ. of Tokyo, 2. Nippon Steel)

4:30 PM - 4:45 PM JST | 7:30 AM - 7:45 AM UTC

[R7-09] Mineral zoning and formation temperatures around the Myokenzan lithium pegmatites

「発表賞エントリー」

*Takumi Ishizaki¹, Takuya Echigo¹, Yasushi Watanabe¹ (1. Akita Univ.)

4:45 PM - 5:00 PM JST | 7:45 AM - 8:00 AM UTC

[R7-10] Copper as a Coloring Factor of Pale Blue Sepiolite in amygdales in Higashimatsuura basalt from Kabeshima, Saga Prefecture, SW Japan

「発表賞エントリー」

*Satsuki KIO¹, Terumi Ezima², Yoshiaki Kon³, Hikari Minamisawa⁴, Masaomi Horita⁴ (1. Shinshu Univ. Sci., 2. Shinshu Univ. Sci., 3. GSJ, AIST, 4. Shinshu Univ. Eng.)

5:00 PM - 5:15 PM JST | 8:00 AM - 8:15 AM UTC

[R7-11] Quantification of Mineral Modal and Zoning Ratios by Powder X-ray Diffraction: Advancing Towards Big Data Era

「発表賞エントリー」

*Satoshi MATSUNO¹, Shuhou Maitani², Kenta Yoshida³, Nobuo Hirano¹, Ryosuke Sinmyo², Keiichi Osaka⁴, Shin-ichi Yamasaki¹, Masaoki Uno¹, Atsushi Okamoto¹ (1. Tohoku University, 2. Meiji University, 3. JAMSTEC, 4. Japan Synchrotron Radiation Research Institute)

5:15 PM - 5:30 PM JST | 8:15 AM - 8:30 AM UTC

[R7-12] Searching for Dark Matter Using Geological Samples: Techniques and Preliminary Results

*Natsue ABE^{1,2}, Shigenobu Hirose¹, Qing CHANG¹, Takeshi Hanyu¹, Noriko Hasebe², Yasushi Hoshino³, Takashi Kamiyama⁴, Yoji Kawamura¹, Kohta Murase⁵, Tatsuhiro Naka⁶, Kenji Oguni¹, Katsuhiko Suzuki¹, Seiko Yamazaki⁷ (1. JAMSTEC, 2. Kanazawa University, 3. Kanagawa University, 4. Hokkaido University, 5. Penn State University, 6. Toho University, 7. AIST)

The Frontier of Ilmenite Geochronology Using the Uranium-Lead Decay System

*Kengo Ito¹, Sota Niki², Tsuyoshi Iizuka³, Takafumi Hirata¹

1. The Univ. of Tokyo. GcRC, 2. Nagoya Univ. ISEE, 3. The Univ. of Tokyo. Sci. EPS

Keywords: Ilmenite, Geochronology, U-Pb dating, LA-ICP-MS

Distribution of rhenium in hydrothermal deposits in Japan, based on LA-ICP-MS analysis

*Mitsuki Ogasawara¹, Junichiro Ohta^{1,2}, Kazutaka Yasukawa¹, Kentaro Nakamura^{2,1}, Yasuhiro Kato^{1,2}

1. Univ. Tokyo. Eng, 2. CIT. ORCeNG

Keywords: Hydrothermal deposit, LA-ICP-MS, Re mapping, Re-Os radiometric dating

Occurrences of the PGE bearing (Fe,Ni,Co)AsS solid solution in the Bushveld Complex, South Africa

*Sakuya Kubota¹, Takuya Echigo¹, Yasushi Watanabe¹

1. Akita Univ. Irs.

Keywords: PGE mineralization, Bushveld Complex, Assimilation

Shiranuiite and placer deposit of platinum group minerals, Haraigawa, Misato machi, Kumamoto Prefecture, Japan

*Daisuke HAMANE¹, Takahiro Tanaka, Tadashi Shinmachi

1. The University of Tokyo

We have discovered a placer deposit of platinum-group minerals (PGM) in the clinopyroxene mass, Haraigawa, Misato machi, Kumamoto Prefecture. Two new minerals, minakawaite and michitoshiite-(Cu), were discovered in this deposit, and an unknown mineral of the thiospinel group was also found. It is named shiranuiite and was approved as a new mineral (IMA2023-072a). Most placer grains are isoferroplatinum: Pt_3Fe , and some grains have been altered at the outer edge (several to several hundred μm) to tulameenite: Pt_2CuFe or tetraferroplatinum: $PtFe$, showing the influence of post-magmatic alteration. Bowieite: Rh_2S_3 , the second most abundant inclusions, rarely appear on the surface of grains, while it is altered to often cuprorhodsite: $(Cu^{+}_{0.5}Fe^{3+}_{0.5})Rh^{3+}_2S_4$ and rarely shiranuiite: $Cu^{+}(Rh^{3+}Rh^{4+})S_4$. The empirical formula of shiranuiite is $(Cu^{+}_{0.95}Fe^{3+}_{0.04}Ni_{0.01})(Rh^{3+}_{1.19}Rh^{4+}_{0.77}Ir^{4+}_{0.06})S_{3.99}$ and the unit cell parameter is $a = 9.757 \text{ \AA}$ on Fd-3m space group.

Keywords: Platinum-group minerals (PGM), Placer PGM, Shiranuiite




図1. 砂白金粒子の表面に露出したパウイー鉱と不知火鉱 (硫銅ロジウム鉱) の分布 (a)、産出鉱物の一覧 (b)、不知火鉱 (硫銅ロジウム鉱) とRh-S鉱物 (パウイー鉱、キングストーン鉱、ミアス鉱) の組成分布 (c)。

Geochemical and Sr-Nd isotopic characteristics of the Aitutaki peridotite xenoliths from the Cook islands

*Masako YOSHIKAWA¹, Norikatsu Akizawa²

1. Hiroshima Univ., 2. Tokyo Univ. AORI

Keywords: Aitutaki island, mantle peridotite xenolith, Sr-Nd isotopic ratios

Hydration of the mantle and magma genesis in the Izu–Bonin–Mariana fore-arc

*Rion Yamaoka¹, Norikatsu Akizawa¹, Yuji Ichiyama², ToTomoaki Morishita³, Akihiro Tamura³,
Hiroyuki Yamashita⁴, Yumiko Harigane⁵, Yasuhiko Ohara^{6,7,8}

1. University of Tokyo, 2. Chiba Univ., 3. Kanazawa Univ., 4. Kanagawa Prefectural Museum of Natural History, 5. AIST,
6. Japan Coast Guard, Hydrographic and Oceanographic Department, 7. JAMSTEC, 8. Nagoya Univ.

Keywords: Subduction zone, Hydration, Fore-arc magma genesis, Peridotite, Amphibole

Oral presentation

R7: Petrology, Mineralogy and Economic geology (Joint Session with Society of Resource Geology)

Chairperson: Norikatsu Akizawa (Atmosphere and Ocean Research Institute, University of Tokyo), Takuya Echigo (Akita University)

Thu. Sep 12, 2024 2:00 PM - 5:30 PM ES Hall (Higashiyama Campus)

岩石学，鉱物学，鉱床学，地球化学などの分野をはじめとして，地球・惑星物質科学全般にわたる岩石及び鉱物に関する研究発表を広く募集する。地球構成物質についての多様な研究成果の発表の場となることを期待する。

3:40 PM - 3:50 PM

[1Lecture-108-19-7add]休憩

Fault-rock mineralogy and weakening mechanisms at deep- to shallow-crustal depths of Main Himalayan Thrust, NW India

*Dyuti Prakash SARKAR¹, Takehiro Hirose¹, Jun-ichi Ando², Kaushik Das², Gautam Ghosh³

1. Japan Agency for Marine-Earth Science and Technology (JAMSTEC), 2. Hiroshima University, 3. Presidency University

Fault rock mineralogy and deformation microstructures of crustal-scale faults provide insights into the stress accommodation mechanisms in terms of fault weakening and possibilities of earthquake nucleation. The Main Himalayan Thrust is considered to be the crustal-scale collisional boundary along which the Indian plate subducts beneath the Asian plate. The near-surface exposure of the present active boundary of the subduction is considered the Main Frontal Thrust, while the Main Central Thrust at the northernmost boundary represents the older post-collisional paleo-thrust boundaries between Indian and Asian plates. Here we present a comparative study of our findings on the fault rocks from the Nahan Thrust and North Almora Thrust belonging to the Main Frontal Thrust and the Main Central Thrust respectively. The deformation zone from North Almora Thrust represents the deep crustal part of the Main Himalayan Thrust exhibiting granite mylonite-ultramylonite zone. Quartz microstructures indicate deformation temperatures of 450–550 °C with evidence of grain-size sensitive creep as the dominant weakening mechanism. Additionally, two-feldspar thermometry has been used to estimate temperatures from sheared myrmekitic feldspar in the mylonites and ultramylonites. Crystallographic preferred orientations of quartz and mica indicate a shift in partitioning of strain from quartz to mica with an increase in mica content. In comparison, the fault zone from the Nahan Thrust shows the dominance of brittle deformation structures consisting of highly brecciated sandstone and a wide gouge zone. Microstructures of the gouge zone indicate distributed deformation and progressive stress localization leading to fracture development, cataclasis, and frictional sliding. Especially the ultra-fine bands of the principle slip zone exhibit a possible fingerprint of frictional heating during the seismic slips. The frictional properties of the fault rocks estimated from the rotary-shear velocity step experiments also indicate a velocity weakening to strengthening behavior based on phyllosilicate content. Our results on fault rocks from deep- to shallow-crustal depth indicate that phyllosilicate content play a pivotal role in frictional behavior and weakening mechanisms along the subduction boundary.

Keywords: Main Himalayan Thrust, Main Central Thrust, Main Frontal Thrust, rock friction, phyllosilicates

On the Relationship between Phosphorus Concentration and Microstructure in Goethite in Australian Iron Ore

*Takashi MIKOUCHI¹, Reiko Murao², Toru Takayama²

1. Univ. Museum, Univ. of Tokyo, 2. Nippon Steel

Keywords: Iron Ore, Goethite, Phosphorous, Microstructure

Mineral zoning and formation temperatures around the Myokenzan lithium pegmatites

*Takumi Ishizaki¹, Takuya Echigo¹, Yasushi Watanabe¹

1. Akita Univ.

Keywords: Pegmatite, Crystallization fractionation, Garnet-biotite geothermometer, Mineral zoning

Copper as a Coloring Factor of Pale Blue Sepiolite in amygdales in Higashimatsuura basalt from Kabeshima, Saga Prefecture, SW Japan

*Satsuki KIO¹, Terumi Ezima², Yoshiaki Kon³, Hikari Minamisawa⁴, Masaomi Horita⁴

1. Shinshu Univ. Sci., 2. Shinshu Univ. Sci., 3. GSJ, AIST, 4. Shinshu Univ. Eng.

Keywords: Kabeshima Island, sepiolite, copper, Higashimatsuura Basalt

Quantification of Mineral Modal and Zoning Ratios by Powder X-ray Diffraction: Advancing Towards Big Data Era

*Satoshi MATSUNO¹, Shuhou Maitani², Kenta Yoshida³, Nobuo Hirano¹, Ryosuke Sinmyo², Keiichi Osaka⁴, Shin-ichi Yamasaki¹, Masaoki Uno¹, Atsushi Okamoto¹

1. Tohoku University, 2. Meiji University, 3. JAMSTEC, 4. Japan Synchrotron Radiation Research Institute

Keywords: Powder X-ray Diffraction, Rietveld Refinement, Mineral Modal, Mineral Zoning, Quantification

Searching for Dark Matter Using Geological Samples: Techniques and Preliminary Results

*Natsue ABE^{1,2}, Shigenobu Hirose¹, Qing CHANG¹, Takeshi Hanyu¹, Noriko Hasebe², Yasushi Hoshino³, Takashi Kamiyama⁴, Yoji Kawamura¹, Kohta Murase⁵, Tatsuhiro Naka⁶, Kenji Oguni¹, Katsuhiko Suzuki¹, Seiko Yamazaki⁷

1. JAMSTEC, 2. Kanazawa University, 3. Kanagawa University, 4. Hokkaido University, 5. Penn State University, 6. Toho University, 7. AIST

The traces of cosmic rays from inside and outside the Earth recorded in geological samples can be effectively used for geological and astrophysical research. For instance, studies using Antarctic ice cores have estimated the number and age of supernova explosions and observed geoneutrinos from Earth's interior. Research utilizing seabed samples to measure cosmic ray intensity is also expected to provide insights into various events over long geological periods. Only about 5% of the matter in the universe can be directly observed, including such cosmic rays. In contrast, approximately 70% is dark energy, and the remaining 25% or more is dark matter. To search for such unknown matter or energy, large detectors using xenon are typically employed for dark matter detection. However, their scalability could be improved, making it challenging to improve detection limits. Considering this situation, we are exploring methods other than large detectors. For example, natural minerals like mica have been around for geological time scales, providing plenty of exposure even in small samples. These minerals can retain nuclear recoil tracks—evidence of dark matter interactions—for periods longer than the Earth's age. When etched, these tracks appear as observable pits. In 1995, Snowden-Ifft and colleagues studied natural Muscovite that was 500 million years old and covered an area of just 0.08 square millimeters. We propose using natural minerals (such as olivine and mica) that have long formation periods and are collected from deep within the ocean floor or continents with minimal surrounding radioactive substances. We are developing the necessary observational techniques to identify traces of dark matter as "Paleo-detectors" in these samples. Although there is still a long way to go before making an actual observation, this presentation will share our current efforts and discuss more efficient observation methods.

Keywords: Paleo-detectors, Dark Matter, Geoneutrino, Cosmic rays

Oral presentation | R1: Characterization and description of minerals (Joint Session with The Gemmological Society of Japan)

📅 Thu. Sep 12, 2024 2:00 PM - 3:15 PM JST | Thu. Sep 12, 2024 5:00 AM - 6:15 AM UTC | 🏠 ES024 Higashiyama Campus

R1: Characterization and description of minerals (Joint Session with The Gemmological Society of Japan)

Chairperson: Koichi Momma, Yohei Shirose

2:00 PM - 2:15 PM JST | 5:00 AM - 5:15 AM UTC

[R1-08] Ultra nano inclusion in blue sapphire from Diego, Madagascar

*Akira MIYAKE¹, Seika Oto¹, Yohei Igami¹, Kentaro Emori² (1. Kyoto University, 2. CGL)

2:15 PM - 2:30 PM JST | 5:15 AM - 5:30 AM UTC

[R1-09] Reexamination of the symmetry of itoigawaite

*Chigako KIYOHARA, Yohei Igami¹, Akira Miyake¹, Norimasa Shimobayashi¹ (1. Kyoto Univ. Sci.)

2:30 PM - 2:45 PM JST | 5:30 AM - 5:45 AM UTC

[R1-10] Zeolite and altered boninite from Miyanojima, Ogasawara, Japan

*Yuki INOUE¹, Mana Yasui², Jun-ichiro Ishibashi³, Hagiya Hiroshi⁴, Kazuhiko Shimada¹, Athushi Yamazaki² (1. Kyushu Univ. Sci., 2. Waseda Univ., 3. Kobe Univ., 4. Tokyo City Univ.)

2:45 PM - 3:00 PM JST | 5:45 AM - 6:00 AM UTC

[R1-11] Willemseite from the Hamayokawa mine, Nagano Prefecture, Japan

*Satoshi Matsubara¹, Koichi Momma¹, Norio Yanagisawa², Yoko Kusaba¹, Akiko Tokumoto¹, Ritsuro Miyawaki¹ (1. National Museum of Nature and Science, 2. Geological Survey of Japan, AIST)

3:00 PM - 3:15 PM JST | 6:00 AM - 6:15 AM UTC

[R1-12] On the Fe²⁺-analogue of zemannite from Kawazu mine, Shizuoka Prefecture, Japan

*Koichi MOMMA¹, Owen Missen², Stuart Mills³, Ritsuro Miyawaki¹, Satoshi Matsubara¹, Eiji Ohtani⁴, Seiji Kamada⁵, Shin Ozawa⁴ (1. National Museum of Nature and Science, 2. University of Tasmania, 3. The Arkenstone, 4. Tohoku Univ., 5. AD Science Inc.)

Ultra nano inclusion in blue sapphire from Diego, Madagascar

*Akira MIYAKE¹, Seika Oto¹, Yohei Igami¹, Kentaro Emori²

1. Kyoto University, 2. CGL

Ultra-nano inclusions in Be-containing natural blue sapphires (corundum) from Diego, Madagascar were observed and analysed by transmission electron microscope (TEM) and 3D atomprobe (3DAP). TEM results show that ultra-nano inclusion has 10 nm length and 2 nm width, and the nano-inclusion is identified by Wolframite structure (P2/c). 3DAP results show the inclusion consists of Ti, Nb, Ta, Fe, Sn.


Keywords: Ultra-nano inclusion, corundum)

Reexamination of the symmetry of itoigawaite

*Chigako KIYOHARA, Yohei Igami¹, Akira Miyake¹, Norimasa Shimobayashi¹

1. Kyoto Univ. Sci.

Keywords: Itoigawaite, symmetry



図：糸魚川石中の細かい筋状組織のTEM明視野像とその部分から得られた電子回折図形。端の方のスポットは2つに分裂しており、そのため結晶格子の軸角が 90° からわずかにずれていると考えられる。

Zeolite and altered boninite from Miyanohama, Ogasawara, Japan

*Yuki INOUE¹, Mana Yasui², Jun-ichiro Ishibashi³, Hagiya Hiroshi⁴, Kazuhiko Shimada¹, Athushi Yamazaki²

1. Kyushu Univ. Sci., 2. Waseda Univ. , 3. Kobe Univ., 4. Tokyo City Univ.

Keywords: zeolite, Boninite, phillipsite


Willemseite from the Hamayokawa mine, Nagano Prefecture, Japan

*Satoshi Matsubara¹, Koichi Momma¹, Norio Yanagisawa², Yoko Kusaba¹, Akiko Tokumoto¹, Ritsuro Miyawaki¹

1. National Museum of Nature and Science, 2. Geological Survey of Japan, AIST

Rare mineral, suzukiite, was reported from the metamorphosed manganese ore deposit of the Hamayokawa mine (Hirowatari & Yoshie, 1978). Although some specimens from the Hamayokawa mine, Nagano Prefecture, Japan donated by amateur mineralogists in the National Museum of Nature and Science and the Geological Survey of Japan are registered as suzukiite, but recent our analyses reveal they are not suzukiite but willemseite $[(\text{Ni}_3\text{Si}_4\text{O}_{10}(\text{OH})_2)]$ which is first found in Japan.

Keywords: suzukiite, willemseite, Hamayokawa mine



On the Fe²⁺-analogue of zemannite from Kawazu mine, Shizuoka Prefecture, Japan

*Koichi MOMMA¹, Owen Missen², Stuart Mills³, Ritsuro Miyawaki¹, Satoshi Matsubara¹, Eiji Ohtani⁴, Seiji Kamada⁵, Shin Ozawa⁴

1. National Museum of Nature and Science, 2. University of Tasmania, 3. The Arkenstone, 4. Tohoku Univ., 5. AD Science Inc.

The Kawazu Mine in Shimoda City, Shizuoka Prefecture, has been known as type locality of two new minerals, kawazulite (Bi₂Te₂Se) and kinichilite. In the original description of kinichilite, its chemical composition was reported as (Fe²⁺_{1.13}Mg_{0.47}Zn_{0.43}Mn²⁺_{0.17})_{Σ2.20}(Te_{2.97}Se_{0.03})_{Σ3.00}O_{9.00}(H_{1.38}Na_{0.22})_{Σ1.60} · 3.2H₂O, and as Fe analogue of zemannite at that time. Later, however, crystal chemical study of zemannite revealed that its ideal formula is Mg_{0.5}[ZnFe³⁺(TeO₃)₃] · 4.5H₂O, *i.e.*, iron is not 2+ but 3+ and both Zn²⁺ and Fe³⁺ are essential. If the originally reported chemical composition of kinichilite is recalculated based on this find, it remains in the compositional range of zemannite. However, R. Miletich (1995), in his study of zemannite group minerals, also examined the "kinichilite" from the Kawazu mine and found that manganese was predominant as a divalent ion in most of the analyzed point. He considered it as kinichilite and redefined its ideal formula as Mg_{0.5}[MnFe³⁺(TeO₃)₃] · 4.5H₂O. Here, we report Fe analogue of zemannite, in which divalent iron exceeds zinc, manganese and other divalent ions. One of the collections of the National Museum of Nature and Science (NSM-M41022) was used in this study. Chemical analysis was performed using a scanning electron microscope (JEOL JSM-6610) equipped with an energy dispersive X-ray spectroscopic detector (EDS). Water content was estimated by the difference from 100% of the EDS analysis. Single crystal X-ray diffraction (SXR) experiments were performed using a Rigaku Synergy Custom equipped with a rotating anode and a multilayer X-ray focusing mirror (VariMax).

The empirical formula obtained from the average of four analytical points is Mg_{0.40}(Fe²⁺_{0.54}Zn_{0.33}Cu_{0.18}Mn_{0.05})_{Σ1.10}Fe³⁺_{1.00}(Te_{2.92}P_{0.06}O₉) · 4.12H₂O. The SXR experiment gave the space group *P6₃* (or *P6₃/m*) with lattice parameters *a* = 9.37765(11), *c* = 7.58379(9), *V* = 577.570(15). The refinement converged with reliability index *R*₁ = 1.56% and it was confirmed to be isostructural with zemannite. Distribution of the channel cations and bond lengths of the framework *M* sites indicate symmetry lowering from *P6₃/m* to *P6₃*.

Keywords: zemannite, kinichilite, Kawazu mine

Oral presentation | R3: High-pressure science and deep Earth's material

📅 Thu. Sep 12, 2024 2:00 PM - 3:15 PM JST | Thu. Sep 12, 2024 5:00 AM - 6:15 AM UTC | 🏠 ES025
Higashiyama Campus

R3: High-pressure science and deep Earth's material

Chairperson: Takeshi Sakai (Ehime University), Ryosuke Sinmyo (Meiji University), Takayuki Ishii (Okayama University), Takaaki Kawazoe (Hiroshima University)

2:00 PM - 2:15 PM JST | 5:00 AM - 5:15 AM UTC

[R3-09] Electrical conductivity of the lower mantle materials under high pressure with implication for valence and spin states of iron

「招待講演」

*Izumi MASHINO¹ (1. Okayama Univ.)

2:15 PM - 2:30 PM JST | 5:15 AM - 5:30 AM UTC

[R3-10] GHz-DAC ultrasonics to measure elastic anomaly with the spin transition of ferrous ion in ferropericlase

*Akira Yoneda¹, Izumi Mashino², Ryosuke Matsui¹, Ryoma Ishida¹, Tadashi Kondo¹ (1. Osaka Univ. Sci, 2. Okayama Univ. IPM)

2:30 PM - 2:45 PM JST | 5:30 AM - 5:45 AM UTC

[R3-11] Reaction between water and iron in the early magma ocean and the present core-mantle boundary

*Yongjae Lee¹ (1. Yonsei University)

2:45 PM - 3:00 PM JST | 5:45 AM - 6:00 AM UTC

[R3-12] Sound velocity of B2-FeNiSi alloy at high pressure and high temperature and constitution of the Earth's inner core

*Eiji OHTANI¹, Daijo IKUTA², Hiroshi FUKUI^{3,4}, Tatsuya SAKAMAKI¹, Daisuke Ishikawa^{3,4}, Alfred Q. R. BARON^{3,4} (1. Tohoku University, 2. Institute for Planetary Materials, Okayama University, 3. JASRI, 4. RIKEN)

3:00 PM - 3:15 PM JST | 6:00 AM - 6:15 AM UTC

[R3-13] On the phase boundary between FeS IV and V

*Satoru URAKAWA¹ (1. Okayama University)

Electrical conductivity of the lower mantle materials under high pressure with implication for valence and spin states of iron

*Izumi MASHINO¹

1. Okayama Univ.


Keywords: Spin transition, The lower mantle, Electrical conductivity

GHz-DAC ultrasonics to measure elastic anomaly with the spin transition of ferrous ion in ferroperricite

*Akira Yoneda¹, Izumi Mashino², Ryosuke Matsui¹, Ryoma Ishida¹, Tadashi Kondo¹

1. Osaka Univ. Sci, 2. Okayama Univ. IPM

Keywords: GHz-DAC ultrasonics, Spin transition, ferroperricite



Reaction between water and iron in the early magma ocean and the present core-mantle boundary

*Yongjae Lee¹

1. Yonsei University

Recent interest in hydrogen as a clean geological resource has been escalating, leading to active research on the formation and exploration of natural hydrogen worldwide. Paradoxically, however, hydrogen is the most abundant element in the universe and is also widely distributed in the Earth, as contained in mineral structures from the crust to the core. Hydrogen in minerals can exist in the form of molecular water (H₂O), hydroxyl group (OH), or monatomic hydrogen (H), and is, therefore, directly related to the global transport and distribution of water. From this perspective, it is notable that the hydrogen budget increases innards to the mantle transition zone, lower mantle, and core, compared to the amount in the hydrosphere on the Earth's surface. In this presentation, I will introduce the reactions between water and iron based on two recent experimental results to simulate the conditions of the early magma ocean [1] and the present core-mantle boundary [2], which could be linked to the initial distribution of hydrogen and its enrichment in the topmost outer core.

[1] J. Choi, R.J. Husband, H. Hwang, T. Kim, Y. Bang, S. Yun, J. Lee, H. Sim, S. Kim, D. Nam, B. Chae, H.-P. Liermann, Y. Lee*, Oxidation of iron by giant impact and its implication on the formation of reduced atmosphere in the early Earth, *Science Advances*, Vol.9, eadi6096, 2023

[2] T. Kim, J.G. O'Rourke, J. Lee, S. Chariton, V. Prakapenka, R.J. Husband, N. Giordano, H.-P. Liermann, S.-H. Shim*, Y. Lee*, "A hydrogen-enriched layer in the topmost outer core sourced from deeply subducted water", *Nature Geoscience*, Vol.16, 1208-1214, 2023

Keywords: Hydrogen, Magma ocean, Core-mantle boundary

Sound velocity of B2-FeNiSi alloy at high pressure and high temperature and constitution of the Earth's inner core

*Eiji OHTANI¹, Daijo IKUTA², Hiroshi FUKUI^{3,4}, Tatsuya SAKAMAKI¹, Daisuke Ishikawa^{3,4}, Alfred Q. R. BARON^{3,4}

1. Tohoku University, 2. Institute for Planetary Materials, Okayama University, 3. JASRI, 4. RIKEN

Elastic properties of an ordered derivative of the body-centered cubic (B2) structure of Fe-7wt% Ni-15wt% Si ($\text{Fe}_{0.67}\text{Ni}_{0.06}\text{Si}_{0.27}$) alloy have been investigated by combining high-resolution inelastic X-ray scattering and powder X-ray diffraction in diamond anvil cells up to 130 GPa and 2300 K. The density (ρ)-compressional wave velocity (V_p) relation of this phase shows a weak or almost negligible temperature dependence, similar to that observed for the body-centered cubic phase of pure iron (Shibazaki et al., 2016) and the B20 phase of iron-silicon alloys (Whitaker et al., 2009). The ρ , V_p and shear wave velocity (V_s) were extrapolated to the inner core conditions and compared with the PREM (preliminary reference Earth model) inner core (Dziewonski and Anderson, 1981). The phase relation of the Fe-Ni-Si system revealed that B2-Fe-7wt% Ni-15wt% Si alloy coexists with hexagonal close-packed (hcp) Fe-Ni alloys with negligible amount of silicon (Ikuta et al., 2021). The V_p and V_s of the two-phase mixture of B2 and hcp phases under the inner core conditions show slightly higher V_p and V_s compared to the PREM inner core. The two-phase mixture with the addition of a small amount of sulfur could potentially explain the properties of the PREM inner core.

Keywords: Sound velocity, B2-FeNiSi alloy, High pressure and high temperature, Inner core

On the phase boundary between FeS IV and V

*Satoru URAKAWA¹

1. Okayama University

Keywords: iron sulfide, second order phase transition, thermal expansivity, high pressure, X-ray diffraction

Oral presentation | R8: Metamorphic rocks and tectonics

📅 Thu. Sep 12, 2024 3:30 PM - 6:00 PM JST | Thu. Sep 12, 2024 6:30 AM - 9:00 AM UTC | 🏠 ES024
Higashiyama Campus

R8: Metamorphic rocks and tectonics

Chairperson: Yui Kouketsu (Nagoya University), Shunsuke Endo

3:30 PM - 3:45 PM JST | 6:30 AM - 6:45 AM UTC

[R8-01] Toward the establishment of "EBSD method for the detection of quartz with phase transition to coesite": Incorporation of the Gongen area, Sanbagawa metamorphic belt, Shikoku, SW-Japan

「発表賞エントリー」

*Momoko Minowa¹, Takayoshi Nagaya¹, Taisuke Ito², Simon Wallis² (1. Tokyo Gakugei University, 2. The University of Tokyo)

3:45 PM - 4:00 PM JST | 6:45 AM - 7:00 AM UTC

[R8-02] Pressure-Temperature-Time Conditions of The Basement Rocks in The Banggai-Sula Microcontinent, Indonesia

「発表賞エントリー」

*Fransiska Ayuni Catur Wahyuandari¹, Nobuhiko Nakano¹, Tatsuro Adachi¹, Nugroho Imam Setiawan² (1. Kyushu Univ., 2. Gadjah Mada Univ.)

4:00 PM - 4:15 PM JST | 7:00 AM - 7:15 AM UTC

[R8-03] Amphibolites from the Mineoka belt revisited

*Yuji ICHIMURA¹, Shun Takamizawa¹, Hisatoshi Ito², Akihiro Tamura³, Tomoaki Morishita³ (1. Chiba University, 2. Central Research Institute of Electric Power Industry, 3. Kanazawa University)

4:15 PM - 4:30 PM JST | 7:15 AM - 7:30 AM UTC

[R8-04] Compositional heterogeneity of garnet in garnet-biotite felsic granulite, Czech Republic

*Daisuke NAKAMURA¹, Yuuka FUJIMURA¹, Yuuki KODERA¹ (1. Okayama University)

4:30 PM - 4:45 PM JST | 7:30 AM - 7:45 AM UTC

[R8-05] Formation and alteration of a zoned calcsilicate vein from the contact aureole of Kasuga area, central Japan: insights from chemical ages of U-Th oxides

*Shunsuke ENDO¹ (1. Shimane University)

4:45 PM - 5:00 PM JST | 7:45 AM - 8:00 AM UTC

[R8-06] Observations of geological structures and microstructures associated with strain release at a continental plate boundary fault

「発表賞エントリー」

*Haruki Yoshiasa¹, Jun-ichi Ando^{1,2}, Kaushik Das^{1,2}, Dyuti Prakash Sarkar³ (1. Hiroshima University, 2. HiPeR, 3. JAMSTEC)

5:00 PM - 5:15 PM JST | 8:00 AM - 8:15 AM UTC

[R8-07] Comparison and verification study of new sample forms for use in Raman carbonaceous material geothermometer

「発表賞エントリー」

*Shunsuke Ogino¹, Yui Kouketsu¹, Satoshi Takahashi¹ (1. Nagoya Univ. Env.)

5:15 PM - 5:30 PM JST | 8:15 AM - 8:30 AM UTC

[R8-08] Pressure and temperature conditions for the occurrence of index minerals in pelitic schists in the Sanbagawa belt, Kanto Mountains, Japan, inferred from Raman carbonaceous material geothermometry

*Yui KOUKETSU¹, Ichiko Shimizu² (1. Nagoya Univ. Env., 2. Kyoto Univ. Sci.)

5:30 PM - 5:45 PM JST | 8:30 AM - 8:45 AM UTC

[R8-09] Microstructural and petrological characteristics of the Ryoke mylonite occurred along the Median Tectonic Line in the Shinshiro area, Aichi Prefecture

*Miharu Niwa^{1,2}, Katsuyoshi Michibayashi^{1,3}, Kenichiro Tani⁴, Takuma Nishimura⁵ (1. Nagoya Univ. Env., 2. TMNH, 3. JAMSTEC, 4. NMNS, 5. HMNH)

5:45 PM - 6:00 PM JST | 8:45 AM - 9:00 AM UTC

[R8-10] Re-proposal of the metamorphic zone division in the Horokanai area of the Kamuikotan Belt based on the mineral paragenesis and phengite K-Ar Age.

*Takao HIRAJIMA¹, Kousuke Naemura², Kenta Yoshida³ (1. Kyoto University, 2. Iwate University, 3. JAMSTEC)

Toward the establishment of "EBSD method for the detection of quartz with phase transition to coesite": Incorporation of the Gongen area, Sanbagawa metamorphic belt, Shikoku, SW-Japan

*Momoko Minowa¹, Takayoshi Nagaya¹, Taisuke Ito², Simon Wallis²

1. Tokyo Gakugei University, 2. The University of Tokyo

Keywords: Sanbagawa metamorphic belt, Gongen area, coesite-quartz phase transition, EBSD (Electron Back Scatter Diffraction), anisotropy

Pressure-Temperature-Time Conditions of The Basement Rocks in The Banggai-Sula Microcontinent, Indonesia

*Fransiska Ayuni Catur Wahyuandari¹, Nobuhiko Nakano¹, Tatsuro Adachi¹, Nugroho Imam Setiawan²

1. Kyushu Univ., 2. Gadjah Mada Univ.

The Banggai-Sula microcontinent in the eastern Indonesia region is known to be derived from the Northwest Australia continental margin (Sula Spur). The basement rocks of Banggai-Sula microcontinents are often associated with the high-temperature/low-pressure metamorphic rocks of Kemum Basement High of the Bird's Head region in Papua which intruded by granitoids of Devonian–Carboniferous and Permian–Triassic age. This study reports the petrological and geochemical characteristics, and monazite EPMA ages of these basement rocks to understand its origin and the relation with tectonic events along the northeastern margin of Gondwana in the Paleozoic. Metamorphic rocks collected from Banggai and Peleng Islands imply differences in metamorphic conditions between Banggai Island in the east (garnet-andalusite-muscovite schist) and the western part of Peleng Island (garnet-staurolite-biotite-muscovite schist). The preservation of garnet growth zoning during prograde metamorphism is indicated by decreasing Mn from the core towards the rim of the garnet grains obtained from metamorphic rocks in this region. The monazite EPMA ages were determined from biotite granite in Banggai Island and staurolite-bearing pelitic schist in Peleng Island. The biotite granite from Banggai Island gives a weighted mean age of 271 ± 8 and 263 ± 14 Ma from the core and rim of monazite. The staurolite-bearing pelitic schist from Peleng Island yields a weighted mean age of 434 ± 3 and 449 ± 12 Ma from the core and mantle, and 403 ± 8 Ma from the rim of monazite. This Silurian-Devonian metamorphic age has not been reported before from the Banggai-Sula microcontinent and Eastern Indonesia region. The differences in metamorphic grade and timing with the Kemum Basement High suggest that an earlier tectonic event influenced the metamorphism in the Banggai-Sula microcontinent. Such early Paleozoic igneous activity and metamorphism happened along the northern margin of East Gondwana related to the subduction of the Proto-Tethys Oceanic crust. Silurian–Devonian metamorphism in the Banggai-Sula microcontinent might represent the eastern continuation of this tectonic event. In the presentation, we will also discuss the pressure-temperature conditions of these basement rocks.

Keywords: Banggai-Sula microcontinent, Paleozoic basement, monazite EPMA ages

Amphibolites from the Mineoka belt revisited

*Yuji ICHIYAMA¹, Shun Takamizawa¹, Hisatoshi Ito², Akihiro Tamura³, Tomoaki Morishita³

1. Chiba University, 2. Central Research Institute of Electric Power Industry, 3. Kanazawa University


Keywords: Mineoka belt, Amphibolites, Metamorphic sole

Compositional heterogeneity of garnet in garnet-biotite felsic granulite, Czech Republic

*Daisuke NAKAMURA¹, Yuuka FUJIMURA¹, Yuuki KODERA¹

1. Okayama University

Keywords: Garnet



Formation and alteration of a zoned calcsilicate vein from the contact aureole of Kasuga area, central Japan: insights from chemical ages of U-Th oxides

*Shunsuke ENDO¹

1. Shimane University

Keywords: Vein

Observations of geological structures and microstructures associated with strain release at a continental plate boundary fault

*Haruki Yoshiasa¹, Jun-ichi Ando^{1,2}, Kaushik Das^{1,2}, Dyuti Prakash Sarkar³

1. Hiroshima University, 2. HiPeR, 3. JAMSTEC

Keywords: Bedding plane slip, MBT, Microstructure, Frictional heat, Dynamic recrystallization

Comparison and verification study of new sample forms for use in Raman carbonaceous material geothermometer

*Shunsuke Ogino¹, Yui Kouketsu¹, Satoshi Takahashi¹

1. Nagoya Univ. Env.

Keywords: Carbonaceous material, Raman spectroscopy, Raman CM geothermometer, Mudstone, Pelitic schist

Pressure and temperature conditions for the occurrence of index minerals in pelitic schists in the Sanbagawa belt, Kanto Mountains, Japan, inferred from Raman carbonaceous material geothermometry

*Yui KOUKETSU¹, Ichiko Shimizu²

1. Nagoya Univ. Env., 2. Kyoto Univ. Sci.

The Sanbagawa belt in the Kanto Mountains has long been studied as a type locality for the Sanbagawa metamorphic belt, but the occurrence of index minerals and the graphitization degree (GD) show a complicated distribution, and the boundaries of isograd vary among researchers. In this study, we applied Raman carbonaceous material (CM) geothermometry to pelitic and siliceous schists collected in the Ayukawa-Sanbagawa area to estimate their temperature structures, and conducted pseudosection modelling to analyze the stability field of index minerals, such as garnet, biotite, and oligoclase. The application of Raman CM geothermometer to the Ayukawa-Sanbagawa area resulted in temperatures ranging from 360°C to 520°C: Chlorite zone is about 400-440°C, Garnet zone is about 360-470°C, and Biotite zone is about 470-520°C, respectively. A pseudosection diagram of the pelitic schist of the Biotite zone (AM41P, Miyashita 1997), which shows the average bulk rock composition of the Kanto Sanbagawa belt, shows that garnet, biotite, and oligoclase occur with increasing temperature. The stability field of garnet strongly affected by the bulk rock composition of MnO, as shown in previous studies, which explains the overlap of the temperature ranges of the Chlorite and Garnet zones. Compared with the temperature data, the biotite zone in the Kanto Mountains was formed at about 0.8 GPa, and may have experienced lower pressure conditions than the Shikoku. Oligoclase ($X_{An} > 0.1$) appears at about +20-30°C after the appearance of biotite under low pressure conditions up to about 0.9 GPa, but the stability field tends to shift to the higher temperature side under high pressure conditions above 0.9 GPa. This result is consistent with the hypothesis that there is a pressure difference between the Kanto Mountains, where the oligoclase appears homogenously in the biotite zone, and the Shikoku, where the Albite-biotite zone and the Oligoclase-biotite zone are clearly separated.

Keywords: Raman carbonaceous material geothermometry, Pseudosection modelling, Garnet, Biotite, Oligoclase

Microstructural and petrological characteristics of the Ryoke mylonite occurred along the Median Tectonic Line in the Shinshiro area, Aichi Prefecture

*Miharu Niwa^{1,2}, Katsuyoshi Michibayashi^{1,3}, Kenichiro Tani⁴, Takuma Nishimura⁵

1. Nagoya Univ. Env., 2. TMNH, 3. JAMSTEC, 4. NMNS, 5. HMNH

The Median Tectonic Line (MTL) in Shinshiro City, Higashimikawa area, Aichi Prefecture, has a complex geology, with the Ryoke metamorphic rocks and plutonic rocks in the inner zone, the Sambagawa metamorphic rocks in the outer zone, and the Shidara Group that overlie the basement rocks. Although the fault rocks are exposed in this area along the MTL as like in Nagano, Shizuoka and Mie Prefecture, detail structural and petrological reports are rare compared to other areas. The fault rocks in this area are mostly cataclasite, and mylonite has been found near in the Horai-cho, Shinshiro City, we found continuous outcrops of mylonite in the Sakurabuchi Park, Shinshiro City, further southwest from Hoai-cho. In this study, we observed microstructure with polarized light microscope, measured quartz crystal orientation with SEM-EBSD system, analyzed geochronological, in order to estimate deformation temperature, shear sense, and original rocks. Mylonite in this area is greenish-white in outcrop, and is characterized plagioclase porphyroclasts with white spots, as like Kashio mylonite. There are tonalitic rocks, composing of main constituently minerals, quartz, plagioclase, and biotite, with minor amount of K-feldspar, allanite, and zircon. The microstructure consists of a typical porphyroclastic texture composing of coarse-grained plagioclase and an asymmetric texture with plagioclase and fine quartz strain shadows. The quartz c-axis fabric shows a pattern of Y-maxima, which is formed at relatively intermediate temperature during deformation. In combination with the plagioclase-quartz porphyroclasts structure, which shows deformation during the progressive retrogression period, it is inferred that the quartz was formed by plastic shear deformation during the peak temperature from 350-450°C. Furthermore, the shear sense of the mylonite along the MTL, which is common in Nagano Prefecture, shows sinistral shear sense, whereas the shear sense of the mylonite in this area shows dextral shear sense. Based on these results, we discuss the regional structural characteristics of the Mikawa area in comparison with other areas.

Keywords: Mylonite, Median Tectonic Line, Ryoke belt, Fault rock

Re-proposal of the metamorphic zone division in the Horokanai area of the Kamuikotan Belt based on the mineral paragenesis and phengite K-Ar Age.

*Takao HIRAJIMA¹, Kousuke Naemura², Kenta Yoshida³

1. Kyoto University, 2. Iwate University, 3. JAMSTEC

According to the K-Ar age of phengite (Phn) and the deformation style of the rock, Sakakibara et al. (2007) proposed a zone division in the Horokanai area of the Kamuikotan belt, such as Horokanai unit (HKU) characterized by the occurrence of blueschist (BS) with Phn K-Ar ages of 135-120 Ma, and Biei unit (BIU) by the missing of BS and Phn K-Ar ages of 115-100 Ma. However, we confirmed the occurrence of BS from Mt. Shirakke through Jyari River to Mt. Numaushi, where Sakakibara et al. (2007) classified as BIU, in addition, the Phn K-Ar dating in the area has not been reported. To clarify these inconsistencies, we performed Phn K-Ar dating collected from the Jyari River and Mt. Numashi and their surrounding area. The results are as follows: HKU: EP256: Etanbetsu Pass, 115.1 Ma, KD12/KD13: Numaushi River East, 123.3/106.6 Ma, 126.9/101.8 Ma. BIU: KHK121: Mt. Numaushi, 120.2 Ma, KHK331/KHK333: Jyari-River, 109.2 Ma/105.6 Ma. Since Phns in the studied BS are generally fine-grained, the particle size of the dated fraction was adjusted to 0.2-2 micron and coarse-grained (50-120 micron) Phn fractions in KD12/KD13 were also dated. The coarse-grained fraction of KD12/KD13 showed 123.3/126.9 Ma, and their fine-grained fractions were 106.6/101.8 Ma, and the difference reached 12-25 Ma. The phenomenon of younger fine-grained fractions has also been confirmed in a few reports (Kurosegawa, Sato et al., 2014; Kanto Mountains, Lu et al., 2022), and there are various theories as to why. In order to eliminate the influence of particle size, the fine-grained fraction age is considered, i.e., 115.1~106.6 Ma for HKU and 120.2~105.6 Ma for BIU. Phn K-Ar ages (107-102 Ma) reported from BS missing area of BIU also overlap with abovementioned BS ages (120-101 Ma). Therefore, the blueschist occurrence region, which Sakakibara et al. (2007) regarded as the Biei unit, has lost the reason to distinguish it from the Horokanai unit in terms of mineral combination and metamorphic age.

Keywords: Phengite K-Ar age, Triple Point blueschist, Horokanai area, Kamuikotan Belt

Oral presentation | S3: Rheology and Material Transfer in Mantle and Crust (Special Session)

📅 Thu. Sep 12, 2024 3:30 PM - 6:00 PM JST | Thu. Sep 12, 2024 6:30 AM - 9:00 AM UTC | 📍 ES025 Higashiyama Campus

S3: Rheology and Material Transfer in Mantle and Crust (Special Session)

Chairperson: Ikuo Katayama (Hiroshima University), Katsuyoshi Michibayashi (Nagoya University)

3:30 PM - 3:50 PM JST | 6:30 AM - 6:50 AM UTC

[S3-01] High-speed time-resolved in-situ stress-strain measurements under high pressure and high temperature using synchrotron radiation X-rays

「招待講演」

*Noriyosi TSUJINO¹ (1. JASRI)

3:50 PM - 4:05 PM JST | 6:50 AM - 7:05 AM UTC

[S3-02] Deformation-induced crystallographic-preferred orientation of ϵ -FeOOH

*Yu NISHIHARA¹, Yui MORI¹, Wentian WU¹, Noriyoshi TSUJINO² (1. GRC, Ehime Univ., 2. JASRI)

4:05 PM - 4:20 PM JST | 7:05 AM - 7:20 AM UTC

[S3-03] Strength of metastable olivine and the rheology of subducting cold slab at mantle transition zone

「発表賞エントリー」

*Rikuto HONDA¹, Tomoaki Kubo¹, Noriyoshi Tsujino², Yuji Higo², Sho Kakizawa², Yuki Shibazaki³, Yu Nishihara⁴ (1. Kyushu University, 2. JASRI, 3. KEK, 4. Ehime Univ. GRC)

4:20 PM - 4:40 PM JST | 7:20 AM - 7:40 AM UTC

[S3-04] High-resolution measurement of ice anelasticity over a broad frequency range with a new cryogenic forced oscillation apparatus

「招待講演」

*Hatsuki Yamauchi¹, Christine McCarthy¹, Benjamin Holtzman¹, David Goldsby², Travis Hager² (1. Columbia Univ. LDEO, 2. U. Penn)

4:40 PM - 4:55 PM JST | 7:40 AM - 7:55 AM UTC

[S3-05] Relationship between creep and grain growth in rock with bicontinuous structure

「発表賞エントリー」

*Shenghao Jiang¹, Takehiko Hiraga¹ (1. The University of Tokyo)

4:55 PM - 5:10 PM JST | 7:55 AM - 8:10 AM UTC

[S3-06] Microstructures of plagioclase in gabbroic ultramylonites and quadruple point analysis

*Itsuki Natsume¹, Katsuyoshi Michibayashi^{2,3}, Yohei Igami⁴ (1. Kanagawa Prefectural Museum of Natural History, 2. Nagoya Univ., 3. JAMSTEC, 4. Kyoto Univ.)

5:10 PM - 5:25 PM JST | 8:10 AM - 8:25 AM UTC

[S3-07] Reconstruction of the uppermost mantle continuous structure of the Oman ophiolite

*Takeo Okuwaki¹, Natsume Itsuki², Katsuyoshi Michibayashi¹ (1. Nagoya Univ. Env, 2. Kanagawa Pref. Mus. Nat. Hist.)

5:25 PM - 5:40 PM JST | 8:25 AM - 8:40 AM UTC

[S3-08] Deformation and melt-rock interaction in the Horoman peridotite: Petrological and structural study of the MHL suite and BDH suite rocks

「発表賞エントリー」

*Aya Hihara¹, Miki Tasaka¹, Keisuke Kurihara¹, Hajime Taniuchi², Tastuhiko Kawamoto¹ (1. Shizuoka Univ., 2. AIST)

Session

2024 Annual Meeting of Japan Association of Mineralogical Sciences (JAMS)

5:40 PM - 5:55 PM JST | 8:40 AM - 8:55 AM UTC

[S3-09] Multicomponent measurements of seismic velocity and electrical resistivity using foliated serpentinite and peridotite

「発表賞エントリー」

*Tomohiro Ito¹, Ikuo Katayama¹, Katsuyoshi Michibayashi², Kazuki Matsuyama² (1. Hiroshima University, 2. Nagoya University)

5:55 PM - 6:00 PM JST | 8:55 AM - 9:00 AM UTC

調整

High-speed time-resolved in-situ stress-strain measurements under high pressure and high temperature using synchrotron radiation X-rays

*Noriyosi TSUJINO¹

1. JASRI

Keywords: High pressure, High-speed time-resolution, In-situ measurement, Rheology

Deformation-induced crystallographic-preferred orientation of ϵ -FeOOH

*Yu NISHIHARA¹, Yui MORI¹, Wentian WU¹, Noriyoshi TSUJINO²

1. GRC, Ehime Univ., 2. JASRI

Keywords: Crystallographic-preferred orientation, Seismic anisotropy, ϵ -FeOOH

Strength of metastable olivine and the rheology of subducting cold slab at mantle transition zone

*Rikuto HONDA¹, Tomoaki Kubo¹, Noriyoshi Tsujino², Yuji Higo², Sho Kakizawa², Yuki Shibazaki³, Yu Nishihara⁴

1. Kyushu University, 2. JASRI, 3. KEK, 4. Ehime Univ. GRC

Keywords: metastable olivine, Peierls mechanism, deep slab, high-pressure deformation experiments, deep-focus earthquake

High-resolution measurement of ice anelasticity over a broad frequency range with a new cryogenic forced oscillation apparatus

*Hatsuki Yamauchi¹, Christine McCarthy¹, Benjamin Holtzman¹, David Goldsby², Travis Hager²

1. Columbia Univ. LDEO, 2. U. Penn

Keywords: ice, anelasticity, attenuation

Relationship between creep and grain growth in rock with bicontinuous structure

*Shenghao Jiang¹, Takehiko Hiraga¹

1. The University of Tokyo

Keywords: Forsterite, Diopside

Microstructures of plagioclase in gabbroic ultramylonites and quadruple point analysis

*Itsuki Natsume¹, Katsuyoshi Michibayashi^{2,3}, Yohei Igami⁴

1. Kanagawa Prefectural Museum of Natural History, 2. Nagoya Univ., 3. JAMSTEC, 4. Kyoto Univ.

Keywords: Quadruple points, Microstructure, Plagioclase, Ultramylonite

Reconstruction of the uppermost mantle continuous structure of the Oman ophiolite

*Takeo Okuwaki¹, Natsume Itsuki², Katsuyoshi Michibayashi¹

1. Nagoya Univ. Env, 2. Kanagawa Pref. Mus. Nat. Hist.

Keywords: mantle flow, Oman ophiolite, peridotite, foliation, Crystallographic Preferred Orientation (CPO)

Deformation and melt-rock interaction in the Horoman peridotite: Petrological and structural study of the MHL suite and BDH suite rocks

*Aya Hihara¹, Miki Tasaka¹, Keisuke Kurihara¹, Hajime Taniuchi², Tostuhiko Kawamoto¹

1. Shizuoka Univ., 2. AIST

Keywords: mantle, peridotite, deformation, crystallographic preferred orientation, EBSD

Multicomponent measurements of seismic velocity and electrical resistivity using foliated serpentinite and peridotite

*Tomohiro Ito¹, Ikuo Katayama¹, Katsuyoshi Michibayashi², Kazuki Matsuyama²

1. Hiroshima University, 2. Nagoya University

Water is involved in various phenomena at subduction zones. When a subducting plate releases water into the mantle wedge, the water reacts with the mantle to form serpentinite. Seismic low velocity and high electrical resistivity in the mantle have been interpreted as mantle hydration; however, these geophysical data can be highly anisotropic. In this study, we performed multicomponent measurements of seismic velocity and electrical resistivity of the foliated serpentinite and peridotite to discuss the fluid movement and mantle hydration at subduction zones. The rocks used in this study are serpentinite and peridotite with the x-axis is parallel to the lineation and the z-axis is normal to the foliation. An intravessel deformation and fluid flow apparatus was used to measure seismic velocity and electrical resistivity at the confining pressures ranging from 5 MPa to 200 MPa. The fluid used was a 0.5 mol/L NaCl solution, and the fluid pressure was controlled at 1 MPa. These measurements represent the physical properties of the rocks, including microcracks. We also measured the crystal orientation using EBSD to evaluate the anisotropy of the minerals. Serpentinite and peridotite show higher P-wave velocity in the x- and y-axes than in the z-axis, although the P-wave velocity of serpentinite is nearly the same in the x- and y-axes. These results are most likely due to crack alignments subparallel to the foliation and crystal preferred orientation. The electrical resistivity of serpentinite tends to be about an order of magnitude lower in the x-axis than in the z-axis, while the electrical resistivity of peridotite is almost the same in the z- and x-axes. From the multicomponent seismic velocities, we calculated the elastic constant tensor and produced pole figures of seismic velocity (Mainprice, 2014). We will discuss these data in more detail with application to seismic tomography and electrical resistivity data at subduction zones.

Keywords: Ultramafic rocks, Seismic velocity, Electrical resistivity, Anisotropy

Poster presentation | S1: Dynamics of igneous processes (Special Session)

📅 Thu. Sep 12, 2024 12:30 PM - 2:00 PM JST | Thu. Sep 12, 2024 3:30 AM - 5:00 AM UTC | 🏢 Entrance Hall Higashiyama Campus

S1: Dynamics of igneous processes (Special Session)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S1-P-01] Oxidation states of HIMU-type ocean island basalts: Insights from μ -XANES analysis of quenched glasses and melt inclusions

*Yuuki HAGIWARA¹, Hidemi Ishibashi², Takeshi Hanyu¹ (1. Japan Agency for Marine-Earth Science and Technology, 2. Shizuoka Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S1-P-02] The composition and pressure of the fluid in crystal mush estimated from cordierite in tonalitic polycrystalline volcanic ejecta

*Shumpei YOSHIMURA¹ (1. Hokkaido University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S1-P-03] Geochemical evolution of Izu-Oshima volcano: Constraints from analysis of melt inclusions in a submarine core

*Morihsa HAMADA¹, Erika TANAKA², Takeshi HANYU¹, Kenji SHIMIZU³, Takayuki USHIKUBO², Qing CHANG¹, Yoshihiko TAMURA¹ (1. IMG, JAMSTEC, 2. Marine Core Research Institute, Kochi Univ., 3. Kochi Institute for Core Sample Research, JAMSTEC)

Oxidation states of HIMU-type ocean island basalts: Insights from μ -XANES analysis of quenched glasses and melt inclusions

*Yuuki HAGIWARA¹, Hidemi Ishibashi², Takeshi Hanyu¹

1. Japan Agency for Marine-Earth Science and Technology, 2. Shizuoka Univ.

Keywords: Ocean island basalt, Oxidation state, Melt inclusion, μ -XENES

The composition and pressure of the fluid in crystal mush estimated from cordierite in tonalitic polycrystalline volcanic ejecta

*Shumpei YOSHIMURA¹

1. Hokkaido University

Keywords: Cordierite, Crystal mush

Geochemical evolution of Izu-Oshima volcano: Constraints from analysis of melt inclusions in a submarine core

*Morihsa HAMADA¹, Erika TANAKA², Takeshi HANYU¹, Kenji SHIMIZU³, Takayuki USHIKUBO², Qing CHANG¹, Yoshihiko TAMURA¹

1. IMG, JAMSTEC, 2. Marine Core Research Institute, Kochi Univ., 3. Kochi Institute for Core Sample Research, JAMSTEC

Background

Magmas erupting from Izu-Oshima volcano have been interacted with those from Izu-Tobu volcano (Ishizuka et al., 2015, EPSL). Ishizuka et al. (2015) clarified geochemical evolution of Izu-Oshima volcano based on whole-rock geochemical analysis of Izu-Oshima onland samples. In order to extend their previous study, we analyzed melt inclusions in a submarine core.

Samples and analytical methods


We recovered a 25-cm long submarine push core from the seafloor located as far as about 10 km east of Izu-Oshima volcano during the research cruise KR21-16. This core comprises of tephra layers deposited from 4,500 y.B.P. to 3,200 y.B.P., where ages were determined by ¹⁴C dating of foraminifera. The push core samples were divided into ten parts, each of which is 2.5 cm long. We then collected minerals (olivine, plagioclase and orthopyroxene) from each part and polished them until the surface of the melt inclusions were exposed. Volatile elements (H₂O, CO₂, S, F and Cl) and P₂O₅ were analyzed by SIMS, and major elements were analyzed by EPMA. For larger melt inclusions, we also analyzed trace elements by LA-ICP-MS.

Results and discussion


We finally analyzed 100 melt inclusions (51-55 wt% SiO₂) in total. Geochemical data of melt inclusions can be separated into 3 groups. Group A melt inclusions are dominant at upper levels of the push core, which corresponds to Izu-Oshima melts (Ishizuka et al., 2015). Systematics of H₂O and CO₂ concentrations of Group A melt inclusions demonstrate that crystallization differentiation took place at P<100 MPa. Group C melt inclusions are dominant at lower levels, which is hybrid magma of Izu-Oshima and Izu-Tobu magmas. Systematics of H₂O and CO₂ concentrations of Group C melt inclusions demonstrate that crystallization differentiation took place at P>100 MPa. Group B melt inclusions are dominant at intermediate levels and show intermediate characteristics between Group A and Group C melt inclusions, suggesting that Group A and Group C mixed. We argue that melt composition of Izu-Oshima volcano evolved from Group C, Group B to Group A from 4,500 y.B.P. to 3,200 y.B.P.

Keywords: Izu-Oshima volcano, melt inclusion



Izu-Oshima volcano (Older Oshima Group; Ishizuka et al., 2015)





Izu-Tobu volcanoes (Ishizuka et al., 2015)

**This study**



Group A (equivalent to Izu-Oshima melts, dominant at younger ages)



Group B (intermediate melts between Group A and Group C)



Group C (closer to Izu-Tobu melts, dominant at older ages)



Poster presentation | S3: Rheology and Material Transfer in Mantle and Crust (Special Session)

📅 Thu. Sep 12, 2024 12:30 PM - 2:00 PM JST | Thu. Sep 12, 2024 3:30 AM - 5:00 AM UTC | 🏢 Entrance Hall Higashiyama Campus

S3: Rheology and Material Transfer in Mantle and Crust (Special Session)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S3-P-01] Preliminary results of deformation experiments on hydrous stishovite using a rotational DAC

*Shintaro AZUMA¹, Keishi Okazaki², Kentaro Uesugi³, Masahiro Yasutake³, Steeve Gréaux⁴, Yoshiyuki Okuda^{1,5}, Bunrin Natsui¹, Eranga Jayawickrama², Kenji Ohta¹ (1. Tokyo Tech., 2. Hiroshima Univ., 3. JASRI, 4. Ehime Univ., 5. University of Hawai'i)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S3-P-02] Water weakening of Mg₂SiO₄ ringwoodite

「発表賞エントリー」

*Yuta Goto¹, Tomoaki Kubo¹, Rikuto Honda¹, Yuki Shibazaki² (1. Kyushu Univ., 2. KEK-PF)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S3-P-03] Toward an understanding of dehydration process of partially serpentinized slab peridotite under conditions where deep earthquakes occur

*Tomoaki KUBO¹, Musashi Ezaki¹, Nobumasa Fujiwara¹, Rikuto Honda¹, Goto Yuta¹, Noriyoshi Tsujino² (1. Kyushu University, 2. JASRI)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S3-P-04] Viscous anisotropy of olivine aggregates using micro Vickers indentation tests

「発表賞エントリー」

*Namu Fujii¹, Miki Tasaka¹ (1. Shizuoka University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S3-P-05] Crystal-fabric analysis using principal component analysis method for the Horoman peridotite

「発表賞エントリー」

*Kazuki Matsuyama¹, Katsuyoshi Michibayashi¹ (1. Nagoya Univ. Env.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S3-P-06] Traces of mantle fluid/melt within olivine phenocrysts from Ohima-Oshima picritic basalts

*Ryo Tsukawaki¹, Terumi Ejima², Atusi Ninomiya³, Shoji Arai⁴ (1. Shinshu Univ. Sci., 2. Shinshu Univ. Sci., 3. Sumiko Res. Exp. & Dev. Co., 4. Kanazawa Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S3-P-07] Microstructural characteristics of ultramafic rocks in the Tosa Megamullion, the Shikoku Basin.

「発表賞エントリー」

*So Inoue¹, Katsuyoshi Michibayashi^{1,2}, Yumiko Harigane³, Yasuhiko Ohara^{1,2,4} (1. GSES, Nagoya Univ., 2. JAMSTEC, 3. GSJ/AIST, 4. JCG)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S3-P-08] Deformation microstructures of granitic mylonite in Hida Metamorphic Belt

「発表賞エントリー」

*Masaaki Horie¹, Katsuyoshi Michibayashi¹ (1. GSES, Nagoya Univ.)

Preliminary results of deformation experiments on hydrous stishovite using a rotational DAC

*Shintaro AZUMA¹, Keishi Okazaki², Kentaro Uesugi³, Masahiro Yasutake³, Steeve Gréaux⁴, Yoshiyuki Okuda^{1,5}, Bunrin Natsui¹, Eranga Jayawickrama², Kenji Ohta¹

1. Tokyo Tech., 2. Hiroshima Univ., 3. JASRI, 4. Ehime Univ., 5. University of Hawai'i

Keywords: Rotational diamond anvil cell, High temperature and pressure, Hydrous stishovite, Rheology, Deformation experiment

Water weakening of Mg_2SiO_4 ringwoodite

*Yuta Goto¹, Tomoaki Kubo¹, Rikuto Honda¹, Yuki Shibasaki²

1. Kyushu Univ., 2. KEK-PF

Keywords: in-situ X-ray observation, high-pressure deformation experiment, water content, water weakening, mantle transition zone

Toward an understanding of dehydration process of partially serpentized slab peridotite under conditions where deep earthquakes occur

*Tomoaki KUBO¹, Musashi Ezaki¹, Nobumasa Fujiwara¹, Rikuto Honda¹, Goto Yuta¹, Noriyoshi Tsujino²

1. Kyushu University, 2. JASRI

Keywords: deep slab, dehydration process, dehydration embrittlement, water weakening

Viscous anisotropy of olivine aggregates using micro Vickers indentation tests

*Namu Fujii¹, Miki Tasaka¹

1. Shizuoka University

Keywords: olivine, low temperature plasticity, Vickers indentation tests, anisotropy, viscosity

Crystal-fabric analysis using principal component analysis method for the Horoman peridotite

*Kazuki Matsuyama¹, Katsuyoshi Michibayashi¹

1. Nagoya Univ. Env.

Keywords: Olivine, Crystal-fabric, Horoman peridotite complex, Principal component analysis

Traces of mantle fluid/melt within olivine phenocrysts from Ohima-Ōshima picritic basalts

*Ryo Tsukawaki¹, Terumi Ejima², Atusi Ninomiya³, Shoji Arai⁴

1. Shinshu Univ. Sci., 2. Shinshu Univ. Sci. , 3. Sumiko Res. Exp. & Dev. Co., 4. Kanazawa Univ.

Keywords: Mantle, Picritic basalt, Olivine, Fluid/melt, Oshima-Ōshima

Microstructural characteristics of ultramafic rocks in the Tosa Megamullion, the Shikoku Basin.

*So Inoue¹, Katsuyoshi Michibayashi^{1,2}, Yumiko Harigane³, Yasuhiko Ohara^{1,2,4}

1. GSES, Nagoya Univ. , 2. JAMSTEC, 3. GSJ/AIST, 4. JCG

Keywords: Megamullion, Olivine, Back-arc basin, ductile shear deformation

Deformation microstructures of granitic mylonite in Hida Metamorphic Belt

*Masaaki Horie¹, Katsuyoshi Michibayashi¹

1. GSES. Nagoya Univ.

Keywords: Granitic mylonite, Deformation microstructure, Crystallographic preferred orientation, Hida Metamorphic Belt

Poster presentation | R1: Characterization and description of minerals (Joint Session with The Gemmological Society of Japan)

📅 Thu. Sep 12, 2024 12:30 PM - 2:00 PM JST | Thu. Sep 12, 2024 3:30 AM - 5:00 AM UTC | 🏢 Entrance Hall Higashiyama Campus

R1: Characterization and description of minerals (Joint Session with The Gemmological Society of Japan)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-01] Chemical quantitative analysis of heulandite using SEM-EDS - How can we accurately estimate the chemical composition of zeolite?

「発表賞エントリー」

*Atsushi ISHIHARA¹, Hiroaki Ohfuji¹ (1. Tohoku university)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-02] Quantitative electron microprobe analysis of xenotime

*Yasuyuki BANNO¹ (1. AIST)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-03] Deformation and compositional changes of plagioclase: A combined electron backscattered diffraction and energy dispersive X-ray spectroscopy approach

「発表賞エントリー」

*Kohei Nimura¹, Katsuyoshi Michibayashi^{1,2} (1. Nagoya University, 2. JAMSTEC)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-04] Mineralogical features of smelting slags from the Early Iron Age Yashin Tepe site, northeastern Iraq

*Masanori KUROSAWA¹, Shin'ichi Nishiyama² (1. Univ. Tsukuba, 2. Chubu Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-05] The origin of abundant graphite in quartz veins in Ishidera area, Wazuka Town, Kyoto Prefecture, Japan

*Masaki Nishio¹, Itaru Mitsukawa¹, Yohei Igami¹, Akira Miyake¹, Norimasa Shimobayashi¹ (1. Kyoto Univ. Sci.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-06] Michitoshiite-(Cu), a new Ge-containing platinum group mineral from Haraigawa, Misato machi, Kumamoto Prefecture, Japan

*Takahiro TANAKA¹, Daisuke Nishio Hamane², Tadashi Shinmachi (1. Nittetsu Mining Co., Ltd., 2. ISSP, Univ. of Tokyo)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-07] Fibrous inclusions in rose quartz

*Yohei SHIROSE¹, Hayato Fudamoto¹, Sayako Inoue² (1. Ehime Univ. Sci., 2. Ehime Univ. GRC)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-08] Rose quartz in gneisses from Uoshima Island, Ehime Prefecture

*Yohei SHIROSE¹, Shoma Sakai¹ (1. Ehime Univ. Sci.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-09] Secondary arsenate minerals from the Takumi Mine, Hyogo Prefecture, Japan

*Yohei SHIROSE¹, Riakako Kamise¹, Katsuichi Nishida, Yoshiteru Fujiwara (1. Ehime Univ. Sci.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-10] Mineralogical properties of lautenthalite and wroewolfeite from the Nii mine, Hyogo Prefecture, Japan

*Masayuki Ohnishi, Norimasa Shimobayashi¹, Daisuke Nishio-Hamane², Keiji Shinoda³, Takeshi Hisano (1. Sci., Kyoto Univ., 2. ISSP, Univ. of Tokyo, 3. Sci., Osaka Metro. Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-11] A re-examination of Sr-rich apatite from Itoigawa, Niigata Prefecture, Japan

*Seiichiro UEHARA¹, Koichi MONMA², Masayuki OHNISHI, Shunsuke OHSUMI, Yoshiya OHKI, Hiroki OKA³ (1. Kyushu Univ. Museum, 2. Nat'l. Mus. Nat. Sci., 3. OYO Corp.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-12] Hydroxylchondrodite from Ogouchi, Hinokage, Nisiusuki, Miyazaki Prefecture, Japan

*Toshiro Okada¹, Seiichiro Uehara², Isao Yukinori³, Yohei Shirose⁴ (1. Kashii 2 JHS, 2. Kyushu Univ, 3. Fukuoka Stc, 4. Ehime Univ)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-13] Arsenmedaite from the Yamato mine, Kagoshima Prefecture, SW Japan

*Shunsuke Ohsumi, Daisuke Nishio-Hamane¹, Hiroki Oka², Masashi Tamura³, Kosuke Takagi⁴ (1. ISSP, Univ. of Tokyo, 2. OYO Corp., 3. Fac. Eng. Tech. Div., Mie Univ., 4. Grad. Sch. of Eng., Mie Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-14] "Common Hornblende" from Mt. Tawarayama (Goou-toge), the outer-rim of Mt. Aso, Kumamoto Prefecture

*Haruki Inoue¹, Seiichiro Uehara² (1. Enecom Co., Ltd., 2. Kyushu Univ. Museum)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-15] Chemical composition of tourmaline and amphibole associated with gabbro from Kajishima, Ehime Prefecture, Japan

「発表賞エントリー」

*Itsuki Ota¹, Kazuya Shimooka², satoshi saitou¹, youhei shirose¹ (1. Ehime Univ. Sci and Eng, 2. Kwansei Gakuin Univ. Sci)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-16] Constituent minerals of clay associated with the pegmatite dike in Nagatare, Fukuoka Prefecture, Japan

*Yuya TAKEDA¹, Seiichiro Uehara², Yoshihiro Kuwahara³ (1. Kyushu Univ. ISGS, 2. Kyushu Univ. Museum, 3. Kyushu Univ. SCS)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R1-P-17] Microstructure of serpentine veins in peridotite in Ooshika Village, Nagano Prefecture, Japan

*Yuya TAKEDA¹, Yoshihiro Kuwahara³, Seiichiro Uehara² (1. Kyushu Univ. ISGS, 2. Kyushu Univ. Museum, 3. Kyushu Univ. SCS)

Chemical quantitative analysis of heulandite using SEM-EDS - How can we accurately estimate the chemical composition of zeolite?

*Atsushi ISHIHARA¹, Hiroaki Ohfuji¹

1. Tohoku university

Keywords: Zeolite, SEM-EDS, heulandite

$$*\text{balance error} : E = \frac{[\text{Al}] - ([\text{Na}] + [\text{K}] + [\text{Ca}] \times 2)}{([\text{Na}] + [\text{K}] + [\text{Ca}] \times 2)} \times 100 \text{ (Passaglia, 1970)}$$

Quantitative electron microprobe analysis of xenotime

*Yasuyuki BANNO¹

1. AIST

The conditions suitable for the quantitative analysis of xenotime from the Takehara mine in Mie Prefecture, Japan, using EPMA were investigated. First, a qualitative analysis was performed to identify the major elements and determine the characteristic X-ray positions and the appropriate background measurement positions. In cases where the overlap of peaks was significant, a pulse height analyzer was used when higher-order lines overlapped. An interference correction factor was determined to correct the measured intensity when first-order lines interfered.

Keywords: EPMA, Xenotime, Chemical composition, Takehara mine

Deformation and compositional changes of plagioclase: A combined electron backscattered diffraction and energy dispersive X-ray spectroscopy approach

*Kohei Nimura¹, Katsuyoshi Michibayashi^{1,2}

1. Nagoya University, 2. JAMSTEC

Keywords: plagioclase, recrystallization, electron backscattered diffraction, energy dispersive X-ray spectroscopy

Mineralogical features of smelting slags from the Early Iron Age Yashin Tepe site, northeastern Iraq

*Masanori KUROSAWA¹, Shin'ichi Nishiyama²

1. Univ. Tsukuba, 2. Chubu Univ.

Microstructures, materials, and chemical compositions of four metal-smelting slags and two specimens of corroded ironware from the Early Iron Age Yashin Tepe site in northeastern Iraq were analyzed with a scanning electron microscope equipped with an energy-dispersive X-ray spectrometer (SEM-EDS) to examine the technical level of the Iron Age iron-smelting in the frontier area of West Asia. Two types of metal-smelting slag were identified in the slag samples: copper smelting slag and iron smelting slag. The surfaces of both types were covered with a thin altered film. The copper smelting slag was vitreous with a light greenish interior and consisted mainly of Ca-Al silicate glass and precipitated crystals of augite. The slag also contained tiny fragments of limestone in the process of melting reaction. The ironmaking slag was black vitreous inside and composed of dendritic crystals of wustite, Ca-Al silicate glass, and Ca-Al silicate mineral precipitates. The Ca-Al silicates were found to have precipitated at 1200-1300 degrees. Small pieces of limestone in the middle of the melting reaction were also observed. Since limestone is abundant around the site, it is thought that limestone was used as a slag-forming agent in metal smelting. The CaO-rich slag produced by limestone incorporation is favorable for the production of high-purity iron, suggesting that high-quality iron was produced in the early Iron Age. The ironware samples were almost completely corroded and consisted mainly of iron hydroxide and small amounts of iron oxides. However, some traces of metallic iron were also observed, and the metallic iron was of high quality with very few impurities. These results indicate that iron suitable for ironware making with high purity was produced in the frontier area of West Asia from around 800 BCE using advanced smelting technology that utilizes high temperatures of over 1200 degrees and limestone as a slag-forming agent.

Keywords: slag, iron smelting, ironware, SEM-EDS, Yashin-Tepe site

The origin of abundant graphite in quartz veins in Ishidera area, Wazuka Town, Kyoto Prefecture, Japan

*Masaki Nishio¹, Itaru Mitsukawa¹, Yohei Igami¹, Akira Miyake¹, Norimasa Shimobayashi¹

1. Kyoto Univ. Sci.

Keywords: Graphite, Fluid inclusion, C-O-H fluid, Quartz vein




図 (a) 光学顕微鏡下で球状の外形をなす、石英粒子内の石墨 (※赤線はFIBで切り出した場所を示す)
(b) (a)から切り出した試料のTEMでのHAADF-STEM像

Michitoshiite-(Cu), a new Ge-containing platinum group mineral from Haraigawa, Misato machi, Kumamoto Prefecture, Japan

*Takahiro TANAKA¹, Daisuke Nishio Hamane², Tadashi Shinmachi

1. Nittetsu Mining Co., Ltd., 2. ISSP, Univ. of Tokyo

Michitoshite-(Cu) is a new mineral of Ge-containing platinum group minerals (PGM) discovered from the Haraigawa, Kumamoto Prefecture, Japan. It is named in honor of Michitoshi Miyahisa (1928-1983), a former professor of the Ehime University. The mineral and name have been approved by the IMA Commission on New Minerals, Nomenclature and Classification (IMA2019-029a). The PGM placer deposit where the michitoshiite-(Cu) was found is located in a small stream that crossing a clinopyroxenite mass (Nishio-Hamane et al., 2019). Michitoshite-(Cu) is opaque, has a metallic luster, and is silver-gray in color. It is light gray with brownish tints and shows no pleochroism and anisotropy in reflected light. The Mohs hardness is 5, and the density calculated from the empirical formula and powder X-ray diffraction (XRD) data is 10.78 g/cm³. The empirical formula is calculated on the basis of 2 apfu is $(\text{Rh}_{0.95}\text{Pt}_{0.03}\text{Ir}_{0.01})_{\Sigma 0.99}[(\text{Cu}_{0.36}\text{Fe}_{0.24})_{\Sigma 0.60}\text{Ge}_{0.41}]_{\Sigma 1.01}$. The powder XRD pattern shows five strong peaks [d in Å (I/10) hkl], 2.103 (100) 110, 1.717 (3) 111, 1.487 (15) 200, 1.332 (6) 210, 1.216 (70) 211, and can be indexed as the cubic *Pm-3m* (#221) with the lattice parameters $a = 2.9771(11)$ Å and $V = 26.39(3)$ Å³ ($Z = 1$). From previous studies, the RhCu-RhFe-RhGe system has a *face-centered cubic (fcc)* structure for RhCu, a CsCl-type structure for RhFe, and a MnP-type structure for RhGe, with the end-member of each compound having a different structure, but the solid solution structure has not been investigated. Since $\text{Rh}[(\text{Cu}_{0.35}\text{Fe}_{0.25})_{\Sigma 0.60}\text{Ge}_{0.40}]$, which was synthesized to mimic the empirical formula of the natural sample, had a CsCl-type structure, the crystal structure of michitoshite-(Cu) was considered to be CsCl-type. Considering that the synthetic materials of Rh(Cu,Fe) composition synthetics were not *bcc-based* crystal structures but rather *fcc* structures in the synthesis experiments, the presence of Ge is considered essential for the formation of the *bcc-based* crystal structure. Therefore, based on the relationship between composition and crystal structure obtained from the behavior of other synthetics, the ideal formula for michitoshite-(Cu) is $\text{Rh}(\text{Cu}_{1-x}\text{Ge}_x)$ $0 < x \leq 0.5$.

Keywords: Michitoshiite-(Cu), Germanium, Platinum group mineral, Synthetic experiment, Kumamoto




図. 三千年鉱の産状 (a: 写真, b: SEM像, c: BSE像, d: BSE像), 三千年鉱の反射顕微鏡写真 (e: オープンニコル, f: クロスニコル, g: オープンニコル, h: クロスニコル), 三千年鉱の取りうる結晶構造 (i), 合成実験における三千年鉱の組成範囲及び固溶体の結晶構造 (j).

Fibrous inclusions in rose quartz

*Yohei SHIROSE¹, Hayato Fudamoto¹, Sayako Inoue²

1. Ehime Univ. Sci., 2. Ehime Univ. GRC

Keywords: rose quartz, dumortierite, dumortierite-like mineral, pegmatite, Fukuyoshi




Fig. SEM-BEI and TEM-BFI of dumortierite-like fibrous inclusion (Dm) in rose quartz (Qz)

Rose quartz in gneisses from Uoshima Island, Ehime Prefecture

*Yohei SHIROSE¹, Shoma Sakai¹

1. Ehime Univ. Sci.

Keywords: rose quartz, gneiss, dumortierite-like mineral, Uoshima Island, boron




Fig. (a) Photograph and (b) crossed polarized photomicrograph of rose quartz in a gneiss from Uoshima Island.

Secondary arsenate minerals from the Takumi Mine, Hyogo Prefecture, Japan

*Yohei SHIROSE¹, Riakako Kamise¹, Katsuichi Nishida, Yoshiteru Fujiwara

1. Ehime Univ. Sci.

Keywords: rooseveltite, parasymphesite, scorodite, bismuthinite, Takumi Mine




Fig. BEI of rooseveltite with bismuthinite (Bin) and arsenopyrite (Apy).

Mineralogical properties of lautenthalite and wroewolfeite from the Nii mine, Hyogo Prefecture, Japan

*Masayuki Ohnishi, Norimasa Shimobayashi¹, Daisuke Nishio-Hamane², Keiji Shinoda³, Takeshi Hisano

1. Sci., Kyoto Univ., 2. ISSP, Univ. of Tokyo, 3. Sci., Osaka Metro. Univ.

Lautenthalite and wroewolfeite have been found in the dump of the Nii mine, Hyogo Prefecture, Japan. The minerals occur in cracks of quartz containing ore minerals such as chalcopyrite, galena and sphalerite.

Lautenthalite is aggregates of platy crystals up to 0.5 mm in length and 0.1 mm in thick. Lautenthalite is always in overgrowth on wroewolfeite. The mineral is blue-green in color with a vitreous luster. It is monoclinic with unit cell parameters of $a = 21.557(6)$, $b = 6.019(1)$, $c = 22.467(5)$ Å, and $\beta = 108.06(3)^\circ$. A WDS analysis yields the empirical formula (based on total cations = 7; water by stoichiometry), $(\text{Pb}_{0.97}\text{Ca}_{0.06})_{\Sigma 1.03}(\text{Cu}_{3.97}\text{Zn}_{0.11}\text{Mn}_{0.02})_{\Sigma 4.10}(\text{SO}_4)_{1.87}(\text{OH})_{6.52} \cdot 3\text{H}_2\text{O}$.

Wroewolfeite is aggregates of platy crystals up to 1 mm in length and 0.1 mm in thick in association with langite, brochantite, linarite and cerussite. The mineral is blue in color with a vitreous luster. It is monoclinic with unit cell parameters of $a = 6.042(8)$, $b = 5.637(6)$, $c = 14.45(2)$ Å, and $\beta = 93.5(1)^\circ$. A WDS analysis yields the empirical formula (based on total cations = 5; water by stoichiometry), $(\text{Cu}_{3.94}\text{Zn}_{0.04}\text{Ca}_{0.02}\text{Fe}_{0.01}\text{Pb}_{0.01}\text{Mn}_{0.01})_{\Sigma 4.04}(\text{SO}_4)_{0.96}(\text{OH})_{6.14} \cdot 2\text{H}_2\text{O}$. Optically, the mineral is biaxial (-) with refractive indices $\alpha = 1.641(2)$, $\beta = 1.691(2)$ and $\gamma = 1.701(2)$, and $2V(\text{calc.}) = 47.0^\circ$.

Keywords: Lautenthalite, Wroewolfeite, Sulfate, Nii mine




Figure 1. (a) and (b) Photomicrographs of aggregates of lautenthalite and wroewolfeite crystals. (c) A photomicrograph of aggregates of wroewolfeite crystals. (d) A back-scattered electron image of a polished section of lautenthalite and wroewolfeite. Abbreviations: Lth, lautenthalite; Wwf, wroewolfeite; Qz, quartz; Cer, cerussite.

A re-examination of Sr-rich apatite from Itoigawa, Niigata Prefecture, Japan

*Seiichiro UEHARA¹, Koichi MONMA², Masayuki OHNISHI, Shunsuke OHSUMI, Yoshiya OHKI, Hiroki OKA³

1. Kyushu Univ. Museum, 2. Nat' l. Mus. Nat. Sci., 3. OYO Corp.

Keywords: Sr-apatite like mineral, OH-analog of stronadelphite, New Mineral, Omi-gawa River, Itoigawa, Niigata Prefecture, Albitite




Fig. 1 SEM images of Sr-apatite ('hydroxylstronadelphite') in albitite from the Omi-gawa River, Itoigawa, Niigata Prefecture.
(a) Large area RGB X-ray mapping image (R:CaK α , G:SrL α , B:PK α). Red grains are pectolite, blue grains are Sr-apatite.
(b) BSE image of Sr-apatite.

Hydroxylchondrodite from Ogouchi ,Hinokage, Nisiusuki,Miyazaki Prefecture,Japan

*Toshiro Okada¹, Seiichiro Uehara², Isao Yukinori³, Yohei Shirose⁴

1. Kashii 2 JHS, 2. Kyushu Univ, 3. Fukuoka Stc, 4. Ehime Univ

Keywords: Hydroxylchondrodite, Humite group, Phlogopite, Skarn, Ogouchi Hinokage Nisiusuki Miyazaki Prefecture




Fig. 1. Hydroxylchondrodite and associated minerals from Ogouchi, Hinokage, Nisiusuki, Miyazaki Prefecture, Japan.

(a) Sample shows occurrence of hydroxylchondrodite of sample No. SD11. Sample width is 6.2 cm.

(b) Hydroxylchondrodite crystal with phlogopite of sample No. SD11.

(c) Thin section image under plane polarized light of sample No. SD01, which contains hydroxylchondrodite, uvite, phlogopite, diopside, clinocllore and titanite. Width is 2.0 cm.

(d) Back scattered electron image of hydroxylchondrodite and associate minerals. The red frame in figure (c) corresponds to this BSE image. Darkest grains are hydroxylchondrodite and clinocllore. Small bright grains are mainly titanite some are zirconolite and apatite. Arrow indicates zirconolite.

Hchn: hydroxylchondrodite. Phl: phlogopite. Dp: diopside. Uvt: uvite. Clc:clinocllore

Arsenmedaite from the Yamato mine, Kagoshima Prefecture, SW Japan

*Shunsuke Ohsumi, Daisuke Nishio-Hamane¹, Hiroki Oka², Masashi Tamura³, Kosuke Takagi⁴

1. ISSP, Univ. of Tokyo, 2. OYO Corp., 3. Fac. Eng. Tech. Div., Mie Univ., 4. Grad. Sch. of Eng., Mie Univ.

The Yamato mine in the Amami Oshima Island, Kagoshima Prefecture, southwestern Japan, is a well-known manganese mine as the type locality of haradaite (Watanabe et al., 1982) and occurrences of V and/or As-rich minerals such as roscoelite (Yoshimura & Momoi, 1964), goldmanite (Momoi, 1964), tiragalloite (Nakao et al., 2005), palenzonaite, nabiasite, tokyoite, vuorelainenite (Yamada et al., 2008), medaite, saneroite (Matsubara et al., 2013), poppiite (Niwa & Ishibashi, 2015), and reppiaite (Yamada et al., 2018). In this study, we report the first occurrence of arsenmedaite in Japan. Arsenmedaite, As⁵⁺-analogue of medaite, was reported in quartz veinlets cutting braunite ores from the Molinello mine, which is hosted within metacherts of the Northern Apennines. It is associated with calcite, As-rich medaite, rhodochrosite, talc, and ganophyllite. In the Yamato mine, arsenmedaite was found as from yellow to orange-reddish granular crystals, up to 1 mm in diameter, with a vitreous lustre in siliceous manganese ores, which consist mainly of massive hausmannite, rhodochrosite, and coarse-grained rhodonite, with absence of medaite. Tiny crystals of rhodonite are ubiquitously included in arsenmedaite. The occurrence of arsenmedaite in the Yamato mine differs from that in the type locality. The representative chemical composition by SEM-EDS is: SiO₂ 30.12, CaO 0.99, V₂O₅ 3.25, MnO 48.03, As₂O₅ 13.26, H₂O_{calc.} 1.01, total 96.65 wt%, and they gave the empirical formula as (Mn²⁺_{6.04}Ca_{0.16})_{Σ6.20}[(As_{1.03}V_{0.32})_{1.35}Si_{4.47}]_{Σ5.82}O₁₈(OH), assuming total cations = 18 and Mn = Mn²⁺. The empirical formula is slightly richer in As and V and less in Si than the ideal formula of arsenmedaite, Mn²⁺₆As⁵⁺Si₅O₁₈(OH). Therefore, it is inferred that As and V partly substitute the Si-tetrahedral framework in arsenmedaite. Raman spectroscopy detected weak peaks at 340–360 and 994 cm⁻¹, moderate peaks at 644 and 658 cm⁻¹, and strong peaks at 859, 879, and 894 cm⁻¹. These Raman peaks are comparable with those of arsenmedaite (Biagioni et al., 2019) and medaite (Lafuente et al., 2015). Although a precise crystallographic analysis has not been completed yet due to its rarity and rhodonite inclusions, the occurrence of arsenmedaite possibly contributes to V/As-mineralization in the Yamato mine.

Keywords: arsenmedaite, Yamato mine

“Common Hornblende” from Mt. Tawarayama (Goou-toge), the outer-rim of Mt. Aso, Kumamoto Prefecture

*Haruki Inoue¹, Seiichiro Uehara²

1. Enecom Co., Ltd., 2. Kyushu Univ. Museum

Keywords: Mt. Tawarayama, Goou-toge, magnesio-hastingsite, common hornblende




Fig.1. Polarizing microscope images and scanning electron microscope images of “Common Hornblende” from Mt. Tawarayama (Goou-toge), the outer-rim of Mt. Aso, Kumamoto Prefecture, Japan.

(a) Optical microscope image of amphibole phenocryst under plane polarized light. The cross section is parallel to the c -axis. (b) EDS X-ray maps of amphibole phenocryst. This image has RGB colors overlaid. The red frame indicates the area in Fig. 1a. (c) Optical microscope image of opacitization of hornblende under plane polarized light. The blue frame indicates the area in Fig. 1d. (d) BSE image of opacitization of “common hornblende”. There are microscopic minerals that cannot be observed with a polarizing microscope.

Chemical composition of tourmaline and amphibole associated with gabbro from Kajishima, Ehime Prefecture, Japan

*Itsuki Ota¹, Kazuya Shimooka², satoshi saitou¹, youhei shirose¹

1. Ehime Univ. Sci and Eng, 2. Kwansei Gakuin Univ. Sci

Keywords: kajishima, gabbro, pegmatite, tourmaline, amphibole




Fig. 1 角閃石の化学組成変化 (分類はHawthorne et al., 2012; Leake, 1997を参考)

Constituent minerals of clay associated with the pegmatite dike in Nagatare, Fukuoka Prefecture, Japan

*Yuya TAKEDA¹, Seiichiro Uehara², Yoshihiro Kuwahara³

1. Kyushu Univ. ISGS, 2. Kyushu Univ. Museum, 3. Kyushu Univ. SCS

Keywords: kaoline, halloysite, nagatare




Fig. 1. Photographs of clays in Nagatare. (a) Bright field TEM image of halloysite. (b) Bright field TEM image of FIB thin section. (c) Bright field TEM image of kaolinite in the red box area of (b). (d) Enlarged TEM image of kaolinite.

Microstructure of serpentine veins in peridotite in Ooshika Village, Nagano Prefecture, Japan

*Yuya TAKEDA¹, Yoshihiro Kuwahara³, Seiichiro Uehara²

1. Kyushu Univ. ISGS, 2. Kyushu Univ. Museum, 3. Kyushu Univ. SCS

Keywords: chrysotile, serpentine vein, iron




Fig. 1. Photographs of serpentine vein I and II. (a) Back scattered electron image by SEM of serpentine vein I and II. (b) Bright field TEM image of chrysotile and brucite in serpentine vein I. (c) Bright field TEM images of chrysotile and SAED pattern of chrysotile in serpentine vein II. (d) Bright field TEM images of chrysotile and brucite and awaruite in serpentine vein I. SAED pattern of awaruite.

Ctl: chrysotile. Brc: brucite. Aw: awaruite. Chr: chromite. Fo: forsterite.

Poster presentation | R3: High-pressure science and deep Earth's material

📅 Thu. Sep 12, 2024 12:30 PM - 2:00 PM JST | Thu. Sep 12, 2024 3:30 AM - 5:00 AM UTC | 🏢 Entrance Hall Higashiyama Campus

R3: High-pressure science and deep Earth's material

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R3-P-01] Influence of co-existing inorganic minerals on chemical reaction of *n*-alkane under high-pressure and high-temperature conditions of subduction zone.

*Ayako SHINOZAKI¹, Kina Takimoto¹, Takaya Nagai¹, Koichi Mimura² (1. Hokkaido University, 2. Nagoya University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R3-P-02] Differential Scanning Calorimetry of Mn₂SiO₄ tephroite

*Yuta Asami¹, Itaru Ohira², Hiroshi Kojitani² (1. Gakushuin Univ. Sci, 2. Gakushuin Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R3-P-03] Ab initio calculation of the polarized IR spectra and hydrogen positions of hydrous Bridgmanite

*Kikuyo Inagaki¹, Jun Tsuchiya¹, Yanyao Zhang³, Jung-Fu Lin², Shun-ichiro Karato⁴, Jennifer Kung⁵, ChingChien Li⁵ (1. GRC Ehime Univ. , 2. Univ. Texas Austin , 3. Stanford Univ., 4. Yale Univ., 5. National Cheng Kung Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R3-P-04] Determination of the stability of silica phases under high pressure by ultra-fast X-ray diffraction measurements

*Ryosuke SINMYO¹, Saori Kawaguchi-Imada², Takayuki Ishii³, Hiroshi Sakuma⁴, Ayase Ogawa¹, Kenta Kobayashi¹, Shuhou Maitani¹ (1. Meiji Univ. Sci. Tech., 2. JASRI, 3. Okayama Univ. IPM, 4. NIMS)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R3-P-05] Crystallographic preferred orientation properties of Ferropicicase polycrystals obtained from large strain deformation experiments under lower mantle pressures

「発表賞エントリー」

*Bunrin Natsui¹, Shintaro Azuma¹, Keishi Okazaki^{2,5}, Kentaro Uesugi³, Masahiro Yasutake³, Saori Kawaguchi³, Ryuichi Nomura⁴, Kenji Ohta¹ (1. Tokyo Tech, 2. Hiroshima Univ., 3. JASRI, 4. Kyoto Univ., 5. JAMSTEC)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R3-P-06] Investigation of hydrogen sealing materials at high temperature and high pressure using neutron imaging

*Sho KAKIZAWA¹, Hiroyuki Kagi², Masahiro Takano², Asami Sano-Furukawa³, Takanori Hattori³, Abe Jun⁴, Kenichi Funakoshi⁴ (1. JASRI, 2. UTokyo Sci., 3. JAEA J-PARC Center, 4. CROSS, Neutron Science and Technology Center)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R3-P-07] Reactions of FeS with hydrogen at high pressure and high temperature revisited

「発表賞エントリー」

*Masahiro Takano¹, Hiroyuki Kagi¹, Yuichiro Mori¹, Katsutoshi Aoki¹, Sho Kakizawa², Noriyoshi Tsujino², Yuji Higo², Asami Sano-Furukawa³ (1. UTokyo, 2. JASRI, 3. J-PARC center, JAEA)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R3-P-08] Extreme pressure generation using toroidal diamond anvil cell

Session

2024 Annual Meeting of Japan Association of Mineralogical Sciences (JAMS)

*Takeshi SAKAI¹, Yuki Nakamoto², Satoru Nakamura¹, Sotaro Iwatsu², Shuto Fukuda², Yuki Kato², Katsuya Shimizu², Hirokazu Kadobayashi³, Saori Kawaguchi-Imada³ (1. GRC, Ehime University, 2. KYOKUGEN, Osaka University, 3. JASRI)

Influence of co-existing inorganic minerals on chemical reaction of n -alkane under high-pressure and high-temperature conditions of subduction zone.

*Ayako SHINOZAKI¹, Kina Takimoto¹, Takaya Nagai¹, Koichi Mimura²

1. Hokkaido University, 2. Nagoya University

Keywords: Hydrocarbons, olivine, Gas specteometry/Mass spectroscopy , Raman, Subduction zone

Differential Scanning Calorimetry of Mn_2SiO_4 tephroite

*Yuta Asami¹, Itaru Ohira², Hiroshi Kojitani²

1. Gakushuin Univ. Sci, 2. Gakushuin Univ.

Keywords: Tephroite, Heat capacity, d-orbital electrons




図 1.オリビン型 M_2SiO_4 (M=Mn、Fe、Co、Ni、Mg) の定圧モル熱容量

Ab initio calculation of the polarized IR spectra and hydrogen positions of hydrous Bridgmanite

*Kikuyo Inagaki¹, Jun Tsuchiya¹, Yanyao Zhang³, Jung-Fu Lin², Shun-ichiro Karato⁴, Jennifer Kung⁵, ChingChien Li⁵

1. GRC Ehime Univ. , 2. Univ. Texas Austin , 3. Stanford Univ., 4. Yale Univ., 5. National Cheng Kung Univ.

Keywords: Bridgmanite, Hydrogen, ab initio calculation

Determination of the stability of silica phases under high pressure by ultra-fast X-ray diffraction measurements

*Ryosuke SINMYO¹, Saori Kawaguchi-Imada², Takayuki Ishii³, Hiroshi Sakuma⁴, Ayase Ogawa¹, Kenta Kobayashi¹, Shuhou Maitani¹

1. Meiji Univ. Sci. Tech., 2. JASRI, 3. Okayama Univ. IPM, 4. NIMS

Keywords: SiO₂, high pressure and high temperature, X-ray diffraction measurement, seifertite

Crystallographic preferred orientation properties of Ferropericlasite polycrystals obtained from large strain deformation experiments under lower mantle pressures

*Bunrin Natsui¹, Shintaro Azuma¹, Keishi Okazaki^{2,5}, Kentaro Uesugi³, Masahiro Yasutake³, Saori Kawaguchi³, Ryuichi Nomura⁴, Kenji Ohta¹

1. Tokyo Tech, 2. Hiroshima Univ., 3. JASRI, 4. Kyoto Univ., 5. JAMSTEC

Keywords: Rheology, Deformation experiment, Lower mantle




図 活性化するすべり面と温度および圧力の関係



Investigation of hydrogen sealing materials at high temperature and high pressure using neutron imaging

*Sho KAKIZAWA¹, Hiroyuki Kagi², Masahiro Takano², Asami Sano-Furukawa³, Takanori Hattori³, Abe Jun⁴, Kenichi Funakoshi⁴

1. JASRI, 2. UTokyo Sci., 3. JAEA J-PARC Center, 4. CROSS, Neutron Science and Technology Center

Keywords: Hydrogen, Neutron Imaging, hydrogen sealing materials




図 1：（左）使用した高圧セルおよび（右）得られた透過像および透過プロフィール

Reactions of FeS with hydrogen at high pressure and high temperature revisited

*Masahiro Takano¹, Hiroyuki Kagi¹, Yuichiro Mori¹, Katsutoshi Aoki¹, Sho Kakizawa², Noriyoshi Tsujino², Yuji Higo², Asami Sano-Furukawa³

1. UTokyo, 2. JASRI, 3. J-PARC center, JAEA

Keywords: neutron diffraction, X-ray diffraction, FeS, hydrogenation

Extreme pressure generation using toroidal diamond anvil cell

*Takeshi SAKAI¹, Yuki Nakamoto², Satoru Nakamura¹, Sotaro Iwatsu², Shuto Fukuda², Yuki Kato², Katsuya Shimizu², Hirokazu Kadobayashi³, Saori Kawaguchi-Imada³

1. GRC, Ehime University, 2. KYOKUGEN, Osaka University, 3. JASRI

Keywords: Toroidal type DAC(t-DAC)

Oral presentation | T1: Comprehensive understanding of the crustal evolution and resource exploration in Asia (Symposium)

📅 Fri. Sep 13, 2024 9:00 AM - 12:00 PM JST | Fri. Sep 13, 2024 12:00 AM - 3:00 AM UTC | 🏢 ES Hall Higashiyama Campus

T1: Comprehensive understanding of the crustal evolution and resource exploration in Asia (Symposium)

Chairperson: Yasuhito Osanai, Masaaki Owada

9:00 AM - 9:25 AM JST | 12:00 AM - 12:25 AM UTC

[T1-01] Decarbonized Society and Essential Metal Resources

「招待講演」

*YOSHITAKA HOSOI¹ (1. JICA)

9:25 AM - 9:50 AM JST | 12:25 AM - 12:50 AM UTC

[T1-02] Japan's current approach to securing mineral resources

「招待講演」

*Kazuhiro YONEMURA¹ (1. JOGMEC)

9:50 AM - 10:15 AM JST | 12:50 AM - 1:15 AM UTC

[T1-03] Critical metal potentiality of Mongolia

「招待講演」

*Sereenen Jargalan¹, M. Arvinzun² (1. Mongolian University of Science and Technology, 2. Mongolian Society of Economic Geologists)

10:15 AM - 10:30 AM JST | 1:15 AM - 1:30 AM UTC

[2Lecture-101-06-4add] 休憩

10:30 AM - 10:55 AM JST | 1:30 AM - 1:55 AM UTC

[T1-04] Geology, sedimentation environment of the Ovoot khural coal bearing depression, in South Mongolia

「招待講演」

*Magsarjav Ochirbat², Sereenen Jargalan¹ (1. Mongolian University of Science and Technology, 2. Mongolian Society of Economic Geologists)

10:55 AM - 11:20 AM JST | 1:55 AM - 2:20 AM UTC

[T1-05] Insights into the mineralogical characteristics of Li-enriched metasomatic albitite from the Iwagi islet, SW Japan

「招待講演」

*Mariko NAGASHIMA¹, Teruyoshi IMAOKA¹ (1. Yamaguchi Univ. Sci.)

11:20 AM - 11:45 AM JST | 2:20 AM - 2:45 AM UTC

[T1-06] Ion adsorption-type REE deposits: the source of HREE

「招待講演」

*Yasushi Watanabe¹ (1. Akita Univ. Int. Res. Sci.)

11:45 AM - 12:00 PM JST | 2:45 AM - 3:00 AM UTC

[2Lecture-101-06-8add] 休憩

Decarbonized Society and Essential Metal Resources

*YOSHITAKA HOSOI¹

1. JICA

Measures to curb the rise in global temperature include energy conservation, low-carbon energy (promotion of the use of wind power, solar power generation, geothermal power generation, etc.), and conversion of energy use (electrification, use of hydrogen, etc.). Here, when trying to reduce the carbon value of energy, it became clear that special metals were needed in unusually large quantities. For example, solar power generation requires gallium and cadmium as solar cells, in addition to copper and aluminum. Wind turbines use generators that use permanent magnets composed of rare earth minerals such as neodymium and dysprosium. Geothermal power generation requires titanium for heat-resistant wells, and chromium is also needed for other technologies. Storage batteries are also needed for electric vehicles and wind power generation, but they also require lithium and vanadium. The demand for electric vehicles is expanding rapidly in various countries, and the demand for storage batteries will expand proportionally. According to the World Bank's 2020 report, if we forecast the amount of production required for 2050 compared to the production volume in FY2018, the amount of graphite 494%, lithium 488%, cobalt 460%, indium 231%, vanadium 189%, etc. It has become necessary. This is not the only metal needed. The World Bank lists 17 mineral types. As for the reserves, production, and consumption of these metals, the author considered the priority countries. Many of these resources are found in developing countries. Here, we consider the challenges of securing critical mineral resources. In addition, there are concerns that many of these limited producer countries are politically unstable, environmental pollution associated with mine development is a problem, and social turmoil occurs frequently. JICA is committed to solving the problems of resource-rich developing countries.

Keywords: Decarbonization, Essential metals, Mining challenges

Japan's current approach to securing mineral resources

*Kazuhiro YONEMURA¹

1. JOGMEC

The global green transformation (GX) is intensifying competition to secure Critical Minerals for batteries, semiconductors, and other applications. In particular, not only existing resource companies but also automakers and battery manufacturers around the world are accelerating their efforts to secure those material source to lithium, nickel, and graphite, which are used in electric vehicles. In some cases, there are concerns about economic security risks due to the ubiquity of supply source and midstream processes for these mineral resources.

Under these circumstances, efforts to diversify supply sources and midstream processes are being promoted worldwide, including financial support from governments and institutional design. There is also a growing movement to promote sustainable resource development by high level ESG standards. In Japan, based on the "Storage Battery Industrial Strategy" and the "Policy for Initiatives to Ensure Stable Supplies of Critical Minerals" based on the Economic Security Promotion Act, Japanese government supports Japanese companies investment for mine development, technology development and smelting-processing, specifically, increasing the ratio of financial support and providing subsidies for development and other activities through JOGMEC. In addition to these measurement, it is also actively conducting resource diplomacy with resource-rich countries and responding to multiple-frameworks. As for diversification of supply sources, while existing resources are being depleted, investment is concentrated on promising projects. As one solution, JOGEMC is focusing on ore minerals that have not been considered as resources (e.g., Awaruite) and areas where exploration has not progressed. Asia, which contains complex tectonics settings and remains un-exploration areas, has great potential of critical minerals.

Keywords: Critical Minerals, Securing Mineral Resources

Critical metal potentiality of Mongolia

*Sereenen Jargalan¹, M. Arvinzun²

1. Mongolian University of Science and Technology, 2. Mongolian Society of Economic Geologists

Critical metals such as copper, lithium, nickel, cobalt and rare earth elements are essential components in many of today's rapidly growing clean energy technologies –from wind turbines and electricity networks to electric vehicles. Lithium, nickel, cobalt, manganese and graphite are crucial to battery performance. rare earth elements are essential for permanent magnets used in wind turbines and EV motors. Mongolia has wide potential on mineral resources in variety of types.

Regarding to critical metal tendency, there is no clear classification in Mongolia, partly identify as high technology minerals and some government official documents use as important minerals. Even though some small projects are carried out to identify how potential is critical metals including REE, Li, Ni, Co as well as graphite in recent years.

Therefore, we carried out geological reconnaissance study to make clear genetic type, regional distribution characteristics and ore mineral identification. As result we have quite good potentiality on REE mineralization, including carbonatite and alkaline metasomatite types. Lithium is not so studied in Mongolia, but recently, we have several discoveries of Li bearing pegmatites in the central-eastern part. Nickel and cobalt are almost not studied instead of small occurrences found during geological mapping at scale 1:200000 and 1:50000, so no clear potentiality is recognized. There are several deposits and occurrence in Mongolia which are closely relate with marbles metamorphic rocks and has possible potentiality. Copper is the most potential resource making in all, almost 1 billion tons of reserves and resources. Annual production is expected to more than double from 300,000 tons of copper concentrate per year to over 600,000 tons per year from 2028 to 2036 once the Oyu Tolgoi mine is fully operational in 2023.

This time we would like to make general introduction of how potential is in critical metal tendency in Mongolia.

Keywords: Metal potentiality, Mongolia

Oral presentation

T1: Comprehensive understanding of the crustal evolution and resource exploration in Asia (Symposium)

Chairperson: Yasuhito Osanai, Masaaki Owada

Fri. Sep 13, 2024 9:00 AM - 12:00 PM ES Hall (Higashiyama Campus)

10:15 AM - 10:30 AM

[2Lecture-101-06-4add]休憩

Geology, sedimentation environment of the Ovoot khural coal bearing depression, in South Mongolia

*Magsarjav Ochirbat², Sereenen Jargalan¹

1. Mongolian University of Science and Technology, 2. Mongolian Society of Economic Geologists

Mongolia has abundant resources of coal, which are distributed mainly in the south and southeastern part, including some deposits in the north and west part of the country. In recent years, the exploration of coal deposits has been intensively carried out, but not much effort has been made to determine the origin, regional regularity of coal distribution as well as relationship between geological condition and coal quality.

The purpose of this study is to clarify geology and sedimentation environment of the Ovoot khural coal bearing depression in order to contribute to the reconstruction of the Mesozoic geologic and geodynamic setting of the Mongolia.

The study area is located at the 1000 km southwest of Ulaanbaatar and 50 km north of the Mongolia-China border, forming latitudinal trending depression which is 40 to 60 km in width and continues more than 200 km. There are five independent coal bearing parts with 16 coal deposits, which are: Bayantes part, contains Elstei, Khurshuut, Khuvd, Gashuu Tolgoi and Khuren tasv deposits; Ovoot Tolgoi part, contains Sunset and Sunrise deposits; Nariin sukhait part contains West Nariin sukhait, Central Nariin sukhait, East Nariin sukhait and Khuren shand deposits; Sumber part contains Central Sumber, Sumber and Biluut deposits and Jargalant part contains Jargalant and South Biluut deposits. Result of geochemical study indicates that sediments deposited in the Ovoot khural depression is sourced by the weathering and transporting of intermediate and felsic composition magmatic rocks, mainly from dacite, andesite including minor amount of metamorphic, sedimentary and intrusive rocks. Spider diagrams of trace element composition of sedimentary rocks of the Orgilokhbulag formation, show Nb-depletion and Pb, Mo enrichment, indicating possible origin of magmatic rocks formed under subduction environment and they are intermediate to felsic in composition. The major trace and rare earth element composition of the Orgilokh bulag formation sedimentary rocks, indicate that the source rocks of sediments might have been formed in the active continental margin tectonic setting. According to provenance model, coal deposition is undertaking with good tissue preservation, in an alternating environment of oxygenic and deoxygenated swamps. Based on the metamorphic degree, the temperature of peat compression, the amount of volatile, and the depth gradient of temperature, peat was buried and deposited at a depth of 3500-7000 meters. Low sulfur content, low ash content and low volatile content of coal at the various parts of the depression indicates that the peat deposition is occurred under two stages.

Keywords: Ovoot khural coal, Mongolia

Insights into the mineralogical characteristics of Li-enriched metasomatic albitite from the Iwagi islet, SW Japan


*Mariko NAGASHIMA¹, Teruyoshi IMAOKA¹

1. Yamaguchi Univ. Sci.

The study of metasomatic rocks is essential for comprehending the nature and origin of metasomatic agents. It might provide a clue for fluid circulation within the crust. In the Setouchi Province, metasomatic rocks are distributed along an approximately E–W trend, and these rocks are associated with Late Cretaceous granites. The Iwagi albitite is one such metasomatic rock. However, unlike other metasomatites in the area, its high lithium content (500 ppm) is unique. Detailed investigations of the mineralogical characteristics of Li-minerals have been conducted to better understand the formation and evolution of Iwagi albitites. The albitites exist as small masses, and the textures of the weakly metasomatized ones resemble those of the host adjacent granite. The transition from granite to albitite occurs gradually and can be understood through the mineral assemblages. The Iwagi albitite is known as the type locality of four Li-analog minerals: sugilite $\text{KNa}_2(\text{Fe}^{3+}, \text{Mn}^{3+}, \text{Al})_2\text{Li}_3\text{Si}_{12}\text{O}_{30}$, katayamalite $\text{KLi}_3\text{Ca}_7\text{Ti}_2(\text{SiO}_3)_{12}(\text{OH})_2$, murakamiite $\text{LiCa}_2\text{Si}_3\text{O}_8(\text{OH})$, and ferro-ferri-holmquistite $\text{Li}_2(\text{Fe}^{2+}_3\text{Fe}^{3+}_2)\text{Si}_8\text{O}_{22}(\text{OH})_2$. The former three minerals were found in the fully albitized rock, while the latter was found in the weakly albitized granite. The albitites display a variety of replacement textures due to Na–Li metasomatism, and they also exhibit noticeable strain-induced textures.

The $\delta^7\text{Li}$ values of murakamiite and Li-rich pectolite show a wide range from -9.1 to +0.4‰ (ave. -2.9‰) and should have resulted from hydrothermal fluid-rock interactions at 300–600 °C. The very low $\delta^7\text{Li}$ values may have originated from intra-crystalline Li isotope diffusion or involvement of deep-seated, Li–Na-enriched subduction-zone fluids with low $\delta^7\text{Li}$ values. This finding highlights the significance of fluid-rock interactions in the formation of metasomatic rocks. Deformation-induced fracturing of the rock may have enhanced fluid circulation, leading to the formation of the metasomatic rocks along the E–W trending lineament.

Keywords: Lithium, albitite, metasomatism



Distribution of metasomatites in Setouchi Province, SW Japan.
(after Murakami 1976)

Ion adsorption-type REE deposits: the source of HREE

*Yasushi Watanabe¹

1. Akita Univ. Int. Res. Sci.

Ion adsorption rare earth deposits were confirmed in southern China including Jiangxi province in late 1970's. This unique deposit type forms by adsorption of rare earth ions on clay minerals represented by kaolinite and halloysite due to weathering of granitic rocks. Although the ore grades of this deposit type is extremely lower (<0.2 wt%) than the other rare earth deposits such as carbonatite, extraction of rare earths from the clay ores is easy and inexpensive. The development of this deposit type has been accelerating since 2000 as the source of heavy rare earths. This is due to the invention of neodymium magnet in 1983, followed by commercialization in 1985, and production of hybrid vehicle (Prius) in 1997. Because major rare earth deposits such as carbonatite and placer deposits are enriched in LREE but poor in HREE, the ion adsorption type deposits became the important HREE supply source. Although exploration of HREE prospects has been conducted worldwide and a few HREE enriched alkaline-rock related deposits were discovered, no deposit is better than the ion adsorption deposits in terms of production cost and easiness in processing. The ion adsorption type deposits are distributed not only in southern China but also in southeast Asia. This type of deposits also present in southern Africa including Malawi and Madagascar and South America such as Brazil and Chile. Presently Myanmar has become the major country that produces ionic ores. For the formation of ion adsorption HREE deposits needs the following three conditions; 1) presence of HREE enriched host rocks, 2) formation of thick (>10 m) weathering crust, and 3) presence of REE minerals in the host rocks that easily dissolve during weathering.

Keywords: ion adsorption-type deposit, heavy rare earth elements, weathering, magnet

Oral presentation

T1: Comprehensive understanding of the crustal evolution and resource exploration in Asia (Symposium)

Chairperson: Yasuhito Osanai, Masaaki Owada

Fri. Sep 13, 2024 9:00 AM - 12:00 PM ES Hall (Higashiyama Campus)

11:45 AM - 12:00 PM

[2Lecture-101-06-8add]休憩

Oral presentation | S2: Water Rock Interaction (Special Session)

📅 Fri. Sep 13, 2024 9:00 AM - 12:00 PM JST | Fri. Sep 13, 2024 12:00 AM - 3:00 AM UTC | 🏠 ES024
Higashiyama Campus

S2: Water Rock Interaction (Special Session)

Chairperson: Noriyoshi Tsuchiya

9:00 AM - 9:05 AM JST | 12:00 AM - 12:05 AM UTC

[2Lecture-201-10-1add] コンビーナ挨拶

9:05 AM - 9:35 AM JST | 12:05 AM - 12:35 AM UTC

[S2-01] Slab-derived fluid infiltrating back-arc mantle

「招待講演」

*Junji YAMAMOTO¹ (1. Kyushu University)

9:35 AM - 9:50 AM JST | 12:35 AM - 12:50 AM UTC

[S2-02] Geochemical diversity and significance of orthopyroxene pseudomorphs in ultramafic rocks derived from mantle wedges

「発表賞エントリー」

*Takumi Wani¹, Yuji Ichiyama¹, Akihiro Tamura², Tomoaki Morishita² (1. Chiba University, 2. Kanazawa University)

9:50 AM - 10:05 AM JST | 12:50 AM - 1:05 AM UTC

[S2-03] Petrogenesis and significance of ophicarbonates in the Kanasaki serpentinite body (Kanto Mountains, Central Japan)

*Ryosuke OYANAGI^{1,2}, Hikaru Sawada^{3,2}, Qing Chang², Madhusoodhan Satish-Kumar⁴ (1. Kokushikan Univ., 2. JAMSTEC, 3. Toyama Univ., 4. Niigata Univ.)

10:05 AM - 10:20 AM JST | 1:05 AM - 1:20 AM UTC

[S2-04] CO₂ mineralization in andesitic rocks revealed by hydrothermal experiments and thermal analyses

*Otgonbayar DANDAR¹, Atsushi Okamoto¹, Masaaki Uno¹, Miku Takeya² (1. Tohoku University, 2. INPEX)

10:20 AM - 10:30 AM JST | 1:20 AM - 1:30 AM UTC

[2Lecture-201-10-6add] 休憩

10:30 AM - 10:45 AM JST | 1:30 AM - 1:45 AM UTC

[S2-05] Massive and foliated serpentinites from the Udonohana ultramafic body, Western Ehime Prefecture, Japan.

「発表賞エントリー」

*Yui Joguchi¹, Satomi Enju¹ (1. Ehime Univ. Sci. & Egn.)

10:45 AM - 11:00 AM JST | 1:45 AM - 2:00 AM UTC

[S2-06] "Element transport and magnetite decomposition during alteration of the gabbroic vein in serpentinite body from the Bayankhongor ophiolite, Mongolia"

「発表賞エントリー」

*Nomin Tumurkhuu¹, Otgonbayar Dandar¹, Masaaki Uno¹, Manzshir Bayarbold¹, Atsushi Okamoto¹ (1. Tohoku University)

11:00 AM - 11:15 AM JST | 2:00 AM - 2:15 AM UTC

[S2-07] Dissolution reprecipitation - re-equilibration process of feldspar in heat source granite and supercritical geothermal reservoir using borehole samples from Kakkonda granite

「発表賞エントリー」

*Masayoshi Hoshida¹, Masaoki Uno¹, Satoshi Matsuno¹, Astin Nurdiana¹, Noriyoshi Tsuchiya^{2,1} (1. Tohoku University, 2. National Institute of Technology, Hachinohe College)

11:15 AM - 11:30 AM JST | 2:15 AM - 2:30 AM UTC

[S2-08] Formation mechanism of "cleavable olivine"

*Jun-ichi ANDO^{1,2}, Naotaka Tomioka^{3,2}, Hirokazu Maekawa⁴ (1. Hiroshima Univ., 2. Hiroshima Univ., HiPeR, 3. JAMSTEC, 4. Osaka Metropolitan Univ.)

11:30 AM - 11:45 AM JST | 2:30 AM - 2:45 AM UTC

[S2-09] Experimental study to elucidate sulfide chimney development process and power generation characteristics in submarine hydrothermal systems

「発表賞エントリー」

*Kentarō Toda¹, Atsushi Okamoto¹, Dandar Otgonbayar¹, Misaki Takahashi¹, Yoshinori Sato¹ (1. Tohoku Univ. Environmental Sci)

11:45 AM - 12:00 PM JST | 2:45 AM - 3:00 AM UTC

[S2-10] Carbonation of Mantle Peridotite: An Approach From Fluid Inclusion Analysis and Hydrothermal Experiments

*Tatsuhiko KAWAMOTO¹ (1. Shizuoka University)

Oral presentation

S2: Water Rock Interaction (Special Session)

Chairperson: Noriyoshi Tsuchiya

Fri. Sep 13, 2024 9:00 AM - 12:00 PM ES024 (Higashiyama Campus)

9:00 AM - 9:05 AM

[2Lecture-201-10-1add] コンビーナ挨拶

Slab-derived fluid infiltrating back-arc mantle

*Junji YAMAMOTO¹

1. Kyushu University

Keywords: slab, mantle transition zone, nitrogen isotope ratio, xenolith, fluid inclusion




Fig. 1. $\text{N}_2/^{36}\text{Ar}$ vs. $\delta^{15}\text{N}$ of Far Eastern Russian xenoliths with data referred from earlier studies (see Yamamoto et al. (2020) EPSL).

Geochemical diversity and significance of orthopyroxene pseudomorphs in ultramafic rocks derived from mantle wedges

*Takumi Wani¹, Yuji Ichiyama¹, Akihiro Tamura², Tomoaki Morishita²

1. Chiba University, 2. Kanazawa University

Keywords: mantle wedge, mantle metasomatism, slab fluid, orthopyroxene pseudomorph

Petrogenesis and significance of ophicarbonates in the Kanasaki serpentinite body (Kanto Mountains, Central Japan)

*Ryosuke OYANAGI^{1,2}, Hikaru Sawada^{3,2}, Qing Chang², Madhusoodhan Satish-Kumar⁴

1. Kokushikan Univ., 2. JAMSTEC, 3. Toyama Univ., 4. Niigata Univ.

Keywords: Serpentine, Ophicarbonate, Carbon cycle

CO₂ mineralization in andesitic rocks revealed by hydrothermal experiments and thermal analyses

*Otgonbayar DANDAR¹, Atsushi Okamoto¹, Masaoki Uno¹, Miku Takeya²

1. Tohoku University, 2. INPEX

Mineral carbonation enables long-term stable CO₂ storage. Among rocks, basalts and ultramafic rocks (high Mg and Ca contents) are known to have high potential to store CO₂. Although, andesitic rocks (composition is wide; microstructures are diverse) are widely distributed in Japan (in a subduction zone), their suitability for mineral carbonation and CO₂ storage is not well understood. In this study, we report preliminary results of hydrothermal reaction experiments (exps) using andesitic agglomerate rocks from the Nagaoka area, Niigata Prefecture and thermal analyses of the products. The andesite agglomerate consists mainly of volcanic glass, plagioclase, clinopyroxene, with small amounts of orthopyroxene, hornblende, and iron oxides. Two types (a batch-type vessel and a stirred apparatus) of exps have been conducted with the powder samples (150-250 μm). In the batch exp (filling ratio = 43% and remained space is CO₂ gas), powder sample is placed at two locations: the top (supercritical CO₂ saturated in water) and bottom (reacted with a CO₂ dissolved solution) of the reaction vessel. The batch exps were conducted at 75 °C, 18 MPa (80 days), 150 °C, 20 MPa (20 days), and 200 °C, 25 MPa (20, 40, and 60 days). The solid phase after exp is observed by EPMA, TG, and TPD-MS, and the solutions are analyzed for element contents by ICP-OES. To accelerate the reaction, a stirred reaction apparatus was used for the exp at 200 °C, 20MPa, and 10 days. In the batch exps at 75 °C and 150 °C (Ca and Mg+Fe contents = 10-80 mg/kgH₂O), dissolution of volcanic glass is observed only with small dissolution of plagioclase and other minerals. Significant carbonate precipitation was observed in a 60-day batch exp at 200 °C (Fe- and Ca-rich magnesite) and a 10-day stirred exp (calcite and dolomite). The precipitation proceeded not only at the bottom (CO₂ dissolved solution), but also at the top (supercritical CO₂), suggesting the reaction accelerated by water vapor. The experimental solution (carbonate precipitated, the pH increased to >7 and the concentrations of Mg and Ca <20 mg/kgH₂O), suggests the pH in the reaction vessel increased and reached to condition of carbonate precipitation. Based on the above exps, the andesitic rocks in the Nagaoka area are considered to have sufficient potential for mineral carbonation and CO₂ storage.

Keywords: Mineral carbonation , CO₂ storage, Andesite, Hydrothermal reaction experiment, the Nagaoka area, Niigata Prefecture

Oral presentation

S2: Water Rock Interaction (Special Session)

Chairperson: Noriyoshi Tsuchiya

Fri. Sep 13, 2024 9:00 AM - 12:00 PM ES024 (Higashiyama Campus)

10:20 AM - 10:30 AM

[2Lecture-201-10-6add]休憩

Massive and foliated serpentinites from the Udonohana ultramafic body, Western Ehime Prefecture, Japan.

*Yui Joguchi¹, Satomi Enju¹

1. Ehime Univ. Sci. &Egn.

Keywords: serpentinite, ultramafic rocks, cumulate, Mikame ultramafic body

"Element transport and magnetite decomposition during alteration of the gabbroic vein in serpentinite body from the Bayankhongor ophiolite, Mongolia"

*Nomin Tumurkhuu¹, Otgonbayar Dandar¹, Masaoki Uno¹, Manzshir Bayarbold¹, Atsushi Okamoto¹

1. Tohoku University

Hydration of mantle rocks is important for elucidating the dynamics of global water circulation, elemental transport, and geological processes. The crust-mantle interface constitutes a significant geological boundary where element transport occurs. However, research on the multi-stage hydration of mantle rocks within this zone, particularly in the oceanic lithosphere (mid-ocean ridge), is still lacking. Therefore, to understand element transport during the hydration of the crustal vein in the mantle at the crust-mantle transition zone, we investigate the mantle-crust section preserved in the Bayankhongor ophiolite (BO; mid-ocean ridge origin) Mongolia. The outcrop of the crust-mantle section (~30 m in diameter) in the BO is characterized by a brownish gabbroic body with a massive and sheared mantle body fully serpentinitized. Mantle rock samples mainly consist of lizardite in two forms: mesh core (Mg# = 0.95-0.98) with fine magnetite (Mgt) and vein (Mg# = 0.94-0.98) with vein Mgt (<30 μ m width), along with spinel (Mg# = 0.42-0.52 & Cr# = 0.46-0.48), and chlorite (Chl; Mg# = 0.87-0.96). The absence of brucite in the serpentinites suggests infiltration of Si-rich fluids. Green veins (80-95 cm in width; it mainly consists of clinopyroxene (Cpx; Mg# = 0.92) replaced by a mixture of Chl-serpentine (Srp) and cut by serpentine and epidote (Ep) veins), along with white veins (~15 cm in width; ~40 cm long; it is mostly consisted of Ep and Cpx with a minor amount of Chl) cut through the mantle rocks. Additionally, black veins (~2 cm in width; it is composed of Chl patches (Mg# = 0.83-0.93) and Chl-Srp patches with clear cleavages and fine Ti-rich minerals) intersect the serpentinite. The reaction zone (~3 mm) between host serpentinite and black vein shows that Mgt disappeared and Mgt is replaced by Al-rich (1.1-6.9 wt%) Srp. Mass balance on black vein (assuming protoliths: Cpx for Chl-Srp and plagioclase for Chl patch shows gain of Fe and Mg, and loss of Si, Al, and Ca whereas that on the reaction zone shows loss of Fe and gain of Si, and Al. This implies that Mg-rich fluid and chl formation cause Mgt disappearance and mobility of Fe, Si, and Al. Reaction zone and mass balance result imply that local mobility of Si, Al, Fe, Mg, and Ca could occur at the crust-mantle section in the oceanic lithosphere during multi-stage hydration.

Keywords: Element transport, Serpentine, Gabbroic vein

Element transport and magnetite decomposition during alteration of gabbroic vein in serpentinite body from the Bayankhongor ophiolite, Mongolia.

Nomin TUMURKHUU, Otgonbayar DANDAR, Masaoki UNO, Manzshir BAYARBOLD, Atsushi OKAMOTO

Graduate School of Environmental Studies, Tohoku University

Hydration of mantle rocks is important for elucidating the dynamics of global water circulation, elemental transport, and geological processes. The crust-mantle interface constitutes a significant geological boundary where element transport occurs. However, research on the multi-stage hydration of mantle rocks within this zone, particularly in the oceanic lithosphere (mid-ocean ridge), is still lacking. Therefore, to understand element transport during the hydration of the crustal vein in the mantle at the crust-mantle transition zone, we investigate the mantle-crust section preserved in the Bayankhongor ophiolite (BO; mid-ocean ridge origin) Mongolia.

The outcrop of the crust-mantle section (~30 m in diameter) in the BO is characterized by a brownish gabbroic body with a massive and sheared mantle body fully serpentinitized. Mantle rock samples mainly consist of lizardite in two forms: mesh core ($Mg\# = 0.95-0.98$) with fine magnetite (Mgt) and vein ($Mg\# = 0.94-0.98$) with vein Mgt (<30 μm width), along with spinel ($Mg\# = 0.42-0.52$ & $Cr\# = 0.46-0.48$), and chlorite (Chl; $Mg\# = 0.87-0.96$). The absence of brucite in the serpentinites suggests infiltration of Si-rich fluids. Green veins (80-95 cm in width; it mainly consists of clinopyroxene (Cpx; $Mg\# = 0.92$) replaced by a mixture of Chl-serpentine (Srp) and cut by serpentine and epidote (Ep) veins), along with white veins (~15 cm in width; ~40 cm long; it is mostly consisted of Ep and Cpx with a minor amount of Chl) cut through the mantle rocks. Additionally, black veins (~2 cm in width; it is composed of Chl patches ($Mg\# = 0.83-0.93$) and Chl-Srp patches with clear cleavages and fine Ti-rich minerals) intersect the serpentinite.

The reaction zone (~3 mm) between host serpentinite and black vein shows that Mgt disappeared and Mgt is replaced by Al-rich (1.1-6.9 wt%) Srp. Mass balance on black vein (assuming protoliths: Cpx for Chl-Srp and plagioclase for Chl patch shows gain of Fe and Mg, and loss of Si, Al, and Ca whereas that on the reaction zone shows loss of Fe and gain of Si, and Al. This implies that Mg-rich fluid and chl formation cause Mgt disappearance and mobility of Fe, Si, and Al. Reaction zone and mass balance result imply that local mobility of Si, Al, Fe, Mg, and Ca could occur at the crust-mantle section in the oceanic lithosphere during multi-stage hydration.

Dissolution reprecipitation - re-equilibration process of feldspar in heat source granite and supercritical geothermal reservoir using borehole samples from Kakkonda granite

*Masayoshi Hoshida¹, Masaaki Uno¹, Satoshi Matsuno¹, Astin Nurdiana¹, Noriyoshi Tsuchiya^{2,1}

1. Tohoku University, 2. National Institute of Technology, Hachinohe College

Keywords: Feldspar, Feldspar thermometer, Supercritical geothermal reservoir, Dissolution reprecipitation, Re-equilibration

Formation mechanism of "cleavable olivine"

*Jun-ichi ANDO^{1,2}, Naotaka Tomioka^{3,2}, Hirokazu Maekawa⁴

1. Hiroshima Univ., 2. Hiroshima Univ., HiPeR, 3. JAMSTEC, 4. Osaka Metropolitan Univ.

Keywords: cleavable olivine, Subgrain boundary, Pipe diffusion

Experimental study to elucidate sulfide chimney development process and power generation characteristics in submarine hydrothermal systems

*Kentaro Toda¹, Atsushi Okamoto¹, Dandar Otgonbayar¹, Misaki Takahashi¹, Yoshinori Sato¹

1. Tohoku Univ. Environmental Sci

Keywords: Hydrothermal Chimney

Carbonation of Mantle Peridotite: An Approach From Fluid Inclusion Analysis and Hydrothermal Experiments

*Tatsuhiko KAWAMOTO¹

1. Shizuoka University

Carbonate rocks in serpentinites are called ophicarbonates. Two types of ophicarbonates are observed as carbonate veins in serpentinite and serpentinite breccias surrounded by carbonate matrix. At Shizuoka University, Japan, we have described and reported saline fluid inclusions in carbonate minerals in ophicarbonates at the localities as follows: (1) carbonate veins in serpentinites of the Oman ophiolite, formed at a fast spreading ridge; (2) carbonate veins and carbonate matrix in serpentinites in the Western Alps ophiolite, formed at a slow spreading ridge; (3) calcite in serpentinite mud volcanoes in the Izu Mariana Trench; and (4) carbonate veins in mantle-wedge serpentinite in the Sambagawa Belt, Chichibu, Japan. The results show that the studied fluid inclusions have salinity similar to that of seawater or slightly higher than seawater, except for the fresh water fluid inclusions found in the later veins of the Oman ophiolite serpentinite in (1), where fluid inclusions with salinity similar to seawater are present in veins formed below the seafloor before. This means that ophicarbonates formed by metamorphism beneath the seafloor seen in (1) and (2) involve saline fluids similar to seawater, and such saline fluids are also brought into the mantle wedge, where lithospheres sink seen in (3) and (4).

Hydrothermal experiments are also conducted to provide constraints on the carbonization of the Earth's mantle. In addition to the hydrothermal experiments, we calculate phase diagrams by use of thermodynamic calculation such as Perple_X. If we will experimentally determine the equilibrium mineral assemblage in iron-bearing system, we can understand the effect of iron on the temperature and pressure conditions by comparing the calculated phase equilibrium in the CaO-MgO-SiO₂-H₂O-CO₂ system. The comparison between natural ophicarbonates and the calculated phase diagram also enables us to know the CO₂/(H₂O+CO₂) molar ratio of the fluids in the carbonation. It will be also possible to constraint on the temperature and pressure conditions from the isochore of fluid inclusions. Combination of natural observation, experiments and calculation allows us to understand the conditions of mantle carbonation more precisely.

Keywords: fluid inclusions, serpentinite, seawater, carbon dioxide, mantle

Oral presentation | S3: Rheology and Material Transfer in Mantle and Crust (Special Session)

📅 Fri. Sep 13, 2024 9:00 AM - 10:00 AM JST | Fri. Sep 13, 2024 12:00 AM - 1:00 AM UTC | 🏠 ES025
Higashiyama Campus

S3: Rheology and Material Transfer in Mantle and Crust (Special Session)

Chairperson: Miki Tasaka (Shizuoka University)

9:00 AM - 9:15 AM JST | 12:00 AM - 12:15 AM UTC

[S3-10] Rheological evolution of olivine during formation of the mantle lithosphere

*Katsuyoshi MICHIBAYASHI^{1,2}, Takeo Okuwaki¹, Itsuki Natsume³ (1. Nagoya University, 2. JAMSTEC, 3. Kanagawa Prefectural Museum of Natural History)

9:15 AM - 9:30 AM JST | 12:15 AM - 12:30 AM UTC

[S3-11] Three-dimensional location analysis on acoustic emissions and faults in olivine under pressure-temperature conditions of subducting slabs

*Tomohiro OHUCHI¹, Masato Hoshino², Kentaro Uesugi², Satoshi Okumura³, Yuji Higo², Noriyoshi Tsujino², Sho Kakizawa² (1. GRC, Ehime Univ., 2. JASRI, 3. Tohoku Univ Sci.)

9:30 AM - 9:45 AM JST | 12:30 AM - 12:45 AM UTC

[S3-12] In-situ observation of grain growth and fluid movement using camphor as a rock analogue

*Junichi Fukuda¹ (1. Dept. Geos. Osaka Metrop. Univ.)

9:45 AM - 10:00 AM JST | 12:45 AM - 1:00 AM UTC

[S3-13] Mantle carbonation through seawater penetration along the outer-rise faults

*Ikuo KATAYAMA¹, Keishi Okazaki¹, Atsushi Okamoto² (1. Hiroshima University, 2. Tohoku University)

Rheological evolution of olivine during formation of the mantle lithosphere

*Katsuyoshi MICHIBAYASHI^{1,2}, Takeo Okuwaki¹, Itsuki Natsume³

1. Nagoya University, 2. JAMSTEC, 3. Kanagawa Prefectural Museum of Natural History

Keywords: Olivine, Mantle

Three-dimensional location analysis on acoustic emissions and faults in olivine under pressure-temperature conditions of subducting slabs

*Tomohiro OHUCHI¹, Masato Hoshino², Kentaro Uesugi², Satoshi Okumura³, Yuji Higo², Noriyoshi Tsujino², Sho Kakizawa²

1. GRC, Ehime Univ., 2. JASRI, 3. Tohoku Univ Sci.


Keywords: Olivine, Acoustic emission, Fault, Intermediate earthquake

In-situ observation of grain growth and fluid movement using camphor as a rock analogue

*Junichi Fukuda¹

1. Dept. Geos. Osaka Metrop. Univ.

Keywords: Rock analogue, Camphor, Grain growth, Fluid



Mantle carbonation through seawater penetration along the outer-rise faults

*Ikuo KATAYAMA¹, Keishi Okazaki¹, Atsushi Okamoto²

1. Hiroshima University, 2. Tohoku University

Keywords: Carbon cycle, Carbonation, Mantle

Oral presentation | R4: Mineral sciences of the Earth surface

📅 Fri. Sep 13, 2024 10:15 AM - 12:00 PM JST | Fri. Sep 13, 2024 1:15 AM - 3:00 AM UTC | 🏠 ES025
Higashiyama Campus

R4: Mineral sciences of the Earth surface

Chairperson: Hiroshi Sakuma(NIMS), Satoko Motai(Yamagata Univ.), Jun Kawano(Hokkaido University)

10:15 AM - 10:35 AM JST | 1:15 AM - 1:35 AM UTC

[R4-01] Understanding and application of spherical concretions: A new durable sealing material learnt from nature

「招待講演」

*Hidekazu Yoshida¹ (1. Nagoya University)

10:35 AM - 10:50 AM JST | 1:35 AM - 1:50 AM UTC

[R4-02] Color change process of Hiroshima granite due to weathering

*Tadashi YOKOYAMA¹, Yuka Inkyo, Masahiro Kaibori¹ (1. Hiroshima University)

10:50 AM - 11:05 AM JST | 1:50 AM - 2:05 AM UTC

[R4-03] Crystallographic preferred orientation and grain size of apatite in terrestrial mammalian bones

*Kyoko N. MATSUKAGE¹, Momoka Ide², Masaya Kurata², Yu Nishihara³ (1. Teikyo Univ. of Sci. Natural and Environmental Sci., 2. Teikyo Univ. of Sci. Ainal Sci., 3. Ehime Univ.)

11:05 AM - 11:20 AM JST | 2:05 AM - 2:20 AM UTC

[R4-04] Microscopic distribution of sodium in biogenic aragonite

*Taiga Okumura¹, Michio Suzuki², Alberto Perez-Huerta³, Eshita Samajpati³, Toshihiro Kogure¹ (1. UTokyo Sci., 2. UTokyo Agri. Life Sci., 3. Univ. Alabama Geol. Sci.)

11:20 AM - 11:40 AM JST | 2:20 AM - 2:40 AM UTC

[R4-05] Structural and functional analyses of organic matrices regulating the formation of minerals in biomineralization.

「招待講演」

*Michio Suzuki¹ (1. UTokyo)

11:40 AM - 11:55 AM JST | 2:40 AM - 2:55 AM UTC

[R4-06] Aragonite formation from amorphous calcium carbonate (ACC) with addition of *n*-butylamine

*Hiroyuki KAGI¹, Kensuke Muraoka¹ (1. The University of Tokyo)

11:55 AM - 12:00 PM JST | 2:55 AM - 3:00 AM UTC

調整

Understanding and application of spherical concretions: A new durable sealing material learnt from nature

*Hidekazu Yoshida¹

1. Nagoya University

Here, we introduce a more durable sealing method for concretion-forming resin developed by learning from natural calcite, CaCO_3 spherical concretion formation. The method was tested by sealing flow paths next to a tunnel in an underground research laboratory at 350 m depth, in Hokkaido, Japan. The flow paths were initially sealed rapidly, then resealed after disturbance by repeated earthquakes just below the underground research laboratory at depths of 2–7 km and maximum magnitude Mw 5.4. The treated rock mass rapidly recovered its very low natural permeability, demonstrating robust self-sealing and healing.

Keywords: Spherical concretion, Calcium carbonate, Durable sealing material

Color change process of Hiroshima granite due to weathering

*Tadashi YOKOYAMA¹, Yuka Inkyo, Masahiro Kaibori¹

1. Hiroshima University

The characteristics and mechanisms of color change associated with weathering of Hiroshima granite were investigated, using a core drilled to a depth of 20 m. Color (L^* a^* b^* values) were measured at approximately every 2-10 cm at each depth using a spectrophotometer. The color is: brighter for larger L^* values, more reddish for larger a^* values (for $a^* > 0$), and more yellowish for larger b^* values (for $b^* > 0$). For core color, whitish unweathered areas at depths greater than 12 m have smaller a^* and b^* values and larger L^* values. From about 12 m to 4 m depth, both a^* and b^* values increase and L^* values decrease closer to the surface, although there is some variation depending on the location. The color at each depth was compared with that of a reference material made from four typical iron-bearing secondary minerals (goethite (yellow), ferrihydrite (dark brown), lepidocrocite (light brown), and hematite (dark red)), where each mineral mixed with SiO_2 powder in various proportions from 0-100%. In the early stage of weathering, a dark brown band close to the color of ferrihydrite is often seen around the fractures. For more weathered areas, the entire rock matrix is often yellowish, the color close to that of goethite. In general, ferrihydrite is known to transform to more stable goethite and hematite with time. From the color measurement results of the drilled cores, it is inferred that ferrihydrite was first formed in the early stage of weathering, and then changed from ferrihydrite to goethite with the progress of weathering. Secondary iron minerals were dissolved and quantified by the selective iron dissolution method at several locations with different degrees of weathering. The amount of secondary iron minerals (all assumed to be goethite) at each location was estimated by comparing the color of the core with that of the reference material and plotted against the values obtained by the selective iron dissolution method, showing a generally proportional relationship. Although the results may change after correction for the effect of ferrihydrite, the results indicate that it may be possible to roughly estimate the amount of secondary iron minerals from a quick and easy core color measurement alone.

Keywords: Granite, Weathering, Color

Crystallographic preferred orientation and grain size of apatite in terrestrial mammalian bones

*Kyoko N. MATSUKAGE¹, Momoka Ide², Masaya Kurata², Yu Nishihara³

1. Teikyo Univ. of Sci. Natural and Environmental Sci., 2. Teikyo Univ. of Sci. Ainal Sci., 3. Ehime Univ.

Keywords: Apatite, terrestrial mammal, Bone, Pregerred orientation, Grain size

Microscopic distribution of sodium in biogenic aragonite

*Taiga Okumura¹, Michio Suzuki², Alberto Perez-Huerta³, Eshita Samajpati³, Toshihiro Kogure¹

1. UTokyo Sci., 2. UTokyo Agri. Life Sci., 3. Univ. Alabama Geol. Sci.

Keywords: Aragonite, Sodium, Biomineralization, STEM-EDS, Atom probe tomography

Structural and functional analyses of organic matrices regulating the formation of minerals in biomineralization.

*Michio Suzuki¹

1. UTokyo

Keywords: biomineralization, organic matrices, calcium carbonate, *Pinctada fucata*


Aragonite formation from amorphous calcium carbonate (ACC) with addition of *n*-butylamine

*Hiroyuki KAGI¹, Kensuke Muraoka¹

1. The University of Tokyo

Direct conversion from amorphous calcium carbonate (ACC) to aragonite has been extremely difficult compared to the other two polymorphs, calcite and vaterite. In the present study, aragonite formation with a high polymorph fraction (>97%) was obtained from ACC immersed in *n*-butylamine under 90% RH (relative humidity) at 30°C for two hours. It is noteworthy that the aragonite with high purity was obtained without the addition of Mg²⁺ ion, which is well known to promote aragonite formation. To understand the effects of hydrophobic and basic properties of *n*-butylamine, hexane, NH₃ aq., and hexane + NH₃ aq. were mixed with ACC and left for two weeks. Aragonite was formed only from the mixture of hexane and NH₃ aq. This result indicates that both hydrophobic and basic properties are required for crystallization from ACC into aragonite.

Keywords: amorphous calcium carbonate, aragonite, polymorph control



Poster presentation | T1: Comprehensive understanding of the crustal evolution and resource exploration in Asia (Symposium)

🏠 Fri. Sep 13, 2024 12:30 PM - 2:00 PM JST | Fri. Sep 13, 2024 3:30 AM - 5:00 AM UTC | 🏠 Entrance Hall Higashiyama Campus

T1: Comprehensive understanding of the crustal evolution and resource exploration in Asia (Symposium)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[T1-P-01] Syenitic veining at the northern Eastern Ghats Belt, India: Formation mechanism, fluid-rock interaction and a review of its economic mineral potential

*Kaushik DAS^{1,5}, Proloy Ganguly², Aparupa Banerjee³, Sankar Bose^{4,5} (1. Hiroshima University, 2. Kazi Nazrul University, 3. Shahid Matangini Hazra Government General Degree College, 4. Presidency University, 5. HiPeR, Hiroshima University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[T1-P-02] Amphibole-bearing granitic rocks in the Ereendavaa block, NE Mongolia: Insights into multi-stage magmatic sources and crystallization conditions

*Munkhdelger Bold¹, Tatsuki Tsujimori¹, Yasuhito Osanai², Nobuhiko Nakano², Tatsuro Adachi², Otgonbayar Dandar¹, Fransiska Ayuni Catur Wahyuandari² (1. Tohoku Univ., 2. Kyushu Univ)

Syenitic veining at the northern Eastern Ghats Belt, India: Formation mechanism, fluid-rock interaction and a review of its economic mineral potential

*Kaushik DAS^{1,5}, Proloy Ganguly², Aparupa Banerjee³, Sankar Bose^{4,5}

1. Hiroshima University, 2. Kazi Nazrul University, 3. Shahid Matangini Hazra Government General Degree College, 4. Presidency University, 5. HiPeR, Hiroshima University

Extensive development of meter-to-micrometer scale felsic bands and veins at the contact between the calc-silicate granulite (clinopyroxene-plagioclase-scapolite-wollastonite-calcite-grandite garnet-titanite-apatite-quartz), charnockite (plagioclase-K-feldspar-orthopyroxene-ilmenite-quartz± garnet), and mafic granulite (clinopyroxene-orthopyroxene-plagioclase±garnet-hornblende±quartz) is observed at a 50-60 km wide area near the northern boundary (south of Mahanadi shear zone) of the Eastern Ghats Province of the Eastern Ghats Belt, India. The calc-silicate and mafic granulites witnessed HT to UHT metamorphism at ca. 1000 Ma, and two pulses of charnockite magmatism between ca. 970 Ma and 950 Ma. Clinopyroxene-bearing syenitic (clinopyroxene-K-feldspar-titanite-REE-phases-minor quartz) bands and veins occur in these rocks sometimes at the rock interface and even as a vein network inside one of the preexisting rocks. Inside the veins close to their wall clinopyroxene (+titanite) formed at the expense of orthopyroxene (+ilmenite). Anorthitic patches and myrmekite intergrowth replaced the albitic plagioclase of the wall zone of charnockite. All these reactions suggest selective mobility of at least Ca and Si in the vein with a definite gradient of these from the interior to the wall. Primary and secondary fluid inclusions in the vein mineralogy are rich in CO₂. Compositional variation in apatite from the host rock to the vein also shows scavenging of REE and new profuse formation of REE-rich megacrystic allanite, titanite, apatite, and zircon inside the vein. The contact rocks of charnockite show reactions of monazite to allanite, apatite, and zircon.

It seems that veins were formed by carbo-fracturing followed by metasomatism post-dating the charnockite magmatism. The nature of the fluid was possibly slightly saline, but carbonic. From our own data and that in the existing literature, we also review the economic potential of these veins.

Keywords: Carbo-fracturing, Syenite veining, Trace element mobility, Metasomatism

Amphibole-bearing granitic rocks in the Ereendavaa block, NE Mongolia: Insights into multi-stage magmatic sources and crystallization conditions

*Munkhdelger Bold¹, Tatsuki Tsujimori¹, Yasuhito Osanai², Nobuhiko Nakano², Tatsuro Adachi², Otgonbayar Dandar¹, Fransiska Ayuni Catur Wahyuandari²

1. Tohoku Univ., 2. Kyushu Univ

The Ereendavaa block (EDB) is located along the southern part of the Mongol-Okhotsk Belt within the eastern Central Asian Orogenic Belt. The EDB has experienced multi-stage magmatism through tectonic cycles, including northward subduction of the Paleo-Asian Ocean during the early Paleozoic and southward subduction of the Mongol-Okhotsk Ocean during the early Mesozoic. However, melt evolution and magma crystallization conditions during magmatic pulses at 540, 470, and 220 Ma are poorly constrained. This study aims to elucidate magmatic crystallization conditions through mineral chemistry (EPMA) analysis of amphibole-bearing granitic rocks from these periods. The granitic rocks in the EDB typically comprise quartz, feldspar, amphibole, and mica, with accessory minerals like opaque. Compositional data for feldspar and hydrous mafic minerals (amphibole and biotite) reveal the following compositions: (1) **~540 Ma granodiorite**: orthoclase, oligoclase, pargasite, and Mg-rich biotite; (2) **~470 Ma granite**: orthoclase, oligoclase, ferro-pargasite, and Fe-rich biotite; (3) **~220 Ma granodiorite**: orthoclase, andesine-oligoclase, edenite, and Mg-rich biotite. The mineral chemistry indicates a sub-alkaline nature for the 540 and 220 Ma granodiorites, while the 470 Ma granite is alkaline. Temperature estimates, based on Ti-in-Ca amphibole geothermometer, range from 957–869°C (540 Ma) and 918–858°C (470 Ma), to 825–782°C (220 Ma). Based on Al-in-Ca amphibole geobarometer, pressure estimates suggest these rocks formed at ~8–6 kbar, ~9–7 kbar, and ~4–3 kbar, respectively. Average oxygen fugacity values are -14.11 (470 Ma granite), -11.82 (540 Ma granodiorite), and -13.04 (220 Ma granodiorite). In summary, the parental magmas for the 540 Ma and 220 Ma granodiorites evolved from orogenic, calc-alkaline to sub-alkaline magmas, emplaced at middle (~26 km) and upper (~12.5 km) crustal levels. They crystallized at high (up to 957°C) and lower temperatures (up to 825°C) under oxidizing conditions. The 470 Ma granitoid likely evolved from more alkaline magmas, emplaced at deep crustal levels (~29 km) approaching the lower crust, and crystallized at high temperatures (up to 918°C) under more reducing conditions than the 540 Ma and 220 Ma granodiorites.

Keywords: CAOB, geothermobarometer, EPMA

Poster presentation | S2: Water Rock Interaction (Special Session)

📅 Fri. Sep 13, 2024 12:30 PM - 2:00 PM JST | Fri. Sep 13, 2024 3:30 AM - 5:00 AM UTC | 🏢 Entrance Hall Higashiyama Campus

S2: Water Rock Interaction (Special Session)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S2-P-01] Scales of extensional-shear fracturing and magnitudes of seismicity induced by magma intrusions into lower crust: Scale comparisons of dike swarm in the high-grade metamorphic rocks and deep low-frequency earthquakes

「発表賞エントリー」

*Takumi Nara¹, Masaoki Uno¹, Tetsuo Kawakami², Fumiko Higashino², Tatsuro Adachi³, Noriyoshi Tsuchiya^{1,4} (1. TOHOKU Univ. Env., 2. Kyoto Univ. Sci., 3. Kyushu Univ. Soc., 4. Hachinohe Kosen)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S2-P-02] Carbonation of serpentinite and formation process of listvenite from Urayama River, Shikokuchuo City, Ehime Prefecture, Japan

「発表賞エントリー」

*Hikaru Takagaki¹, Yohei Shirose¹ (1. Ehime Univ. Sci.&Egn.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S2-P-03] Texture transition during serpentinization in Hodono, Ehime Prefecture

「発表賞エントリー」

*Hinano Wada¹, Enju Satomi¹ (1. Ehime Univ. S/E)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S2-P-04] Water-rock interaction recorded in episyenites from Hakatajima Island, Ehime Prefecture

「発表賞エントリー」

*Toko FUKUI¹, Kazuya SHIMOOKA², Toshiro TAKAHASHI³, Satoshi SAITO¹ (1. Ehime Univ., 2. Kwansei Gakuin Univ., 3. Niigata Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[S2-P-05] Mechanisms of Reaction-Induced Fracturing in Serpentinite Carbonation; Insights from Hydrothermal Experiments and Geochemical Modeling

「発表賞エントリー」

*Taiki Taiki¹, Masaoki Uno¹, Atsushi Okamoto¹ (1. Tohoku University)

Scales of extensional-shear fracturing and magnitudes of seismicity induced by magma intrusions into lower crust: Scale comparisons of dike swarm in the high-grade metamorphic rocks and deep low-frequency earthquakes

*Takumi Nara¹, Masaoki Uno¹, Tetsuo Kawakami², Fumiko Higashino², Tatsuro Adachi³, Noriyoshi Tsuchiya^{1,4}

1. TOHOKU Univ. Env., 2. Kyoto Univ. Sci, 3. Kyushu Univ. Soc., 4. Hachinohe Kosen

Keywords: volcanic earthquakes, magma intrusion, Antarctica

Carbonation of serpentinite and formation process of listvenite from Urayama River, Shikokuchuo City, Ehime Prefecture, Japan

*Hikaru Takagaki¹, Yohei Shirose¹

1. Ehime Univ. Sci.&Egn.

Keywords: listvenite, antigorite, carbonation, serpentinite, Urayama River




Figure. Carbonation and listvenite formation from serpentinite.

BSE images of (a) serpentinite, (b) carbonated serpentinite, (c) carbonated serpentinite in contact with listvenite, (d) listvenite.
Atg : antigorite, Qz : quartz, Mgs : magnesite, Dol : dolomite.

Texture transition during serpentinization in Hodono, Ehime Prefecture

*Hinano Wada¹, Enju Satomi¹

1. Ehime Univ. S/E

Keywords: peridotite, serpentinite, antigorite, Higashi-akaishi peridotite body




図1 : 試料中に含まれる蛇紋石とかんらん石の共存組織 (左 オープンニコル, 右 クロスニコル), (a)直線的な輪郭を持つくさび状-板状のかんらん石と短冊状アンチゴライト, (b)曲線的な輪郭を持つ粒状のかんらん石と短冊状アンチゴライト, (c)メッシュ組織のコアとして存在するかんらん石とリザーダイトリム

Ol : かんらん石, Atg : アンチゴライト, Lz : リザーダイト, Mag : 磁鉄鉱

Water-rock interaction recorded in episyenites from Hakatajima Island, Ehime Prefecture

*Toko FUKUI¹, Kazuya SHIMOOKA², Toshiro TAKAHASHI³, Satoshi SAITO¹

1. Ehime Univ., 2. Kwansei Gakuin Univ., 3. Niigata Univ.

Keywords: Episyenite, metasomatism, Hakatajima Island

Mechanisms of Reaction-Induced Fracturing in Serpentine Carbonation; Insights from Hydrothermal Experiments and Geochemical Modeling

*Taiki Taiki¹, Masaoki Uno¹, Atsushi Okamoto¹

1. Tohoku University

Keywords: Carbonation, Carbon Mineralization, Serpentine, Reaction-induced fracturing

Poster presentation | R4: Mineral sciences of the Earth surface

📅 Fri. Sep 13, 2024 12:30 PM - 2:00 PM JST | Fri. Sep 13, 2024 3:30 AM - 5:00 AM UTC | 🏢 Entrance Hall Higashiyama Campus

R4: Mineral sciences of the Earth surface

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R4-P-01] Design and synthesis of aragonite particles as a reinforcement of plastic materials

*Hiroshi SAKUMA¹, Shigeru SUEHARA¹, Masumi KAMON¹, Kenji TAMURA¹ (1. NIMS)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R4-P-03] Experiments on the Inhibitory Effect of Polysaccharides on Cation Ordering of Dolomite During Dolomitization Reaction at 200°C: Preliminary Results

Hiromi KONISHI¹, *Yao Chen¹ (1. Niigata Univ. Sci.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R4-P-04] Formation Process of Carbonate Minerals in Non-aqueous Solvents: Consideration of the Effects of Different Hydration States of Cations

Naoki IWANE¹, *Jun KAWANO¹, Hiroyuki KAGI², Ayako SHINOZAKI¹, Takaya NAGAI¹ (1. Hokkaido Univ. Sci., 2. UTokyo Sci.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R4-P-05] Impact of evaporation on CO₂ mineralization during enhanced rock weathering

*Naoki NISHIYAMA¹, Masao SORAI¹, Keisuke FUKUSHI², Yuto NISHIKI¹ (1. National Institute of Advanced Industrial Science and Technology (AIST), 2. Kanazawa University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R4-P-06] Framboidal pyrite in modern stromatolite from Fukiage-Jigoku, Onikobe Spring, Miyagi, Japan

「発表賞エントリー」

*Tatsuya Kamada¹, Hiroaki Ohfuji¹ (1. Tohoku Univ. Sci.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R4-P-07] Observations of nano-texture for amosite asbestos by using high-resolution STEM imaging

*Hayato Miura¹, Ichiro Ohnishi¹ (1. JEOL Ltd.)

Design and synthesis of aragonite particles as a reinforcement of plastic materials

*Hiroshi SAKUMA¹, Shigeru SUEHARA¹, Masumi KAMON¹, Kenji TAMURA¹

1. NIMS

Keywords: Calcium carbonate, Elastic modulus, Aspect ratio, Carbon dioxide

Experiments on the Inhibitory Effect of Polysaccharides on Cation Ordering of Dolomite During Dolomitization Reaction at 200°C: Preliminary Results

Hiromi KONISHI¹, *Yao Chen¹

1. Niigata Univ. Sci.

Dolomite is a mineral that consists of calcium-magnesium carbonate, with the chemical formula $\text{CaMg}(\text{CO}_3)_2$. The "dolomite problem" pertains to the challenge of understanding the significant variability in dolomite production over geological time periods and the difficulty of replicating dolomite formation under lab conditions, even though it has been found in surface environments in the past (e.g., Warren 2000). Recent research has shown that polysaccharides such as carboxymethyl cellulose (CMC), agar, and biomass can promote the formation of disordered dolomite at room temperature (e.g., Zhang et al., 2015; 2021). However, Wei and Konishi (submitted) discovered that CMC and agar inhibit the dolomitization reaction at 200°C, although the impact on cation ordering was not clear.

In our study, we are examining the impact of CMC on cation ordering during the dolomitization reaction at 200°C. The experiment involved heating two sets of solutions for varying durations. Both sets had a fixed concentration of Mg and Ca cations at 0.5M and a carbonate ion concentration of 0.1M. One set included 0.2g/L CMC, while the other set did not for comparison. The pH of the solutions was adjusted to 8 before heating. We measured the change in the ratio of 015 peak intensity to 110 peak intensity with different heating durations, which serves as an ordering index indicating the Ca and Mg ordering state in the dolomite structure. In 13 durations of the experiment, 6 showed significantly lower ordering index values with CMC, 6 had similar values, and 1 was higher without CMC. Overall, the trend indicated that the presence of CMC resulted in a lower ordering index in dolomite, suggesting an inhibitory effect on cation ordering.

Keywords: dolomite, Polysaccharides, dolomitization, Inhibitory Effect

Formation Process of Carbonate Minerals in Non-aqueous Solvents: Consideration of the Effects of Different Hydration States of Cations

Naoki IWANE¹, *Jun KAWANO¹, Hiroyuki KAGI², Ayako SHINOZAKI¹, Takaya NAGAI¹

1. Hokkaido Univ. Sci., 2. UTokyo Sci.

Keywords: calcium carbonate, hydration, polymorph, amorphous

Impact of evaporation on CO₂ mineralization during enhanced rock weathering

*Naoki NISHIYAMA¹, Masao SORAI¹, Keisuke FUKUSHI², Yuto NISHIKI¹

1. National Institute of Advanced Industrial Science and Technology (AIST), 2. Kanazawa University

Keywords: Enhanced rock weathering, CO₂ mineralization, Evaporation, Mafic rock, Dissolution

Framboidal pyrite in modern stromatolite from Fukiage-Jigoku, Onikobe Spring, Miyagi, Japan

*Tatsuya Kamada¹, Hiroaki Ohfuji¹

1. Tohoku Univ. Sci.

Keywords: Framboidal pyrite, stromatolite




Figure. 1 Framboidal pyrite enclosed in amorphous silica (opal) in a modern stromatolite from Fukiage-onsen as indicated by the arrow

Observations of nano-texture for amosite asbestos by using high-resolution STEM imaging

*Hayato Miura¹, Ichiro Ohnishi¹

1. JEOL Ltd.

Keywords: Asbestos, TEM, STEM




図 1 アモサイト石綿中に存在する層状ケイ酸塩の高分解能 ADF-STEM 像

Poster presentation | R7: Petrology, Mineralogy and Economic geology (Joint Session with Society of Resource Geology)

📅 Fri. Sep 13, 2024 12:30 PM - 2:00 PM JST | Fri. Sep 13, 2024 3:30 AM - 5:00 AM UTC | 🏢 Entrance Hall Higashiyama Campus

R7: Petrology, Mineralogy and Economic geology (Joint Session with Society of Resource Geology)

岩石学，鉱物学，鉱床学，地球化学などの分野をはじめとして，地球・惑星物質科学全般にわたる岩石及び鉱物に関する研究発表を広く募集する。地球構成物質についての多様な研究成果の発表の場となることを期待する。

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R7-P-01] Ezochiite and placer deposit of platinum group minerals in northwestern Hokkaido, Japan

*Daisuke HAMANE¹, Katsuyuki Saito (1. The University of Tokyo)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R7-P-02] Review for Mineralogical Science: Mineral Resources, Heritage Stone, and SDGs

*Yuhei Takahashi¹ (1. NUE)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R7-P-03] Formation process of olivine with remarkable parting and apparently oblique extinction in the Iherzolite of Ochiai-Hokudo peridotite complex, Okayama Prefecture, Japan

*Terumi EJIMA¹, Takashima Chihiro², Arai Shoji³ (1. Shinshu University, 2. DAIYA SEIKI Co., Ltd., 3. Kanazawa University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R7-P-04] The effect of Na ion on carbonation reaction of forsterite

*Hiroki Hasegawa¹, Atsushi Kyono², Satoru Okada¹, Kosuke Yamaguchi¹ (1. Univ of Tsukuba, Grad. sch. of Life and Environmental. Sci, 2. Univ of Tsukuba, Life and Environmental Science)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R7-P-05] Mineralogical characteristics of Pothole Reef and Pseudo Merensky Reef at the western limb of the Bushveld Complex, South Africa

「発表賞エントリー」

*Amu Umesato¹, Takuya Echigo¹, Yasushi Watanabe¹ (1. Akita Univ. Int.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R7-P-06] High-Ti biotite in the alkali volcanic rock from the Akiyoshi Belt and its significance

*Kosuke Kimura¹, Kaushik Das², Yasutaka Hayasaka³ (1. Osaka Metro. Univ. Sci., 2. Hiroshima Univ., 3. Amakusa Mus. Goshoura Dinosaur Isl.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R7-P-07] Estimation of the origin of SDW in the Horoman peridotite complex by analysis of micro-inclusions in the olivine

「発表賞エントリー」

*Masaharu Aketa¹, Akira Miyake¹, Norikatsu Akizawa², Megumi Matsumoto³, Yohei Igami¹, Itaru Mitsukawa¹ (1. Kyoto University, 2. University of Tokyo, 3. Tohoku University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R7-P-08] Fe-Ni-S-As minerals in the Imono peridotite body, Besshi area, Niihama city, Ehime prefecture.

「発表賞エントリー」

*Masato Kuniyoshi¹, Satomi Enju¹ (1. Ehime Univ.Sci and Eng.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R7-P-09] Fluorite mineralization associated with alkaline metasomatism in the Jinmu-Mihara deposit, Hiroshima, Japan.

「発表賞エントリー」

*Masahiro SUNADA¹, Takuya Echigo¹, Yasushi Watanabe¹ (1. Akita Univ. IRS.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R7-P-10] Petrological study of ultramafic rocks from the Kiyama area, eastern Kumamoto City Narumichi Nishio¹, *Satoko ISHIMARU² (1. Kumamoto Univ. Sci., 2. Kumamoto Univ. FAST)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R7-P-11] FLUID INCLUSION STUDIES IN QUARTZ VEINS WITH TIN MINERALIZATION IN THE KIBARAN INTRUSIVE ROCKS IN KALEHE (SOUTH KIVU, DR CONGO)

「発表賞エントリー」

*MUSA Moise-Kam's SAIDI¹, MADDHUSOODHAN Satish Kumar¹ (1. Niigata Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R7-P-12] Depleted peridotite and melt reaction as recorded by layered dunite-harzburgite in the Horoman peridotite, Hidaka Metamorphic Belt, Hokkaido, Japan.

「発表賞エントリー」

*Keisuke Kurihara¹, Tatsuhiko Kawamoto¹, Aya Hihara¹, Miki Tasaka¹, Hajime Taniuchi², Takeshi Kuritani³, Akiko Matsumoto³ (1. Shizuoka Univ., 2. AIST, 3. Hokkaido Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R7-P-13] Research and development for the exploration of unknown cosmic ray events using Paleo-Detector

*Yuki Ido¹, Tatsuhiro Naka², Shota Futamura³, Tohma Ori⁴, Takenori Kato⁵ (1. Nagoya Univ. Env., 2. Toho Univ. Sci, 3. Nagoya Univ. Sci., 4. N.I.T. Suzuka, 5. Nagoya Univ. ISEE)

Ezochiite and placer deposit of platinum group minerals in northwestern Hokkaido, Japan

*Daisuke HAMANE¹, Katsuyuki Saito

1. The University of Tokyo

Placer deposits of platinum-group minerals (PGM) were once distributed along the central axis of Hokkaido, but due to overhunting during the war, it has become difficult to collect. On the other hand, the several PGM placer deposits have been discovered in northwestern Hokkaido. In this study, we report on the PGM placer deposits in northwestern Hokkaido and the new thiospinel group mineral ezochiite (IMA2022-101) discovered there. Most of placers consist of rutheniridosmine, osmium, iridium, and osmium, while small amount of isoferroplatinum is included. Isoferroplatinum-based grains contain a variety of inclusions, including ezochiite, which is included as a spherical aggregate with other sulfide minerals, suggesting that ezochiite crystallized from melt trapped by isoferroplatinum. The empirical formula of ezochiite is $(\text{Cu}^{+}_{0.85}\text{Fe}^{3+}_{0.15})(\text{Rh}^{3+}_{1.09}\text{Pt}^{4+}_{0.78}\text{Ir}^{3+}_{0.08}\text{Pt}^{2+}_{0.05})\text{S}_{4.00}$, ideally $\text{Cu}^{+}(\text{Rh}^{3+}\text{Pt}^{4+})\text{S}_4$. The unit cell parameter is $a = 9.8559(14) \text{ \AA}$ on Fd-3m space group. Ezochiite was confirmed in ophiolites, Ural-Alaskan intrusions, and orthomagmatic deposits, suggesting that ezochiite is a common mineral in PGM deposit.

Keywords: Platinum-group minerals (PGM), Placer PGM, Ezochiite




図1. 模式標本の蝦夷地鉱 (苫前町海岸) (a)、初山別川試料の蝦夷地鉱 (b)、これまでに確認された白金族鉱物と主な産状 (c)、北海道産試料の硫銅ロジウム鉱—蝦夷地鉱の組成分布と鉱物分類 (d)。産地：茂初山別川 (M)、初山別川 (S)、アイヌ沢川 (A)、苫前町海岸 (TC)、小平町海岸 (OC)、上記念別沢川 (K)、小平薬川 (O)、沼田ポン川 (N)、雨竜川 (U)

Review for Mineralogical Science: Mineral Resources, Heritage Stone, and SDGs

*Yuhei Takahashi¹

1. NUE

Keywords: Mineral Resources, Heritage Stone

Formation process of olivine with remarkable parting and apparently oblique extinction in the Iherzolite of Ochiai-Hokudo peridotite complex, Okayama Prefecture, Japan

*Terumi EJIMA¹, Takashima Chihiro², Arai Shoji³

1. Shinshu University, 2. DAIYA SEIKI Co., Ltd., 3. Kanazawa University

Keywords: Ochiai-Hokudo, peridotite complex, olivine

The effect of Na ion on carbonation reaction of forsterite

*Hiroki Hasegawa¹, Atsushi Kyono², Satoru Okada¹, Kosuke Yamaguchi¹

1. Univ of Tsukuba, Grad. sch. of Life and Environmental. Sci, 2. Univ of Tsukuba, Life and Environmental Science

Keywords: Forsterite, Magnesite, Carbonation, Geological Carbon Storage

Mineralogical characteristics of Pothole Reef and Pseudo Merensky Reef at the western limb of the Bushveld Complex, South Africa

*Amu Umesato¹, Takuya Echigo¹, Yasushi Watanabe¹

1. Akita Univ. Int.

Keywords: Bushveld Complex, Platinum Group Elements, Merensky Reef, Monosulfide Solid Solution, Intermediate Solid Solution




Fig. 1. Moncheite (white) formed on the rim of Nickel sulfide minerals (gray) in Pothole Reef. BSE image.




Fig. 2. Zvyagintsevite (white) formed on the rim of pentlandite (gray), in Pseudo Merensky Reef. BSE image.

High-Ti biotite in the alkali volcanic rock from the Akiyoshi Belt and its significance

*Kosuke Kimura¹, Kaushik Das², Yasutaka Hayasaka³

1. Osaka Metro. Univ. Sci., 2. Hiroshima Univ., 3. Amakusa Mus. Goshoura Dinosaur Isl.

Keywords: Akiyoshi Belt, Greenstone, Zircon U-Pb age, High titanium biotite

Estimation of the origin of SDW in the Horoman peridotite complex by analysis of micro-inclusions in the olivine

*Masaharu Aketa¹, Akira Miyake¹, Norikatsu Akizawa², Megumi Matsumoto³, Yohei Igami¹, Itaru Mitsukawa¹

1. Kyoto University, 2. University of Tokyo, 3. Tohoku University


Keywords: Horoman peridotites, olivine, inclusion, Electron Microscope, XnCT

Fe-Ni-S-As minerals in the Imono peridotite body, Besshi area, Niihama city, Ehime prefecture.

*Masato Kuniyoshi¹, Satomi Enju¹

1. Ehime Univ.Sci and Eng.

Keywords: Fe-Ni-S-As mineral, Serpentinite, Peridotite, Imono peridotite body



図：蛇紋岩中の磁鉄鉱，ペントランド鉱，アワルワ鉱，クロンステッドタイト様鉱物の共生組織

Fluorite mineralization associated with alkaline metasomatism in the Jinmu-Mihara deposit, Hiroshima, Japan.

*Masahiro SUNADA¹, Takuya Echigo¹, Yasushi Watanabe¹

1. Akita Univ. IRS.

Keywords: Fluorite, Alkali (Na-K) metasomatism, Jinmu-Mihara deposit

Petrological study of ultramafic rocks from the Kiyama area, eastern Kumamoto City

Narumichi Nishio¹, *Satoko ISHIMARU²

1. Kumamoto Univ. Sci., 2. Kumamoto Univ. FAST

Keywords: Serpentinite, Dunite, Chromian spinel, Kiyama metamorphic rock

FLUID INCLUSION STUDIES IN QUARTZ VEINS WITH TIN MINERALIZATION IN THE KIBARAN INTRUSIVE ROCKS IN KALEHE (SOUTH KIVU, DR CONGO)

*MUSA Moise-Kam's SAIDI¹, MADDHUSOODHAN Satish Kumar¹

1. Niigata Univ.

The Mesoproterozoic Kibaran granitoid formations host numerous mineral resources which are highly demanded in new technology industries, mineral such as Sn-W, Nb, Ta and Au. This mobile belt is formed by collision of the Western Congo with the Eastern Tanzanian, Bangwelu and Zimbabwe cratons, producing four generations of granite. These ore deposits are hosted in granite pegmatite (Nb, Ta) and Sn-W in quartz veins.

Fluid inclusions were studied in quartz vein samples in order to unravel the evolution of the mineralizing fluid controlling the ore deposits in Kalehe(DR Congo). In the study area the field and petrographic studies indicate that major rocks constitute of a variety of metapelites including sericite schists, micas schists, quartzites and igneous rocks such as granites, greisens and pegmatites. The quartz veins crosscutting the metapelite show a NE-SW trending, whereas the pegmatitic vein trend in a NW-SE direction crosscutting the Kibaran metasediments.

Fluid inclusions are abundant in quartz veins and metallic mineralized quartz veins. Primary, secondary and pseudo secondary inclusion are observed, in which the biphasic inclusions are most common. The mineralizing fluids of the quartz-vein in the sericite schist have a heterogeneous nature at the time of trapping. The temperature of homogenization(T_h) and salinities of the fluid inclusions were determined; some were having high T_h with high salinity, others were with high T_h but with low salinity, whereas another group have low T_h with moderate salinity. The biphasic Inclusions homogenization temperature range is 17.7- 303°C and the salinity range is 0.88-16.24°Wt%. The temperature of first ice melting(T_{mice})values obtained so far from fluid inclusion microthermometry indicate various fluid compositions mainly CO₂, CO₂ + H₂O rich, CO₂+H₂O+/-CH₄. These phases were confirmed by Raman spectroscopy and consistent with T_{mice} as well as the freezing point depression indicating the presence of salt (NaCl or other salts).

The mineralizing fluids trapped as fluid inclusions in the metallic quartz veins crosscutting shale formations can also be categorized as primary, secondary and pseudo secondary, these fluids were multiphase fluid inclusion (solid, liquid and vapor). Detailed studies on these inclusions are being carried out and will be presented.

Keywords: Fluid inclusion, microthermotry, Kibaran mobile belt, Tin, Raman spectroscopy

Depleted peridotite and melt reaction as recorded by layered dunite-harzburgite in the Horoman peridotite, Hidaka Metamorphic Belt, Hokkaido, Japan.

*Keisuke Kurihara¹, Tatsuhiko Kawamoto¹, Aya Hihara¹, Miki Tasaka¹, Hajime Taniuchi², Takeshi Kuritani³, Akiko Matsumoto³

1. Shizuoka Univ., 2. AIST, 3. Hokkaido Univ.

Keywords: peridotite, Horoman, banded dunite-harzburgite, melt-rock reaction, chemical composition

Research and development for the exploration of unknown cosmic ray events using Paleo-Detector

*Yuki Ido¹, Tatsuhiro Naka², Shota Futamura³, Tohma Ori⁴, Takenori Kato⁵

1. Nagoya Univ. Env., 2. Toho Univ. Sci, 3. Nagoya Univ. Sci., 4. N.I.T. Suzuka, 5. Nagoya Univ. ISEE

Keywords: Muscovite, Olivine, Particle physics

Poster presentation | R8: Metamorphic rocks and tectonics

📅 Fri. Sep 13, 2024 12:30 PM - 2:00 PM JST | Fri. Sep 13, 2024 3:30 AM - 5:00 AM UTC | 🏢 Entrance Hall Higashiyama Campus

R8: Metamorphic rocks and tectonics

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R8-P-01] Mineralogical characterization of serpentinite varieties in Sangun-Renge Belt exposed at Sasaguri, Fukuoka Prefecture, and their geological implications

「発表賞エントリー」

*Swarna ANNADURAI MUNUSAMY¹, Jun-ichi ANDO^{1,2}, Yuki IWASAKI³, Kaushik DAS^{1,2}, Dyuti Prakash SARKAR⁴, Seiichiro UEHARA⁵ (1. Hiroshima Univ., 2. HiPeR, Hiroshima, 3. NIPPON STEEL CORP., 4. JAMSTEC, 5. The Kyushu Univ. Museum)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R8-P-02] Metamorphic temperature structure of Sanbagawa Metamorphic Belt in the southern part of Shinshiro City, Aichi Prefecture, Japan

「発表賞エントリー」

*Akane Matsuzaki¹, Yui Kouketsu¹, Katsuyoshi Michibayashi¹ (1. Nagoya Univ. Env.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R8-P-03] Origin and pyrometamorphism of gneissose granitoid xenoliths from Mt. Daisen, Tottori Prefecture, SW Japan

「発表賞エントリー」

*Mizuki TAKAHASHI¹, Shunsuke Endo¹ (1. Shimane University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R8-P-04] Petrography of monazite in a metapelite in the eastern Nepal Himalaya and Development of Th-Pb dating method for monazite

「発表賞エントリー」

*Shumpei KUDO¹, Tetsuo Kawakami¹, Sota Niki², Toru Nakajima³, Takafumi Hirata⁴, Takeshi Imayama⁵ (1. Kyoto Univ. Sci., 2. Nagoya Univ. ISEE., 3. JAEA, 4. UTokyo. Sci., 5. Okayama Univ. of Sci.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R8-P-05] Petrological description of ultramafic rocks in the low-grade metamorphic zone of the Sanbagawa belt: A case study of the Ina area, Nagano Prefecture, central Japan

「発表賞エントリー」

*Kaho Nobuhara¹, Hiroshi Mori¹, Takayoshi Nagaya² (1. Shinshu Univ., 2. Tokyo Gakugei Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R8-P-06] Detrital zircon U-Pb dating and Raman spectral analysis of carbonaceous material in the boundary area of the Sanbagawa-Chichibu belts, central Kii Peninsula

*Hiroshi MORI¹, Kojiro USUI^{1,2}, Tetsuya Tokiwa¹, Kazuhiro Ozawa³ (1. Shinshu University, 2. Nippon Koei Co., Ltd., 3. Precision Forestry Measurement Ltd.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R8-P-07] New finding of wakefieldite from an amphibolite in the Horokanai area, Kamuikotan HP metamorphic belt, Hokkaido, Japan

*Taro Kato¹, Kosuke NAEMURA¹, Toru Takeshita² (1. Iwate University, 2. Pacific Consultants Co., Ltd.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R8-P-08] Thermal history and protolithic detritus provenance of a sillimanite-chrysoberyl-bearing gneiss from the Ashio mountains in the western part of Tochigi prefecture

*Ippei KITANO¹ (1. Hokkaido Univ. Mus.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R8-P-09] Petrological characterization and geochronology of metamorphic rocks from the Northern Subzone of the Maizuru Terrane

*Sota Muroi¹, Kaushik Das¹, Kenta Kawaguchi¹, Yasutaka Hayasaka² (1. Hiroshima University, 2. Amakusa Museum of Goshoura Dinosaur Island)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R8-P-10] The fate of organic carbon during subduction: Raman micro-spectroscopy and C-isotope geochemistry of carbonaceous materials in Sambagawa pelitic schists, central Shikoku, Japan

*Hironobu Harada¹, Tatsuki Tsujimori¹, Akizumi Ishida¹, Takeshi Kakegawa¹, Tetsumaru Itaya² (1. Tohoku Univ., 2. jGnet)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R8-P-11] Fluid inclusions of ophicarbonates in the Apennine Mountains, Italy

HiroYuki Kaneko¹, *Tatsuhiko KAWAMOTO¹, Francesca Meneghini², Yosuke Osawa¹ (1. Shizuoka University, 2. Università di Pisa | UNIPI · Department of Earth Sciences)

Mineralogical characterization of serpentinite varieties in Sangun-Renge Belt exposed at Sasaguri, Fukuoka Prefecture, and their geological implications

*Swarnaa ANNADURAI MUNUSAMY¹, Jun-ichi ANDO^{1,2}, Yuki IWASAKI³, Kaushik DAS^{1,2}, Dyuti Prakash SARKAR⁴, Seiichiro UEHARA⁵

1. Hiroshima Univ., 2. HiPeR, Hiroshima, 3. NIPPON STEEL CORP., 4. JAMSTEC, 5. The Kyushu Univ. Museum

Serpentinite (serp) of Sangun-Renge Belt are widely exposed around the Narubuchi dam in Sasaguri area, Fukuoka Prefecture. The purpose of this study is to investigate the mineralogical characteristics of different types of serp in this region in order to understand the tectonic and metamorphic history of the Sangun-Renge Belt serp. Optical microscopy reveals the petrographic characteristics, EPMA is used to measure the composition of Cr-spinel, and Raman spectroscopy is utilized to distinguish serpentine polymorphs. The serp in the Sasaguri area primarily consists of serpentine group minerals, with minor amounts of magnetite, calcite, chlorite, talc, and Cr-spinel. Three types are identified on the basis of their dominant serpentine polymorph. Type 1: Lizardite serp, showing the hourglass textures with chrysotile veins. Type 2: Lizardite-Chrysotile serp, dominated by a mesh texture where the core and rim minerals are lizardite and chrysotile, and lizardite, respectively. And the antigorite (atg) veins are observed. Type 3: Atg serp, exposed to the north and south of the study area. Their microstructures are different. The serp in the north is composed of atg a few mm in grain size, showing undulose extinction and dynamic recrystallization, but no CPO. The serp in the south is composed of atg less than 200 μm in size, with atg veins. At the north, the mylonitized zones can be identified where the atg grain size is less than 100 μm , with (001) and [010] oriented parallel to foliation and lineation, respectively. The chemical composition of the Cr-spinel in the north serp indicates a forearc peridotite origin. The ferritchromite rim in the Cr-spinel suggests that the peridotite is re-equilibrated at greenschist to amphibolite temperatures (400-700°C). A magnetite overgrowth around the ferritchromite rim indicates the serpentinization after ferritchromitization, which occurs at 250-400°C. Based on the above results, the geological implications of the identified serpentine polymorphs and associated mineral assemblages for understanding the tectonic and metamorphic history of the Sangun-Renge Belt serp will be discussed in our presentation.

Keywords: Serpentinization, Antigorite, Serpentine polymorphs, Cr-spinel

Metamorphic temperature structure of Sanbagawa Metamorphic Belt in the southern part of Shinshiro City, Aichi Prefecture, Japan

*Akane Matsuzaki¹, Yui Kouketsu¹, Katsuyoshi Michibayashi¹

1. Nagoya Univ. Env.

Keywords: Quartz, Raman carbonaceous material geothermometer, Sanbagawa Metamorphic Belt

Origin and pyrometamorphism of gneissose granitoid xenoliths from Mt. Daisen, Tottori Prefecture, SW Japan

*Mizuki TAKAHASHI¹, Shunsuke Endo¹

1. Shimane University

Keywords: Mt. Daisen, xenolith, pyrometamorphism

Petrography of monazite in a metapelite in the eastern Nepal Himalaya and Development of Th-Pb dating method for monazite

*Shumpei KUDO¹, Tetsuo Kawakami¹, Sota Niki², Toru Nakajima³, Takafumi Hirata⁴, Takeshi Imayama⁵

1. Kyoto Univ. Sci., 2. Nagoya Univ. ISEE., 3. JAEA, 4. UTokyo. Sci., 5. Okayama Univ. of Sci.

Keywords: Monazite, Th-Pb dating method, Higher Himalayan Crystallines

Petrological description of ultramafic rocks in the low-grade metamorphic zone of the Sanbagawa belt: A case study of the Ina area, Nagano Prefecture, central Japan

*Kaho Nobuhara¹, Hiroshi Mori¹, Takayoshi Nagaya²

1. Shinshu Univ. , 2. Tokyo Gakugei Univ.

Keywords: Sanbagawa (Sambagawa) belt, ultramafic rock, chlorite zone, Ina area

Detrital zircon U–Pb dating and Raman spectral analysis of carbonaceous material in the boundary area of the Sanbagawa–Chichibu belts, central Kii Peninsula

*Hiroshi MORI¹, Kojiro USUI^{1,2}, Tetsuya Tokiwa¹, Kazuhiro Ozawa³

1. Shinshu University, 2. Nippon Koei Co., Ltd., 3. Precision Forestry Measurement Ltd.

Keywords: Sanbagawa Belt, Chichibu Belt, detrital zircon U–Pb dating, Raman spectral analysis, carbonaceous material

New finding of wakefieldite from an amphibolite in the Horokanai area, Kamuikotan HP metamorphic belt, Hokkaido, Japan

*Taro Kato¹, Kosuke NAEMURA¹, Toru Takeshita²

1. Iwate University, 2. Pacific Consultants Co., Ltd.

Kamuikotan amphibolite exposed at the Horokanai area experienced transformation from amphibolite to blueschist facies metamorphism. Similar occurrences are found throughout the world, and it is generally regarded as a result of an isobaric cooling (counterclockwise) pathway. In other words, amphibolite is thought to have remained at depth and experienced cooling. To address this issue, we studied an epidote amphibolite exposed at the Horokanai hill. The rock displays a foliation consisting of hornblende/actinolite, epidote, plagioclase, rutile, white mica (Si=3.3-3.5 apfu, O=11) ±chlorite, and these primary minerals are more or less replaced by secondary blueschist facies minerals along pull apart fractures and rims of primary minerals. The latter consists of glaucophane, chlorite, titanite, epidote, pumpellyite, and white mica (Si=3.5-3.8 apfu, O=11). Yttrium vanadate (wakefieldite) grains occur along pull apart fractures of epidote and hornblende, and the remaining fractures were filled by phengite rich white mica. This suggests wakefieldite was formed before or during blueschist facies metamorphism. According to previous studies, wakefieldite could be stable under ultraoxidation conditions at shallow depths. Following this idea, it is unlikely that the Kamuikotan amphibolite was transformed into blueschist by isobaric cooling at depth, but rather once exhumed to shallow depth and sunk back to the depth.

Keywords: Kamuikotan metamorphic belt, yttrium vanadate, wakefieldite-(Y), P-T-t path

Thermal history and protolithic detritus provenance of a
sillimanite–chrysoberyl-bearing gneiss from the Ashio mountains in the
western part of Tochigi prefecture

*Ippei KITANO¹

1. Hokkaido Univ. Mus.

Keywords: sillimanite–chrysoberyl-bearing gneiss, andalusite-bearing granite, thermal history, detritus
provenance, Ashio mountains

Petrological characterization and geochronology of metamorphic rocks from the Northern Subzone of the Maizuru Terrane

*Sota Muroi¹, Kaushik Das¹, Kenta Kawaguchi¹, Yasutaka Hayasaka²

1. Hiroshima University, 2. Amakusa Museum of Goshoura Dinosaur Island

Keywords: Maizuru Terrane, Northern Subzone, Komori-Kuwagai metamorphic rocks, P-T estimates, CHIME age of monazite

The fate of organic carbon during subduction: Raman micro-spectroscopy and C-isotope geochemistry of carbonaceous materials in Sambagawa pelitic schists, central Shikoku, Japan

*Hironobu Harada¹, Tatsuki Tsujimori¹, Akizumi Ishida¹, Takeshi Kakegawa¹, Tetsumaru Itaya²

1. Tohoku Univ., 2. jGnet

Keywords: carbonaceous materials, carbon isotope, Raman micro-spectroscopy, Sambagawa Belt, pelitic schist

Fluid inclusions of ophicarbonates in the Apennine Mountains, Italy

Hiroyuki Kaneko¹, *Tatsuhiko KAWAMOTO¹, Francesca Meneghini², Yosuke Osawa¹

1. Shizuoka University, 2. Università di Pisa | UNIPI ·Department of Earth Sciences



The Bracco ophicarbonate from the Apennines, Italy, is studied for petrography and microthermometry of fluid inclusions in the calcite. The studied serpentinite body is a part of an ophiolite that has undergone relatively low grade metamorphism, considered to be pumpellyite facies, and retains hydrothermal metamorphic vein textures and seafloor depositional textures that may record interactions with Jurassic seawater and mantle peridotite/ serpentinite (Cannao et al., 2020, in Chemical Geology).

Raman spectroscopy and micro-XRF analysis indicate that the rocks are mainly composed of antigorite, lizardite, and calcite. We performed microthermometry of saline fluid inclusions in relatively large crystals of calcite veins. Other fine-grained calcite crystals also exist, but microthermometry of their fluid inclusions has been difficult up to now. We also have sedimentary ophicarbonate (Cannao et al., 2020) in the same outcrop, and we would like to obtain data on these fluid inclusions in the future.

Microthermometry results show that the salinity estimated from freezing point depression is 5.0 ± 1.8 NaCl wt% (n=9) and the homogenization temperature between gas and liquid phases is $179 \pm 16^\circ\text{C}$ (n=9). The salinity varies from 1 to 8 NaCl wt% within a thin section, and even within a crystal there are 3 NaCl wt% variations, indicating that saline fluids with such degrees of variation existed during the carbonation of the serpentinite.

We would like to discuss the carbonation process of serpentinite at the seafloor by comparing the salinity and homogenization temperature of fluid inclusions in calcite in ophiolites from low grade metamorphic regions in the Western Alps reported by Inukai et al. (2023, Mineralogical Society Abstracts) and other previous studies.

Keywords: fluid inclusions, serpentinite, seawater, carbon dioxide, ophicarbonate



JAMS General Meeting, Award ceremony

📅 Fri. Sep 13, 2024 2:00 PM - 3:15 PM JST | Fri. Sep 13, 2024 5:00 AM - 6:15 AM UTC | 🏢 ES Hall
Higashiyama Campus

JAMS General Meeting, Award ceremony

2:00 PM - 3:15 PM JST | 5:00 AM - 6:15 AM UTC

[2Lecture-1-1400-1add] 定時総会・授賞式

JAMS General Meeting, Award ceremony

JAMS General Meeting, Award ceremony

Fri. Sep 13, 2024 2:00 PM - 3:15 PM ES Hall (Higashiyama Campus)

2:00 PM - 3:15 PM

[2Lecture-1-1400-1add]定時総会・授賞式

Lectures of JAMS Awardees

🎵 Fri. Sep 13, 2024 3:30 PM - 5:00 PM JST | Fri. Sep 13, 2024 6:30 AM - 8:00 AM UTC | 🏢 ES Hall
Higashiyama Campus

Lectures of JAMS Awardees

3:30 PM - 5:00 PM JST | 6:30 AM - 8:00 AM UTC

受賞者講演

Oral presentation | R6: Plutonic rocks, volcanic rocks and subduction factory

📅 Sat. Sep 14, 2024 9:00 AM - 12:00 PM JST | Sat. Sep 14, 2024 12:00 AM - 3:00 AM UTC | 🏢 ES Hall Higashiyama Campus

R6: Plutonic rocks, volcanic rocks and subduction factory

Chairperson: Tatsuhiko Kawamoto (Shizuoka University), Takashi Yuguchi, Atsushi Kamei

9:00 AM - 9:15 AM JST | 12:00 AM - 12:15 AM UTC

[R6-01] Compositional evolution of slab-derived fluids during ascent and origin of subduction-zone primary magmas: Implications from trace-element partition between hydrous melts and Cl-free or Cl-rich aqueous fluids

*Hajime TANIUCHI¹, Tatsuhiko KAWAMOTO², Takayuki NAKATANI¹, Osamu ISHIZUKA¹, Toshihiro SUZUKI¹, Akihiko Tomiya¹ (1. GSJ, AIST, 2. Shizuoka Univ.)

9:15 AM - 9:30 AM JST | 12:15 AM - 12:30 AM UTC

[R6-02] New crust and mantle physical properties research from the development of rock synthesis methods

「招待講演」

*Sanae Koizumi¹ (1. The University of Tokyo)

9:30 AM - 9:45 AM JST | 12:30 AM - 12:45 AM UTC

[R6-03] The use of contact metamorphism to identify plutons associated with major contemporaneous eruption and estimate magma flux

*Simon R WALLIS¹, Ken Yamaoka², Akira Miyake³, Catherine Annen⁴ (1. The University of Tokyo, 2. AIST, 3. Aichi University of Education, 4. Institute of Geophysics, Czech Academy of Sciences)

9:45 AM - 10:00 AM JST | 12:45 AM - 1:00 AM UTC

[R6-04] Petrography and geochronology of the Kuki granite, Kitakami Mountains, northeastern Japan: Shallow crustal intrusion and emplacement processes of granitic magma

*Satoshi SUZUKI¹, Takashi YUGUCHI², Keito ISHIGURO¹, Kyoka ENDO¹, Asuka KATO¹, Kosuke YOKOYAMA¹, Yasuhiro OGITA³, Tatsunori YOKOYAMA³, Shuhei SAKATA⁴, Takeshi OHNO⁵, Eiji SASAO³ (1. Yamagata University, 2. Kumamoto University, 3. Japan Atomic Energy Agency, 4. University of Tokyo, 5. Gakushuin University)

10:00 AM - 10:15 AM JST | 1:00 AM - 1:15 AM UTC

[R6-05] Zircon U-Pb dating of granitoids in the Hobenzan area, western Yamaguchi Prefecture, southwest Japan

*Yukiyasu TSUTSUMI¹, Kenichiro TANI¹ (1. National Museum of Nature and Science)

10:15 AM - 10:30 AM JST | 1:15 AM - 1:30 AM UTC

[R6-06] Estimation of temperature and pressure conditions of granitoids in the eastern part of Shimane Prefecture, San'in Belt.

*Kai Nakahashi¹, Satoshi SAITO¹ (1. Ehime University Graduate School of Science and Engineering, Earth Science Division)

10:30 AM - 10:45 AM JST | 1:30 AM - 1:45 AM UTC

[R6-07] Primary magma of Cretaceous Northern Kyushu batholith: New approach from Cumulus rocks

*Keisuke ESHIMA¹, Masaaki OWADA¹, Atsushi KAMEI² (1. Yamaguchi Univ. Sci., 2. Shimane Univ.)

10:45 AM - 11:00 AM JST | 1:45 AM - 2:00 AM UTC

[R6-08] Origin of the Cretaceous ignimbrite flare-up in SW Japan

*Ken Yamaoka¹, Tokiyuki Morohoshi², Daisuke Sato¹, Kazuto Mikuni¹ (1. AIST, 2. Univ. Tokyo. Sci.)

11:00 AM - 11:15 AM JST | 2:00 AM - 2:15 AM UTC

[3Lecture-101-11-9add] 休憩

11:15 AM - 11:30 AM JST | 2:15 AM - 2:30 AM UTC

[R6-09] The genesis of garnet in the Quaternary rhyolite of Hime-shima volcanic group, EEast Kyushu, Southwest Japan Arc.

*Takehiro HIRAYAMA^{1,2}, Tomoyuki SHIBATA², Masako YOSHIKAWA² (1. Tokyo University, 2. Hiroshima University)

11:30 AM - 11:45 AM JST | 2:30 AM - 2:45 AM UTC

[R6-10] K–Ar ages and geochemical compositions of Early Pleistocene andesites from southern Kyushu, Japan

*Raiki YAMADA¹, Masakazu Niwa¹, Koji Umeda², Koshi Yagi³, Taisei Fujiwara³, Ikuo Okada³, Fukuka Kida¹ (1. Japan Atomic Energy Agency, 2. Hirosaki University, 3. Hiruzen Institute for Geology and Geochronology)

11:45 AM - 12:00 PM JST | 2:45 AM - 3:00 AM UTC

[R6-11] Thermal history of pluton formation from Sr diffusion in plagioclase: implications for magma flux estimation and identification of plutons associated with major eruptions

「発表賞エントリー」

*Tokiyuki Morohoshi¹, Simon Richard Wallis¹ (1. The University of Tokyo)

Compositional evolution of slab-derived fluids during ascent and origin of subduction-zone primary magmas: Implications from trace-element partition between hydrous melts and Cl-free or Cl-rich aqueous fluids

*Hajime TANIUCHI¹, Tatsuhiko KAWAMOTO², Takayuki NAKATANI¹, Osamu ISHIZUKA¹,
Toshihiro SUZUKI¹, Akihiko Tomiya¹

1. GSJ, AIST, 2. Shizuoka Univ.

Keywords: Primary magma, Slab-derived fluid, Partition coefficient, High-pressure and high-temperature experiment

New crust and mantle physical properties research from the development of rock synthesis methods

*Sanae Koizumi¹

1. The University of Tokyo

Keywords: Olivine, Mantle, Crust, Sample synthesis

The use of contact metamorphism to identify plutons associated with major contemporaneous eruption and estimate magma flux

*Simon R WALLIS¹, Ken Yamaoka², Akira Miyake³, Catherine Annen⁴

1. The University of Tokyo, 2. AIST, 3. Aichi University of Education, 4. Institute of Geophysics, Czech Academy of Sciences

Magmatic flux is an important factor controlling the heat and material transport in the continental crust; it is also thought to be a major control on the potential for any magma to erupt, and duration of eruption and time scale between major events. Plutons are a clear record of magma input into the earth's crust and volume estimates combined with geochronological studies have been used to estimate past magma flux rates. However, areas suitable for such studies are limited and this approach cannot assess possible loss of magma through associated volcanic eruption. Contact metamorphic aureoles developed around plutons record the thermal effects of magma input into the crust and their study provides a potential way to use the geological record to estimate ancient magma flux rates even if material has been lost from the system by eruption.

A compilation of aureoles surrounding silicic plutons around the world shows that there is a wide variation in normalized thicknesses. Comparison with thermal modelling shows some aureoles are compatible with rapid input of the entire volume of magma recorded in the pluton whereas other aureoles are very narrow and imply slow rates of magma input. There are also numerous examples of aureoles broader than can be explained easily by either instantaneous intrusion models or incremental growth of plutons. Such broad aureoles may be explained by 1) unusually high crustal temperatures just before intrusion and 2) the contribution of magma that has subsequently been lost from the pluton system due to eruption.

The Shinshiro tonalite and the Busetsu granite are two plutons formed in the same Hongusan area and have similar zircon U–Pb ages. Although the Busetsu granite has a larger volume, it is associated with a much narrower aureole than the Shinshiro tonalite. Because the intrusions formed at the same time in the same area, the difference in aureole widths cannot be explained by contrasting background crustal temperatures. Thermal modeling shows that only eruptive loss of magma can adequately account for the characteristics of the Shinshiro tonalite aureole. Combining petrological studies of metamorphic aureoles with thermal modeling can identify plutons that developed in association with major volcanic eruptions and constrain ancient magma flux.

Keywords: Magma flux, Pluton, Contact metamorphism

Petrography and geochronology of the Kuki granite, Kitakami Mountains, northeastern Japan: Shallow crustal intrusion and emplacement processes of granitic magma

*Satoshi SUZUKI¹, Takashi YUGUCHI², Keito ISHIGURO¹, Kyoka ENDO¹, Asuka KATO¹, Kosuke YOKOYAMA¹, Yasuhiro OGITA³, Tatsunori YOKOYAMA³, Shuhei SAKATA⁴, Takeshi OHNO⁵, Eiji SASAO³

1. Yamagata University, 2. Kumamoto University, 3. Japan Atomic Energy Agency, 4. University of Tokyo, 5. Gakushuin University

Keywords: Flare-up, Non-adakitic magma, Fractional crystallization, P-T history, t-T history




図1 久喜花崗岩体のホルンブレンドから導出した温度と圧力（深度）の関係（a）、ジルコンU-Pb年代・Ti濃度・Th/U比の同時測定による t - T 履歴（b）とTh/U比と温度の関係（c）。

Zircon U-Pb dating of granitoids in the Hobenzan area, western Yamaguchi Prefecture, southwest Japan

*Yukiyasu TSUTSUMI¹, Kenichiro TANI¹

1. National Museum of Nature and Science

Zircons U-Pb ages were obtained from granitoids in the Hobenzan area, western Yamaguchi Prefecture, southwest Japan. Granitoid bodies in the area consist of granodiorite and granite. The 5 granodioritic samples indicate 95.4 ± 0.7 Ma, 96.3 ± 1.0 Ma, 96.3 ± 0.8 Ma, 96.2 ± 0.6 Ma and 97.1 ± 0.7 Ma whereas the 2 granitic samples indicate 93.9 ± 0.7 Ma and 90.2 ± 0.8 Ma. Errors are 95% confidence intervals. These ages are equivalent of the granitoids in northern Kyushu, but there is no report for granitoid which age is more than 95 Ma from eastern Chugoku District. Additional research is necessary to verify the chronological continuation of granitoids in SW Japan.

Keywords: SW Japan, San-yo Province, Cretaceous, granitoid, zircon U-Pb age

Estimation of temperature and pressure conditions of granitoids in the eastern part of Shimane Prefecture, San'in Belt.

*Kai Nakahashi¹, Satoshi SAITO¹

1. Ehime University Graduate School of Science and Engineering, Earth Science Division

Keywords: granitoids, Amphibole only geothermobarometer, San-in granitoids

Primary magma of Cretaceous Northern Kyushu batholith: New approach from Cumulus rocks

*Keisuke ESHIMA¹, Masaaki OWADA¹, Atsushi KAMEI²

1. Yamaguchi Univ. Sci., 2. Shimane Univ.

The Jurassic to Cretaceous igneous rocks associated with the subduction of the Izanagi Plate are widely developed in the Northeast Asian continental margin and have been intensively studied for their igneous activity and tectonics (Kim et al., 2016, *Lithos* 262, 88–106; Yu et al. et al., 2021, *Int. Geol. Rev.*, 1881920). It is clear that the igneous rocks in the Japanese Islands, as well as the continental side (Northeastern China and Korean Peninsula), are related to the subduction (200–50 Ma) of the Izanagi Plate into Eurasia (Seton et al., 2015, *Geophys. Res. Lett.* 42, 1732–1740), and approximately 30% of Japan archipelago is occupied by plutons, typically granitoid. In addition, large-scale granitic complexes with an area of more than 100 km², known as batholiths, are exposed throughout the Japanese island and compose the major part of this island arc crust. Furthermore, about 80% of the granitoids exposed in Japan are concentrated in the geochronological age range 50–130 Ma (mainly Cretaceous to Paleogene). These continental-scale periodic large-scale igneous events are called Magmatic flare-ups and are very important for the mechanism of generation of large amounts of silicic magma and the evolution of crustal growth processes (Ducea, 2001, *GSA Today*, 11, 4–10). In this presentation, we discuss a comprehensive study of the Cretaceous Northern Kyushu Batholith, including magma generation mechanisms, especially the formation process and generation mechanisms of source magmas of the granitoids that form the batholith-scaled pluton. In addition, the source magmas will be examined using the Cumulus rocks from the Kita-Taku mafic complex associated with the Cretaceous Northern Kyushu batholith. We believe that the use of the cumulus minerals from the primary magma and the tracing of the characteristics of the primary magma is the best approach in the Northern Kyushu batholith, where there are very few exposures of coeval volcanic rocks (i.e., it is difficult to determine the melt composition from volcanic rocks).

Keywords: Cretaceous Northern Kyushu Batholith, Kita-taku mafic complex, Cumulate, Primary magma, Sanukitic HMA

Origin of the Cretaceous ignimbrite flare-up in SW Japan

*Ken Yamaoka¹, Tokiyuki Morohoshi², Daisuke Sato¹, Kazuto Mikuni¹

1. AIST, 2. Univ. Tokyo. Sci.

Keywords: Cretaceous, flare-up, isotope ratio, magma, southwest Japan

Oral presentation

R6: Plutonic rocks, volcanic rocks and subduction factory

Chairperson: Tatsuhiko Kawamoto (Shizuoka University), Takashi Yuguchi, Atsushi Kamei

Sat. Sep 14, 2024 9:00 AM - 12:00 PM ES Hall (Higashiyama Campus)

11:00 AM - 11:15 AM

[3Lecture-101-11-9add]休憩

The genesis of garnet in the Quaternary rhyolite of Hime-shima volcanic group, EEast Kyushu, Southwest Japan Arc.

*Takehiro HIRAYAMA^{1,2}, Tomoyuki SHIBATA², Masako YOSHIKAWA²

1. Tokyo University, 2. Hiroshima University

Keywords: Garnet, Rhyolite, Xenocryst, Hime-shima volcanic group

K–Ar ages and geochemical compositions of Early Pleistocene andesites from southern Kyushu, Japan

*Raiki YAMADA¹, Masakazu Niwa¹, Koji Umeda², Koshi Yagi³, Taisei Fujiwara³, Ikuo Okada³, Fukuka Kida¹

1. Japan Atomic Energy Agency, 2. Hirosaki University, 3. Hiruzen Institute for Geology and Geochronology

Keywords: Kyushu, Early Pleistocene, Volcanic front, Andesite

Thermal history of pluton formation from Sr diffusion in plagioclase: implications for magma flux estimation and identification of plutons associated with major eruptions

*Tokiyuki Morohoshi¹, Simon Richard Wallis¹

1. The University of Tokyo

Keywords: pluton formation, chemical diffusion, magma flux, Ryoke plutonic rocks

Oral presentation | R2: Crystal structure, crystal chemistry, physical properties of minerals, crystal growth and applied mineralogy

📅 Sat. Sep 14, 2024 9:00 AM - 12:00 PM JST | Sat. Sep 14, 2024 12:00 AM - 3:00 AM UTC | 🏠 ES024 Higashiyama Campus

R2: Crystal structure, crystal chemistry, physical properties of minerals, crystal growth and applied mineralogy

Chairperson: Fumiya Noritake, Mariko Nagashima, Makoto Tokuda

9:00 AM - 9:20 AM JST | 12:00 AM - 12:20 AM UTC

[R2-01] Changes of physical and structural properties of hydrous minerals induced by dehydration of coordinated water molecules

「招待講演」

*Ryo YAMANE¹ (1. Tohoku University)

9:20 AM - 9:35 AM JST | 12:20 AM - 12:35 AM UTC

[R2-02] Phase change of priceite $\text{Ca}_2\text{B}_5\text{O}_7(\text{OH})_5 \cdot \text{H}_2\text{O}$ during thermal decomposition

*Atsushi KYONO¹, Kosuke Yamaguchi¹, Satoru Okada¹, Hiroki Hasegawa¹ (1. Life & Environmental Sciences, University of Tsukuba)

9:35 AM - 9:50 AM JST | 12:35 AM - 12:50 AM UTC

[R2-03] The incorporation mechanisms of water in aluminous orthoenstatite clarified via comprehensive NMR measurements and first-principles calculations

*Xianyu XUE¹, Masami Kanzaki¹, Rongzhang Yin² (1. Okayama University, 2. Peking University)

9:50 AM - 10:05 AM JST | 12:50 AM - 1:05 AM UTC

[R2-04] Evaluation of the influence of polishing condition on hydrogen diffusion in fluorapatite and estimation of the diffusion mechanism

「発表賞エントリー」

*Chikashi Yoshimoto¹, Isao Sakaguchi², Shoichi Itoh¹ (1. Kyoto University, 2. NIMS)

10:05 AM - 10:20 AM JST | 1:05 AM - 1:20 AM UTC

[R2-05] Evaporation kinetics of forsterite in low-pressure H_2 - H_2O atmosphere

「発表賞エントリー」

*Shiori Inada¹, Shogo Tachibana¹ (1. The University of Tokyo)

10:20 AM - 10:30 AM JST | 1:20 AM - 1:30 AM UTC

[3Lecture-201-11-6add] 休憩

10:30 AM - 10:45 AM JST | 1:30 AM - 1:45 AM UTC

[R2-06] Origine of framboidal goethite in sandstone in the Saiki Subgroup of the Lower Shimanto Group

*Hiroaki OHFUJI¹, Seira Konishi¹, Hiroshi Yamamoto² (1. Tohoku Univ. Sci, 2. Kagoshima Univ. Sci)

10:45 AM - 11:00 AM JST | 1:45 AM - 2:00 AM UTC

[R2-07] Precursor phenomenon of ultrananolite crystallization: liquid immiscibility in the volcanic glass

*Mayumi Mujin¹, Keiko Matsumoto², Akira Miyake³, Yohei Igami³, Michihiko Nakamura¹, Akihiko Yokoo³, Mitsuhiro Yoshimoto⁴ (1. Tohoku University, 2. National Institute of Advanced Industrial Science and Technology (AIST), 3. Kyoto University, 4. Mount Fuji Research Institute, Yamanashi Prefectural Government)

11:00 AM - 11:15 AM JST | 2:00 AM - 2:15 AM UTC

[R2-08] Calorimetric constraint of cation disordering enthalpy in MgAl_2O_4 spinel

*Hiroshi KOJITANI¹, Ryuji Matsuki¹, Yuta Nishimura¹, Itaru Ohira¹ (1. Gakushuin University)

11:15 AM - 11:30 AM JST | 2:15 AM - 2:30 AM UTC

[R2-09] Solid solubilities of Na^+ and K^+ in perovskite type SrTiO_3

Ryogo KITAI², *Takaya NAGAI¹, Jun KAWANO¹, Ayako Shinozaki¹ (1. Hokkaido Univ. Faculty Sci., 2. Hokkaido Univ. School Sci.)

11:30 AM - 11:45 AM JST | 2:30 AM - 2:45 AM UTC

[R2-10] Study on the consistency of Li conductivity measurements of $(\text{Li,Li})\text{TiO}_3$ with perovskite structure and Li migration during SIMS measurements

*Isao SAKAGUCHI¹, Tsuyoshi Onishi¹ (1. National Institute for Materials Science)

11:45 AM - 12:00 PM JST | 2:45 AM - 3:00 AM UTC

[R2-11] Structure transition mechanism under high pressure of Fe-Ti-O system phases

*Takamitsu YAMANAKA^{1,2}, Takanori Hattori³, Ho-kwang Mao¹ (1. Center for High Pressure Science and Technology Advanced Research, 2. Graduate School of Science Osaka University, 3. J-PARC)

Changes of physical and structural properties of hydrous minerals induced by dehydration of coordinated water molecules

*Ryo YAMANE¹

1. Tohoku University


Keywords: hydrous mineral, dehydration, vivianite, coordinated water molecule

Phase change of priceite $\text{Ca}_2\text{B}_5\text{O}_7(\text{OH})_5\cdot\text{H}_2\text{O}$ during thermal decomposition

*Atsushi KYONO¹, Kosuke Yamaguchi¹, Satoru Okada¹, Hiroki Hasegawa¹

1. Life & Environmental Sciences, University of Tsukuba

Keywords: Priceite, Borate mineral, Thermal decomposition, Synchrotron X-ray Diffraction



The incorporation mechanisms of water in aluminous orthoenstatite clarified via comprehensive NMR measurements and first-principles calculations

*Xianyu XUE¹, Masami Kanzaki¹, Rongzhang Yin²

1. Okayama University, 2. Peking University

Aluminum is an important minor element in natural orthopyroxene, which has been shown to significantly enhance its water incorporation capacity. However, the incorporation mechanisms remain speculative. We performed a comprehensive one- and two-dimensional ¹H, ²⁹Si and ²⁷Al NMR measurements on hydrous aluminous orthoenstatite (OEn) samples containing 4 to 8 wt% Al₂O₃ synthesized at 1.5 GPa and 900°C, and first-principles calculations on the energy, and NMR and polarized infrared spectra for anhydrous and hydrous aluminous OEn models to clarify the issue. The combined ¹H MAS and static NMR, ¹H double-quantum and triple-quantum MAS NMR, and ²⁷Al-¹H CP MAS NMR and HETCOR results, along with first-principles calculation results, unambiguously revealed that a large part of the incorporated water are present as proton pairs in Mg vacancies adjacent to Al, with one proton of each pair exhibiting significantly weaker hydrogen bonding, and accordingly smaller ¹H chemical shifts and higher OH stretching frequencies, than those in Al-free OEn, as a result of interaction with Al (see Figure). Proton pairs in Mg vacancies remote from Al are minor or absent. Coupled substitutions of Al + H for 1Si and 2Mg (both with weak hydrogen bonding) were also detected, but are less abundant than hitherto considered. Thus, the enhancement of water solubility by Al is achieved dominantly through the modification of the hydrogen bonding of protons in Mg vacancies. Such a mechanism may also be important in other nominally anhydrous mantle minerals. The obtained polarized IR spectral characteristics from our first-principles calculations also allow us to decipher the incorporation mechanisms of water in synthetic OEn of lower Al concentrations and natural orthopyroxenes of diverse origins from the polarized infrared data reported thus far.

Keywords: orthopyroxene, water, aluminum, NMR, first-principles calculation





Figure. ^1H double-quantum (DQ)-single quantum (1Q) correlation MAS NMR spectrum for hydrous aluminous OEn with 8 wt% Al_2O_3 synthesized at 1.5 GPa and 900°C (a), compared with that for a hydrous MgSiO_3 OEn synthesized at 7 GPa and 1200°C (b). The horizontal dashed lines connect the observed proton pairs (labelled pair 1 to 4). Vertical lines are guides for the proton pair 1. Also shown in (c) is an example of energetically favorable 2H in an M_2 vacancy ($(2\text{H})_{\text{M}_2}$) model for aluminous OEn from first-principles calculation. The $\text{Al}^{\text{IV}}\text{-O}_{2\text{b}}\text{-H}\cdots\text{O}_{1\text{b}}$ linkage yields smaller ^1H chemical shift and higher OH stretching frequency than $(2\text{H})_{\text{M}_2}$ defects in Al-free OEn.

Evaluation of the influence of polishing condition on hydrogen diffusion in fluorapatite and estimation of the diffusion mechanism

*Chikashi Yoshimoto¹, Isao Sakaguchi², Shoichi Itoh¹

1. Kyoto University, 2. NIMS

Keywords: apatite, hydrogen, diffusion, Secondary Ion Mass Spectrometry



Evaporation kinetics of forsterite in low-pressure H₂-H₂O atmosphere*Shiori Inada¹, Shogo Tachibana¹

1. The University of Tokyo

Keywords: forsterite, evaporation, kinetics




Fig. 1. 単位面積あたりの蒸発速度 $J \text{ (m}^{-2} \text{ s}^{-1}\text{)}$ と雰囲気ガス組成 P_{H_2}/P_{H_2O} の関係 (全圧 1 Pa, 温度 1400 K) .

Oral presentation

R2: Crystal structure, crystal chemistry, physical properties of minerals, crystal growth and applied mineralogy

Chairperson: Fumiya Noritake, Mariko Nagashima, Makoto Tokuda

Sat. Sep 14, 2024 9:00 AM - 12:00 PM ES024 (Higashiyama Campus)

10:20 AM - 10:30 AM

[3Lecture-201-11-6add]休憩

Origine of framboidal goethite in sandstone in the Saiki Subgroup of the Lower Shimanto Group

*Hiroaki OHFUJI¹, Seira Konishi¹, Hiroshi Yamamoto²

1. Tohoku Univ. Sci, 2. Kagoshima Univ. Sci

The raspberry-like framboidal aggregate is a unique crystallization texture that can be frequently seen in microscopic pyrite in sedimentary rocks and modern sediments. In this study, we conducted micro- and nano-scale analyses using SEM and TEM on iron oxide framboids, which were recently found by Egami et al. (2021) in sandstones of the Saiki Subgroup of the Lower Shimanto Group distributed in western Kagoshima, Japan, to understand their origin. The results show that framboidal aggregate are distributed locally in the sandstone and often form clusters and/or infilled textures. Framboids are mostly surrounded by clay minerals produced by hydrothermal alteration, but no sulfur-containing minerals such as pyrite and sulfate minerals were observed. TEM observation on thin foils prepared by FIB showed that microcrystals that constitute framboids are not single crystals, but are filled by assemblages of nano-sized crystals with a concentric layer pattern, which consists of several alternating layers of goethite nanocrystals and a mixture of clay minerals and amorphous silica. The goethite nanocrystals grew inward from the microcrystal outline, indicating that they were formed by alternation (replacement) of pyrite microcrystals. These observations suggest goethite framboids are pseudomorphs after pyrite framboids, which formed in the interstitial spaces in the formation during diagenesis.

Keywords: Framboida goethite, Framboidal pyrite, Clay mineral, Alteration

Precursor phenomenon of ultrananolite crystallization: liquid immiscibility in the volcanic glass

*Mayumi Mujin¹, Keiko Matsumoto², Akira Miyake³, Yohei Igami³, Michihiko Nakamura¹, Akihiko Yokoo³, Mitsuhiro Yoshimoto⁴

1. Tohoku University, 2. National Institute of Advanced Industrial Science and Technology (AIST), 3. Kyoto University, 4. Mount Fuji Research Institute, Yamanashi Prefectural Government

Keywords: liquid immiscibility, ultrananolite, nanocrystal, volcanic glass

Calorimetric constraint of cation disordering enthalpy in MgAl_2O_4 spinel

*Hiroshi KOJITANI¹, Ryuji Matsuki¹, Yuta Nishimura¹, Itaru Ohira¹

1. Gakushuin University

Keywords: spinel, cation disorder, calorimetry, enthalpy of disordering, thermodynamics




Fig. 1. 様々な陽イオン無秩序化の程度を持つ MgAl_2O_4 スピネルの落下溶解エンタルピー測定結果

Solid solubilities of Na⁺ and K⁺ in perovskite type SrTiO₃

Ryogo KITAI², *Takaya NAGAI¹, Jun KAWANO¹, Ayako Shinozaki¹

1. Hokkaido Univ. Faculty SCi., 2. Hokkaido Univ. School SCi.

In this study, we report the results of synthetic experiments investigating the solid solubilities of Na⁺ and K⁺ in perovskite-type SrTiO₃. According to Shannon (1976), the ionic radii of Na⁺ and K⁺ at a 6-coordinate site are significantly larger than that of Ti⁴⁺, while their ionic radii at a 12-coordinate site are comparable to that of Sr²⁺. Assuming that Na⁺ and K⁺ only exchange with Sr²⁺, this solid solution mechanism would result in the formation of oxygen vacancies.

Keywords: perovskite structure, SrTiO₃, Na, K, oxygen vacancy

Study on the consistency of Li conductivity measurements of (Li,La)TiO₃ with perovskite structure and Li migration during SIMS measurements

*Isao SAKAGUCHI¹, Tsuyoshi Onishi¹

1. National Institute for Materials Science

Introduction

Research and development for carbon neutrality is widely conducted. Among of these, many resources are devoted to these regarding all-solid-state lithium secondary batteries. Many oxides are proposed as electrolytes for all-solid-state lithium batteries. Among of these, we focus on (Li,La)TiO₃ with perovskite structure. Secondary ion mass spectrometry, which is advantageous for light element analysis, is impossible to analyze Li in this material at room temperature. This time, we tried a method to Li battery electrolytes.

Experiment

(Li,La)TiO₃ (hereinafter referred to as LLT) can be purchased in plates, disks, powder, etc. from Toho Titanium Co., Ltd. At this time, we used a plate shape as a standard sample. A comparative sample for this was newly synthesized. The analysis used SIMS (CAMECA, IMS-4f) equipped with NIMS, and O⁻ as primary ions was used and positive secondary ions were detected.

Results and Discussion

Figure 1 shows the analysis results of the purchased LLT. Figure 1(a) is Li image immediately after primary ion irradiation, and Li showed the uniform intensity, and some precipitates with high intensity were observed. Figure 1(b) is Li image after continuous irradiation with O⁻. Obvious non-uniformity of Li was observed in the image. Considering this, the Li intensity distribution in Figure 1(b) reflects the micro-structure within the sample. When primary ions, O⁻, are irradiated during analysis, the irradiated area becomes negative charge, and then Li with positive charge moves to compensate for its charge. The region where Li has changed to a high intensity is a large grain, and the Li conductivity reaches 1×10^{-3} S/cm. It is thought that because the current density of primary ions is small, Li that has migrated to the surface reacts with oxygen and generates LiO_x. it is considered that such non-uniform Li intensity was generated because the amount of Li movement within the disk was small.

Summary

In materials such as the electrolyte in Li batteries, Li moves through the material at room temperature. This research revealed that Li can be moved by changing the charge state on the surface through primary ion irradiation, and that the migration path of Li is the same as that of movement known by electrical measurements.

Keywords: Perovskite structure




Fig. 1. Li images obtained by SIMS. (a) is the first image by O- irradiation. (b) is the image obtained after about 1.5 hours.

Structure transition mechanism under high pressure of Fe-Ti-O system phases

*Takamitsu YAMANAKA^{1,2}, Takanori Hattori³, Ho-kwang Mao¹

1. Center for High Pressure Science and Technology Advanced Research, 2. Graduate School of Science Osaka University, 3. J-PARC

Keywords: ulvöspinel, ilmenite, pseudobrookite

Oral presentation | R5: Extraterrestrial materials

📅 Sat. Sep 14, 2024 9:00 AM - 12:00 PM JST | Sat. Sep 14, 2024 12:00 AM - 3:00 AM UTC | 🏠 ES025
Higashiyama Campus

R5: Extraterrestrial materials

Chairperson: Megumi Matsumoto, Daiki Yamamoto, Yusuke Seto, Toru Matsumoto (Kyoto University)

9:00 AM - 9:15 AM JST | 12:00 AM - 12:15 AM UTC

[R5-01] Revealing Chemical Compositional Variety of Amorphous Silicate Dust around AGB Stars by Condensation Experiment and Spectroscopic Analysis

「発表賞エントリー」

*Hanako Enomoto¹, Aki Takigawa¹, Hiroki Chihara², Chiyoe Koike² (1. UTokyo, 2. Osaka Sangyo univ.)

9:15 AM - 9:30 AM JST | 12:15 AM - 12:30 AM UTC

[R5-02] Formation process of circumstellar dust studied from presolar oxides in carbonaceous chondrite DOM 08006.

「発表賞エントリー」

*Hiroyuki Hashizume¹, Aki Takigawa¹ (1. UTokyo EPSS)

9:30 AM - 9:45 AM JST | 12:30 AM - 12:45 AM UTC

[R5-03] Progress of irreversible chemical reactions on dust moving in a protoplanetary disk

*Lily Ishizaki¹, Shogo Tachibana¹ (1. UT EPS)

9:45 AM - 10:00 AM JST | 12:45 AM - 1:00 AM UTC

[R5-04] Experimental study of the effect of evaporation during temperature rise on type B CAI formation

「発表賞エントリー」

*Yasuaki Tsuruoka¹, Shogo Tachibana¹ (1. UTokyo Sci.)

10:00 AM - 10:20 AM JST | 1:00 AM - 1:20 AM UTC

[R5-05] Chondrule formation by collisions of planetesimals containing volatiles

「招待講演」

*Sin-iti Sirono¹ (1. Navoya Univ.)

10:20 AM - 10:35 AM JST | 1:20 AM - 1:35 AM UTC

[3Lecture-301-10-6add] 休憩

10:35 AM - 10:55 AM JST | 1:35 AM - 1:55 AM UTC

[R5-06] Constraining the true timescale of chondrule formation through chondrule mineral chemistry

「招待講演」

*Kohei Fukuda¹, Noriko T Kita², Makoto Kimura³ (1. Osaka Univ. Sci., 2. UW-Madison, 3. NIPR)

10:55 AM - 11:10 AM JST | 1:55 AM - 2:10 AM UTC

[R5-07] Reproduction Experiments of Radial Pyroxene Chondrules Using a Gas-jet Levitation System under Reducing Conditions

「発表賞エントリー」

*Kana Watanabe¹, Tomoki Nakamura¹, Tomoyo Morita¹ (1. Tohoku Univ. Sci.)

11:10 AM - 11:25 AM JST | 2:10 AM - 2:25 AM UTC

[R5-08] Stratigraphy of poikilitic shergottite parent body(ies) as deduced from their textural and mineral compositional diversities

「発表賞エントリー」

*Sojiro Yamazaki¹, Akira Yamaguchi², Takashi Mikouchi³ (1. Fac. Sci., Univ. Tokyo, 2. Natl. Inst. Polar Res., 3. Univ. Museum, Univ. Tokyo)

11:25 AM - 11:40 AM JST | 2:25 AM - 2:40 AM UTC

[R5-09] High pressure phase transition of Ti-Cr-rich spinel in Asuka 12325 Martian meteorite

*Atsushi TAKENOUCHI¹, Yohei Igami², Takashi Mikouchi³, Akira Miyake², Akira Yamaguchi⁴ (1. The Kyoto Univ. Museum, 2. Kyoto Univ. Sci., 3. The Univ. Museum, The Univ. of Tokyo, 4. NIPR)

11:40 AM - 11:55 AM JST | 2:40 AM - 2:55 AM UTC

[R5-10] Estimation of thermal history using silica minerals in mesosiderites

*Haruka OONO¹, Atsushi Takenouchi², Takashi Mikouchi³, Akira Yamaguchi⁴, Naoji Sugiura⁵, Akira Miyake¹ (1. Kyoto Univ., 2. Museum of Kyoto Univ., 3. Museum of Univ. Tokyo, 4. NiPR, 5. Chiba Int. Tech.)

11:55 AM - 12:00 PM JST | 2:55 AM - 3:00 AM UTC


調整

Revealing Chemical Compositional Variety of Amorphous Silicate Dust around AGB Stars by Condensation Experiment and Spectroscopic Analysis


*Hanako Enomoto¹, Aki Takigawa¹, Hiroki Chihara², Chiyoe Koike²

1. UTokyo, 2. Osaka Sangyo univ.

Keywords: circumstellar dust, amorphous silicate, presolar grain, experiment, spectroscopy



200 nm



200 nm

実験で生成した非晶質ナノ粒子のSTEM-BF像(左)とSTEM-EDS元素マップ(右)

Formation process of circumstellar dust studied from presolar oxides in carbonaceous chondrite DOM 08006.

*Hiroyuki Hashizume¹, Aki Takigawa¹

1. UTokyo EPSS

Keywords: Presolar grains, meteorite, spinel, circumstellar dust, oxides

Progress of irreversible chemical reactions on dust moving in a protoplanetary disk

*Lily Ishizaki¹, Shogo Tachibana¹

1. UT EPS

Keywords: protoplanetary disk, protosolar disk, dust, Monte Carlo simulation

Experimental study of the effect of evaporation during temperature rise on type B CAI formation

*Yasuaki Tsuruoka¹, Shogo Tachibana¹

1. UTokyo Sci.

Keywords: CAIs, melting, evaporation, crystal growth, protosolar disk




図1. メリライトマンタルの組成ゾーニング. 最高温度からの冷却過程のみ水素を導入した試料 (a) では, 組成がほぼ一定の部分が見られる一方, 昇温時から水素を導入した試料 (b) では, 試料中心に向かって緩やかにオケルマナイト (Åk) 成分の割合が上昇している.

Chondrule formation by collisions of planetesimals containing volatiles

*Sin-iti Sirono¹

1. Navoya Univ.

Keywords: chondrule, planetesimal, collision

Oral presentation

R5: Extraterrestrial materials

Chairperson: Megumi Matsumoto, Daiki Yamamoto, Yusuke Seto, Toru Matsumoto (Kyoto University)

Sat. Sep 14, 2024 9:00 AM - 12:00 PM ES025 (Higashiyama Campus)

10:20 AM - 10:35 AM

[3Lecture-301-10-6add]休憩

Constraining the true timescale of chondrule formation through chondrule mineral chemistry

*Kohei Fukuda¹, Noriko T Kita², Makoto Kimura³

1. Osaka Univ. Sci., 2. UW-Madison, 3. NIPR

Keywords: chondrule, Al-Mg chronology, plagioclase, excess Si

Reproduction Experiments of Radial Pyroxene Chondrules Using a Gas-jet Levitation System under Reducing Conditions

*Kana Watanabe¹, Tomoki Nakamura¹, Tomoyo Morita¹

1. Tohoku Univ. Sci.

Keywords: chondrule, pyroxene, levitation experiment, partition coefficients, reduction reaction




図1. 天然RPコンドリュールと実験合成試料の比較。(a)-(b)偏光顕微鏡による観察像(クロスニコル像)、(c)-(d)SEM-BSE像。(d)の実験合成試料は1536°Cで3600秒間温度保持した。

Stratigraphy of poikilitic shergottite parent body(ies) as deduced from their textural and mineral compositional diversities

*Sojiro Yamazaki¹, Akira Yamaguchi², Takashi Mikouchi³

1. Fac. Sci., Univ. Tokyo, 2. Natl. Inst. Polar Res., 3. Univ. Museum, Univ. Tokyo

Keywords: Poikilitic shergottite, Olivine, Pyroxene, Intrusion, Cooling rate

High pressure phase transition of Ti-Cr-rich spinel in Asuka 12325 Martian meteorite

*Atsushi TAKENOUCHI¹, Yohei Igami², Takashi Mikouchi³, Akira Miyake², Akira Yamaguchi⁴

1. The Kyoto Univ. Museum, 2. Kyoto Univ. Sci., 3. The Univ. Museum, The Univ. of Tokyo, 4. NIPR

Keywords: Chromite, Martian meteorite, shock metamorphism, high-pressure phase transition

Estimation of thermal history using silica minerals in mesosiderites

*Haruka OONO¹, Atsushi Takenouchi², Takashi Mikouchi³, Akira Yamaguchi⁴, Naoji Sugiura⁵, Akira Miyake¹

1. Kyoto Univ., 2. Museum of Kyoto Univ., 3. Museum of Univ. Tokyo, 4. NiPR, 5. Chiba Int. Tech.

Keywords: Mesosiderite, Silica minerals, Tridymite, Cristobalite, Meteorites

Oral presentation | R6: Plutonic rocks, volcanic rocks and subduction factory

📅 Sat. Sep 14, 2024 2:00 PM - 3:00 PM JST | Sat. Sep 14, 2024 5:00 AM - 6:00 AM UTC | 🏢 ES Hall
Higashiyama Campus

R6: Plutonic rocks, volcanic rocks and subduction factory

Chairperson: Takashi Hoshide (Akita University)

2:00 PM - 2:20 PM JST | 5:00 AM - 5:20 AM UTC

[R6-12] Zirconium minerals in ultramafic rocks

「招待講演」

*Hikaru Sawada^{1,2}, Ryosuke Oyanagi^{3,2} (1. University of Toyama, 2. JAMSTEC, 3. Kokushikan University)

2:20 PM - 2:35 PM JST | 5:20 AM - 5:35 AM UTC

[R6-13] Estimation of physical conditions during granitic magma crystallization process using melt inclusions in zircon

*Yuka TANIWAKI¹, Satoshi SAITO² (1. Ciatec., Co LTD, 2. Ehime Univ.)

2:35 PM - 2:50 PM JST | 5:35 AM - 5:50 AM UTC

[R6-14] Estimation of TiO_2 activity for accurate determination of quartz crystallization temperature

*Yasuhiro OGITA¹, Takenori Kato², Takashi Yuguchi³ (1. Yamagata Univ., 2. Nagoya Univ., 3. Kumamoto Univ.)

2:50 PM - 3:00 PM JST | 5:50 AM - 6:00 AM UTC

調整

Zirconium minerals in ultramafic rocks

*Hikaru Sawada^{1,2}, Ryosuke Oyanagi^{3,2}

1. University of Toyama, 2. JAMSTEC, 3. Kokushikan University

Keywords: Serpentinite, Peridotite, Zircon, Baddeleyite

Estimation of physical conditions during granitic magma crystallization process using melt inclusions in zircon

*Yuka TANIWAKI¹, Satoshi SAITO²

1. Ciatec.,Co LTD, 2. Ehime Univ.

Keywords: melt inclusion, granite, zircon, physical conditions

Estimation of TiO_2 activity for accurate determination of quartz crystallization temperature

*Yasuhiro OGITA¹, Takenori Kato², Takashi Yuguchi³

1. Yamagata Univ., 2. Nagoya Univ., 3. Kumamoto Univ.

No abstract in English.

Keywords: Quartz, Cathodoluminescence pattern, Ti concentration, TiO_2 activity, Tono Plutonic Complex

Oral presentation | R2: Crystal structure, crystal chemistry, physical properties of minerals, crystal growth and applied mineralogy

📅 Sat. Sep 14, 2024 2:00 PM - 3:30 PM JST | Sat. Sep 14, 2024 5:00 AM - 6:30 AM UTC | 🏢 ES024
Higashiyama Campus

R2: Crystal structure, crystal chemistry, physical properties of minerals, crystal growth and applied mineralogy

Chairperson: Kazuki 小松 Komatsu

2:00 PM - 2:30 PM JST | 5:00 AM - 5:30 AM UTC

[R2-12] Structural stabilization of pyrite-type compounds and chemical bonds of anions.

「招待講演」

*Akira YOSHIASA¹, Makoto Tokuda¹, Ginga Kitahara¹, Akihiko Nakatsuka², Kazumasa Sugiyama³ (1. Kumamoto University, 2. Yamaguchi University, 3. Tohoku University)

2:30 PM - 2:45 PM JST | 5:30 AM - 5:45 AM UTC

[R2-13] Formation of Fundamental Building Blocks Composed of Calcium Borate Minerals

「発表賞エントリー」

*Kosuke Yamaguchi¹, Atsushi Kyono², Satoru Okada¹, Hiroki Hasegawa¹ (1. Univ of Tsukuba. Grad. Sch. of Life and Env., Sci, 2. Univ of Tsukuba. Life and Environmental Science)

2:45 PM - 3:00 PM JST | 5:45 AM - 6:00 AM UTC

[R2-14] Crystal structures of kimuraite-(Y) and lokkaite-(Y)

*Ritsuro Miyawaki¹, Koichi Momma¹ (1. National Museum of Nature and Science)

3:00 PM - 3:15 PM JST | 6:00 AM - 6:15 AM UTC

[R2-15] Thermal Vibration of Oxygen atoms in Keatite Crystal

*Fumiya NORITAKE¹ (1. University of Yamanashi)

3:15 PM - 3:30 PM JST | 6:15 AM - 6:30 AM UTC

[R2-16] Development of a simulation method for electron back-scattered diffraction based on dynamical theory and its application to mineralogy

*Yusuke SETO¹, Masahiro Ohtsuka² (1. Osaka Metropolitan Univ. Sci., 2. Nagoya Univ. Sci. IMaSS.)

Structural stabilization of pyrite-type compounds and chemical bonds of anions.

*Akira YOSHIASA¹, Makoto Tokuda¹, Ginga Kitahara¹, Akihiko Nakatsuka², Kazumasa Sugiyama³

1. Kumamoto University, 2. Yomaguchi University, 3. Tohoku University

Precise structural analyses of various compounds with pyrite-type structure were performed by selecting single crystals from synthetic and natural mineral crystals. The measurements of Debye temperatures of the compounds will be reported as one of the usefulness of single crystal diffraction method. Certain regularities regarding structural stabilization and physical properties are found in pyrite-type compounds.

Keywords: AuSb₂, RuS₂, pyrite-type compounds, single crystal diffraction, Debye temperature




Fig. 1. Compositional dependences in M-S and S-S bonding distances with respect to Os content in the erlichmanite - laurite solid solutions ($(\text{Ru,Rh})_{1-y}\text{Os}_y\text{S}_2$) ($y=0.07-0.10$)[3].

Formation of Fundamental Building Blocks Composed of Calcium Borate Minerals

*Kosuke Yamaguchi¹, Atsushi Kyono², Satoru Okada¹, Hiroki Hasegawa¹

1. Univ of Tsukuba. Grad. Sch. of Life and Env., Sci, 2. Univ of Tsukuba. Life and Environmental Science

Keywords: hydrothermal method, dissolution-reprecipitation, gowerite, priceite, nobleite

Crystal structures of kimuraite-(Y) and lokkaite-(Y)

*Ritsuro Miyawaki¹, Koichi Momma¹

1. National Museum of Nature and Science

The crystal structures of kimuraite-(Y) and lokkaite-(Y) were re-examined by means of the twining and statistical ordering models with the space groups following to the individual extinction rules in their single crystal XRD data. The crystal structure of kimuraite-(Y) determined with the space group $I222$ is harmonic with that previously determined with the space group $I2cb$. The crystal structure of lokkaite-(Y) was analyzed with the monoclinic space group Cm to determine the mean structure in the $C222$. Each crystal structure of kimuraite-(Y) and lokkaite-(Y) consists of two types of layers: a flat calcium layer and a corrugated rare earth layer (Figure). The rare earth atoms are coordinated to 9 oxygen atoms. The 9-coordinated polyhedra are connected with the wedging triangle of carbonate anion to form the corrugated rare earth layer, as those in the crystal structure of tenerite-(Y). Calcium atoms statistically order into the Ca sites at the middle of the flat calcium layer. The C and O atoms neighboring to the Ca atoms should be shifted with the statistical occupancies. The crystal structures of kimuraite-(Y) and lokkaite-(Y) differ from each other in the sequence of stacking layers corresponding to the ratio of calcium and rare earths atoms.


Keywords: kimuraite-(Y), lokkaite-(Y), crystal structure

Kimuraite-(Y)

$a = 9.2673(7)$

$b = 24.0435(2)$


$c = 6.0537(2)$

**Lokkaite-(Y)**

$a = 9.242(4)$

$b = 39.328(18)$

$c = 6.110(2)$

**Tengerite-(Y)**

$a = 9.157(2)$

$b = 15.114(6)$

$c = 6.078(4)$




図 木村石、ロッカ石、テンゲル石の結晶構造

Thermal Vibration of Oxygen atoms in Keatite Crystal

*Fumiya NORITAKE¹

1. University of Yamanashi

Keywords: Keatite, Negative Thermal Expansion, Molecular Dynamics Simulation

Development of a simulation method for electron back-scattered diffraction based on dynamical theory and its application to mineralogy

*Yusuke SETO¹, Masahiro Ohtsuka²

1. Osaka Metropolitan Univ. Sci., 2. Nagoya Univ. Sci. IMaSS.

Keywords: Electron back-scattered diffraction, Bloch wave, Dynamical theory




図. 実測のEBSDパターンと発表者らのシミュレーション結果との比較

Oral presentation | R5: Extraterrestrial materials

📅 Sat. Sep 14, 2024 2:00 PM - 3:30 PM JST | Sat. Sep 14, 2024 5:00 AM - 6:30 AM UTC | 🏠 ES025
Higashiyama Campus

R5: Extraterrestrial materials

Chairperson: Megumi Matsumoto, Daiki Yamamoto, Yusuke Seto, Toru Matsumoto (Kyoto University)

2:00 PM - 2:15 PM JST | 5:00 AM - 5:15 AM UTC

[R5-11] Mineralogy of micrometeoroids captured in silica aerogel aboard the ISS by the Dandelion Project

*Takaaki NOGUCHI¹, Akira MIYAKE¹, Hikaru Yabuta², Yoko Kebukawa³, Daiki Suga⁴, Makoto Tabata⁵, Kyoko Okudaira⁶, Akihiko Yamagishi⁷, Hajime Yano⁸ (1. Kyoto University, 2. Hiroshima University, 3. Tokyo Institute of Technology, 4. JASRI, 5. Chiba University, 6. University of Aizu, 7. Tokyo University of Pharmacy and Life Sciences, 8. JAXA)

2:15 PM - 2:30 PM JST | 5:15 AM - 5:30 AM UTC

[R5-12] STEM analysis of sodium-rich grains recovered from asteroid Ryugu

*Toru MATSUMOTO¹, Takaaki Noguchi¹, Toru Araki², Hayato Yuzawa², Akira Miyake¹ (1. Kyoto University, 2. UVSOR)

2:30 PM - 2:45 PM JST | 5:30 AM - 5:45 AM UTC

[R5-13] Important occurrence of amorphous hydrated Na-bearing Mg phosphate in Asteroid Ryugu samples and CI chondrites

*Takashi MIKOUCHI¹, Hideto Yoshida², Satoru Hayashi², Minami Masuda², Hiroyuki Kagi², Kazumasa Sugiyama³, Tomoki Nakamura⁴, Michael Zolensky⁵ (1. Univ. Museum, Univ. of Tokyo, 2. Fac. of Sci., Univ. of Tokyo, 3. Inst. Mat. Res., Tohoku Univ., 4. Fac. of Sci., Tohoku Univ., 5. NASA-JSC)

2:45 PM - 3:00 PM JST | 5:45 AM - 6:00 AM UTC

[R5-14] On the relationship between lithological classification and degree of aqueous alteration in Asteroid Ryugu samples

*Minami Masuda¹, Takashi Mikouchi², Hideto Yoshida¹, Tomoki Nakamura³, Michael Zolensky⁴ (1. Univ. Tokyo, Sci., 2. Univ. Tokyo, Univ. Museum, 3. Tohoku Univ., Sci., 4. NASA-JSC)

3:00 PM - 3:15 PM JST | 6:00 AM - 6:15 AM UTC

[R5-15] Highly primitive clasts with GEMS-like grains in Ryugu C0002 sample

*Megumi MATSUMOTO¹, Akira Tsuchiyama^{2,3}, Masahiro Yasutake⁴, Tomoki Nakamura¹ (1. Tohoku Univ., 2. Ritsumeikan Univ., 3. GIG, 4. JASRI/SPring-8)

3:15 PM - 3:30 PM JST | 6:15 AM - 6:30 AM UTC

[R5-16] Aqueous alteration experiments using GEMS simulant particles

*Akira TSUCHIYAMA^{1,2}, Satomi Enju³, Haiyang Xian¹, Mingqi Sun¹, Ding Xing¹, Yuanyuan Wang⁴, Zhaolu He¹, Jing Du¹, Shan Li¹, Jiabin Xi¹, Xiaojun Lin¹, Jian Di¹, Yuto Imura⁵ (1. Guangzhou Inst. Geochem., 2. Ritsumeikan Univ., 3. Ehime Univ. Sci., 4. Tokyo Univ. Sci., 5. Kyoto Univ. Sci.)

Mineralogy of micrometeoroids captured in silica aerogel aboard the ISS by the Dandelion Project

*Takaaki NOGUCHI¹, Akira MIYAKE¹, Hikaru Yabuta², Yoko Kebukawa³, Daiki Suga⁴, Makoto Tabata⁵, Kyoko Okudaira⁶, Akihiko Yamagishi⁷, Hajime Yano⁸

1. Kyoto University, 2. Hiroshima University, 3. Tokyo Institute of Technology, 4. JASRI, 5. Chiba University, 6. University of Aizu, 7. Tokyo University of Pharmacy and Life Sciences, 8. JAXA

Keywords: micrometeoroid, TANPOPO mission, International Space Station, pyroxene

STEM analysis of sodium-rich grains recovered from asteroid Ryugu

*Toru MATSUMOTO¹, Takaaki Noguchi¹, Toru Araki², Hayato Yuzawa², Akira Miyake¹

1. Kyoto University, 2. UVSOR

Keywords: Ryugu

Important occurrence of amorphous hydrated Na-bearing Mg phosphate in Asteroid Ryugu samples and CI chondrites

*Takashi MIKOUCHI¹, Hideto Yoshida², Satoru Hayashi², Minami Masuda², Hiroyuki Kagi², Kazumasa Sugiyama³, Tomoki Nakamura⁴, Michael Zolensky⁵

1. Univ. Museum, Univ. of Tokyo, 2. Fac. of Sci., Univ. of Tokyo, 3. Inst. Mat. Res., Tohoku Univ., 4. Fac. of Sci., Tohoku Univ., 5. NASA-JSC

Keywords: Mg phosphate, Asteroid Ryugu, CI chondrites, Aqueous alteration, Asteroid Bennu

On the relationship between lithological classification and degree of aqueous alteration in Asteroid Ryugu samples

*Minami Masuda¹, Takashi Mikouchi², Hideto Yoshida¹, Tomoki Nakamura³, Michael Zolensky⁴

1. Univ. Tokyo, Sci., 2. Univ. Tokyo, Univ. Museum, 3. Tohoku Univ., Sci., 4. NASA-JSC

Keywords: Asteroid Ryugu, CI chondrite, aqueous alteration, brecciation

Highly primitive clasts with GEMS-like grains in Ryugu C0002 sample

*Megumi MATSUMOTO¹, Akira Tsuchiyama^{2,3}, Masahiro Yasutake⁴, Tomoki Nakamura¹

1. Tohoku Univ., 2. Ritsumeikan Univ., 3. GIG, 4. JASRI/SPring-8

Keywords: asteroid (162173) Ryugu, GEMS, Aqueous alteration

Aqueous alteration experiments using GEMS simulant particles

*Akira TSUCHIYAMA^{1,2}, Satomi Enju³, Haiyang Xian¹, Mingqi Sun¹, Ding Xing¹, Yuanyuan Wang⁴, Zhaolu He¹, Jing Du¹, Shan Li¹, Jiaxin Xi¹, Xiaoju Lin¹, Jian Di¹, Yuto Imura⁵

1. Guangzhou Inst. Geochem., 2. Ritsumeikan Univ., 3. Ehime Univ. Sci, 4. Tokyo Univ. Sci., 5. Kyoto Univ. Sci.

Samples from the asteroid Ryugu by the Hayabusa2 spacecraft correspond to CI chondrites (e.g., [1]). In this study, to reproduce the initial stages of aqueous alteration of Ryugu parent body [2], we performed an aqueous alteration experiment using GEMS simulant particles, prepared by the induction thermal plasma (ITP) [3] with CI or GEMS average composition. Aqueous solutions of formic acid, hexamethylenetetramine, ammonium acetate, and/or ammonium hydrogen carbonate were heated at 200°C for 168~1550 hours in a sealed Teflon container with the GEMS simulant with the rock/water ratio of 5 (or 1). Run products were analyzed by XRD, SEM/EDS, TEM/EELS, and micro-Raman spectroscopy. In all runs, low-crystallinity hydrous layered silicates (Mg silicate hydrate: M-S-H) were formed. Hematite, maghemite or magnetite + hematite formed as Fe oxides consisted of fine particles (~a few 10 nm) and occasionally dendrites within a M-S-H matrix (Fig. 1a). The run products may represent the very early state of aqueous alteration. In most runs, small amounts of anhydrite were formed, but no carbonate formed. Organic crystals and ammonium sulfates also formed. IOM with G and D bands were confirmed by Raman spectroscopy. The run products are aggregates of several 10 μm sized clasts with different porosities (Fig. 1b), which reflects the original texture of the starting material. Ryugu sample matrix of similar texture [2] may be inherited from that formed by accumulation in the parent body. The present experiment was not fully successful in reproducing the minerals and their morphologies as seen in the Ryugu samples. In particular, the formation of maghemite and anhydrite indicates that further experiments under more reducing conditions are necessary. [1] Nakamura T. et al. (2023) *Science*, 379, eabn867. [2] Tsuchiyama A. et al. (2024) *GCA*, 375, 146-172. [3] Enju S. et al. (2022) *A&A*, 661, A121.

Keywords: Ryugu samples, Mg-S-H, carbonaceous chondrites, GEMS-like material, induction thermal plasma




Figure 1. (a) SEM/BSE image of Run 11 (Cl composition, water + NH_4HCO_3 : $X(\text{CO}_2)/X(\text{H}_2\text{O})=0.0317$, $X(\text{NH}_3)/X(\text{H}_2\text{O})=0.0317$, $W/R=5$, 200°C , 1550 hrs.). Different brightness of matrix roughly corresponds to different porosity. (b) TEM/bright field image of Run 07 (Cl composition, water + CH_2O_2 : $X(\text{CO}_2)/X(\text{H}_2\text{O})=0.0435$, $X(\text{NH}_3)/X(\text{H}_2\text{O})=0$, $W/R=5$, 200°C , 168 hrs.) showing a dendritic aggregate of maghemite.

Poster presentation | R2: Crystal structure, crystal chemistry, physical properties of minerals, crystal growth and applied mineralogy

📅 Sat. Sep 14, 2024 12:30 PM - 2:00 PM JST | Sat. Sep 14, 2024 3:30 AM - 5:00 AM UTC | 🏢 Entrance Hall Higashiyama Campus

R2: Crystal structure, crystal chemistry, physical properties of minerals, crystal growth and applied mineralogy

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R2-P-01] Advanced four-circle goniometer system for single-crystal diffraction measurements at BL-10A, PF: anomalous X-ray scattering method & in situ high-pressure experiments

*Takahiro KURIBAYASHI¹, Ryo Yamane², Keita Harada¹, Toshiro Nagase³, Kazumasa Sugiyama² (1. Tohoku University (Sci), 2. Tohoku University (IMR), 3. Tohoku University (Mus))

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R2-P-02] Fine Structural Analysis using Single-Crystal Anomalous X-ray Scattering in KEK Photon Factory BL-6C

*Makoto TOKUDA¹, Takumi Ichimura², Koichi Momma⁶, Ritsuro Miyawaki⁶, Takashi Mikouchi⁴, Akira Yoshiasa³, Kazumasa Sugiyama⁵ (1. IINa, Kumamoto Univ., 2. Tohoku Univ. Eng, 3. Kumamoto Univ. Sci, 4. Univ. of Tokyo UMUT, 5. IMR, Tohoku Univ., 6. Nat'l. Mus. Nat. Sci.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R2-P-03] Evaluation of Fe K-edge XANES anisotropy and accurate determination of micro-area Fe²⁺/Fe³⁺ in clinopyroxene using multivariate analysis

「発表賞エントリー」

*Taisuke Ito¹, Simon Richard Wallis¹, Yoshio Takahashi¹ (1. The Univ. of Tokyo, EPS)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R2-P-04] Fine structural analysis of Diopside (Violan) containing Mn and Fe

「発表賞エントリー」

*Hayato Chiba¹, Kazumasa Sugiyama², Makoto Tokuda³, Takashi Mikouchi⁴ (1. Tohoku Univ. Eng., 2. Tohoku Univ. IMR, 3. Kumamoto Univ. IINa, 4. UMUT)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R2-P-05] Single-crystal X-ray diffraction study of a largely Cs-exchanged natural Ca-chabazite: crystal-chemical factors for its excellent Cs-exchange ability

Naomi KAWATA¹, Ryo FURUHASHI¹, Keiko FUJIWARA¹, Makio OHKAWA², *Akihiko NAKATSUKA¹ (1. Yamaguchi Univ. Sci. Tech. Innov., 2. Hiroshima Univ. Adv. Sci. Eng.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R2-P-06] Distribution of Mn³⁺ and Mn⁴⁺ studied by X anomalous scattering: the details of 3×4 tunneled structure formed by MnO₆ octahedron in woodruffite

「発表賞エントリー」

*Christopher Miyazaki¹, Ryo Yamane², Kazumasa Sugiyama², Makoto Tokuda³, Takeshi Mikouchi⁴ (1. Tohoku Univ. Eng., 2. Tohoku Univ. IMR, 3. Kumamoto Univ. IINa, 4. UMUT)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R2-P-07] Mg²⁺ and Ba²⁺ ion-exchange of GTS-type Na titanosilicate and its structural characterization

*Keiko FUJIWARA¹, Naomi Kawata¹, Akihiko Nakatsuka¹ (1. Yamaguchi Univ. Sci. Tech. Innov.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R2-P-08] Distribution of Fe³⁺ among tetrahedral sites and crystal structural change of gehlenite-Fe³⁺ gehlenite

*Maki HAMADA¹, Mariko Nagashima² (1. Kanazawa University, 2. Yamaguchi University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R2-P-09] Crystallographic site preference of impurity elements in olivine by ALCHEMI method

*Yohei IGAMI¹, Akira MIYAKE¹, Ryo WAKABAYASHI¹, Norikatsu AKIZAWA² (1. Kyoto University, 2. The University of Tokyo)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R2-P-10] Precipitated structures of pyroxene lamellar with various directions

*Shoichi TOH¹ (1. Fukuoka University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R2-P-11] Crystal chemistry of bridgmanite with Al/Fe contents expected in MORB component

*Akihiko NAKATSUKA¹, Hiroshi FUKUI², Seiji KAMADA³, Naohisa HIRAO², Makio OHKAWA⁴, Kazumasa SUGIYAMA⁵, Takashi YOSHINO⁶ (1. Yamaguchi Univ. Sci. Tech. Innov., 2. JASRI, 3. AD Science Inc., 4. Hiroshima Univ. Adv. Sci. Eng., 5. Tohoku Univ. IMR, 6. Okayama Univ. IPM)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R2-P-12] Mechanism of the pressure-induced crystallization of amorphous calcium carbonate

*Chiho Morita¹, Hiroki Kobayashi¹, Hiroyuki Kagi¹ (1. UTokyo. Sci.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R2-P-13] Differences in the effects of Li and Cs on the calcitization of vaterite.

*Noboru FURUKAWA¹, Ami Kinoda² (1. Chiba University, 2. NIWAKA)

Advanced four-circle goniometer system for single-crystal diffraction measurements at BL-10A, PF: anomalous X-ray scattering method & in situ high-pressure experiments

*Takahiro KURIBAYASHI¹, Ryo Yamane², Keita Harada¹, Toshiro Nagase³, Kazumasa Sugiyama²

1. Tohoku University (Sci), 2. Tohoku University (IMR), 3. Tohoku University (Mus)


Keywords: anomalous X-ray scattering, In situ single-crystal X-ray diffraction, synchrotron radiation

Fine Structural Analysis using Single-Crystal Anomalous X-ray Scattering in KEK Photon Factory BL-6C

*Makoto TOKUDA¹, Takumi Ichimura², Koichi Momma⁶, Ritsuro Miyawaki⁶, Takashi Mikouchi⁴, Akira Yoshiasa³, Kazumasa Sugiyama⁵

1. IINa, Kumamoto Univ., 2. Tohoku Univ. Eng, 3. Kumamoto Univ. Sci, 4. Univ. of Tokyo UMUT, 5. IMR, Tohoku Univ., 6. Nat'l. Mus. Nat. Sci.

Keywords: Anomalous X-ray Scattering, Single-crystal X-ray Diffraction, Element Distribution, Valence-Difference Contrast



Evaluation of Fe K-edge XANES anisotropy and accurate determination of micro-area Fe²⁺/Fe³⁺ in clinopyroxene using multivariate analysis

*Taisuke Ito¹, Simon Richard Wallis¹, Yoshio Takahashi¹

1. The Univ. of Tokyo, EPS


Keywords: XAFS spectroscopy, clinopyroxene, Fe²⁺/Fe³⁺, multivariate analysis

Fine structural analysis of Diopside (Violan) containing Mn and Fe

*Hayato Chiba¹, Kazumasa Sugiyama², Makoto Tokuda³, Takashi Mikouchi⁴

1. Tohoku Univ. Eng., 2. Tohoku Univ. IMR, 3. Kumamoto Univ. IINa, 4. UMUT

Keywords: Diopside, Violan, AXS, XANES



Single-crystal X-ray diffraction study of a largely Cs-exchanged natural Ca-chabazite: crystal-chemical factors for its excellent Cs-exchange ability

Naomi KAWATA¹, Ryo FURUHASHI¹, Keiko FUJIWARA¹, Makio OHKAWA², *Akihiko NAKATSUKA¹

1. Yamaguchi Univ. Sci. Tech. Innov., 2. Hiroshima Univ. Adv. Sci. Eng.

The amount of contaminated water generated by the Fukushima Daiichi Nuclear Power Plant accident, caused by the Great East Japan Earthquake (March 11, 2011), continues to increase and has become a serious problem. Among the radioactive elements contained in contaminated water, ¹³⁷Cs (30.1 years) and ⁹⁰Sr (28.8 years) with a long half-life were released in large quantities by the nuclear accident. The search and development of ion-exchangers for highly efficient recovery and removal of these radioactive elements are eagerly desired. Currently, natural zeolite minerals such as chabazite, mordenite and clinoptilolite are practically utilized at the crippled nuclear power plant as the radioactive element removers. To design and develop highly-effective radioactive element removers, it is significant to reveal the relationship between the crystal structural features and the exchange abilities for radioactive ions. Here we conduct the single-crystal X-ray diffraction study of a hydrated natural Ca-chabazite from Burnt Cabin Creek, Spray, Wheeler Co., Oregon, USA and its largely Cs-exchanged form. The sample compositions are $(\text{Ca}_{1.86}\text{Na}_{0.13}\text{K}_{0.09})(\text{Al}_{3.98}\text{Si}_{8.03})\text{O}_{24}\cdot 12.38\text{H}_2\text{O}$ for the former and $(\text{Cs}_{2.66}\text{Ca}_{0.45}\text{Na}_{0.04}\text{K}_{0.10})(\text{Al}_{4.04}\text{Si}_{8.04})\text{O}_{24}\cdot 8.52\text{H}_2\text{O}$ for the latter.

We revealed that the Cs-exchanged form has the essentially ten occupied-sites in extraframework: four water sites (OW2', OW3, OW4, OW5), essentially two Cs sites (Cs1/Cs1', Cs2) and four Ca sites (Ca1, Ca2, Ca3, Ca4). The Cs⁺ ions more preferentially occupy the Cs1/Cs1' site, located at/around the centers of the 8-membered ring windows, than the Cs2 site. In terms of interatomic distances, the coordination environments of the extraframework species in the chabazite crystals before and after the Cs-exchange treatment are discussed. In particular, both samples have a common feature that possible hydrogen bonds are relatively weak between water molecules and framework O atoms, whereas those are relatively strong between water molecules. On the basis of these findings, we discuss the crystal-chemical key factors for an excellent Cs-exchange ability of chabazite as a highly efficient radioactive-element remover.

Keywords: Ca-chabazite, Single-crystal X-ray Diffraction, Cs-exchange ability

Distribution of Mn^{3+} and Mn^{4+} studied by X anomalous scattering: the details of 3×4 tunneled structure formed by MnO_6 octahedron in woodruffite

*Christopher Miyazaki¹, Ryo Yamane², Kazumasa Sugiyama², Makoto Tokuda³, Takeshi Mikouchi⁴

1. Tohoku Univ. Eng., 2. Tohoku Univ. IMR, 3. Kumamoto Univ. IINa, 4. UMUT

Keywords: woodruffite




Fig. 1 (a) Woodruffiteの結晶構造 (VESTAにより描画)
 (b) MnのX線異常散乱によって求めたコントラストマップ (等高線間隔: $1.2 \text{ e}/\text{\AA}^3$)

Mg²⁺ and Ba²⁺ ion-exchange of GTS-type Na titanosilicate and its structural characterization

*Keiko FUJIWARA¹, Naomi Kawata¹, Akihiko Nakatsuka¹

1. Yamaguchi Univ. Sci. Tech. Innov.

GTS-type microporous titanosilicates have a three-dimensional tunnel-type structure. In this structure, four TiO₆ octahedra linked by edge-sharing form a Ti₄O₁₆ cluster; the clusters are linked through SiO₄ tetrahedra to form a three-dimensional framework with an interconnected pore system involving cavities of 8-ring channels, occupied by the alkali-metal ions and adsorbed water molecules. It is reported that Na-GTS, Na₄[(TiO)₄(SiO₄)₃]·6H₂O, crystallizes in a rhombohedral phase with space group *R3m*; its unit cell [*a* = 7.8123 (6) Å, *α* = 88.794 (9) °] is very close to cubic cell and often described as a pseudocubic cell. In this study, we focused on Mg²⁺ and Ba²⁺, which have the different ionic size from and the same valence as the radioactive elements Sr²⁺. We investigated the effects of ionic size and valence on the Mg²⁺- and Ba²⁺-exchange properties of Na-GTS.

We have prepared the ion-exchanged forms, Na_{4(1-x)}}M_{2x}[(TiO)₄(SiO₄)₃]·yH₂O (M: Mg, Ba), by treating Na-GTS with MgCl₂ and BaCl₂ aqueous solutions (C_M = 0 ~ 7.9 mol/L) at 25 °C for 24 hours. The maximum ion-exchange rates of Mg and Ba were *x* = 0.76 and 0.86, respectively. We evaluated the unit-cell parameters from the powder X-ray diffraction (XRD) by a profile fitting method assuming a rhombohedral unit-cell and the H₂O content (*y*) by TG. The increase in the ion-exchange rate (*x*) decreases the unit-cell volumes (*V*) and the H₂O contents (*y*) of the Ba²⁺-exchanged samples, but increases those of the Mg²⁺-exchanged samples. The increase in *V* of the Mg²⁺-exchanged samples can be due to the increase in the *y* value; it can be brought about by the increase in the free space in the pores due to the exchange of smaller divalent Mg²⁺ with larger monovalent Na⁺. The decrease in *V* of Ba²⁺-exchanged samples can be explained by the interpretation that the decrease in the number of exchangeable-cations due to the exchange of divalent Ba²⁺ with monovalent Na⁺ yields the decrease in the number of water molecules coordinating to it. In these samples, the cation distribution of Mg²⁺ and Ba²⁺ in the cavities will be discussed from the comparison of the observed and simulated XRD patterns.

Keywords: Na-GTS, Ion-exchange, Powder XRD




Fig. 1. Observed XRD patterns of Mg^{2+} and Ba^{2+} -exchanged Na-GTS.

Distribution of Fe³⁺ among tetrahedral sites and crystal structural change of gehlenite-Fe³⁺ gehlenite

*Maki HAMADA¹, Mariko Nagashima²

1. Kanazawa University, 2. Yamaguchi University

Keywords: synthetic melilite, crystal structural change, gehlenite

Crystallographic site preference of impurity elements in olivine by ALCHEMI method

*Yohei IGAMI¹, Akira MIYAKE¹, Ryo WAKABAYASHI¹, Norikatsu AKIZAWA²

1. Kyoto University, 2. The University of Tokyo


Keywords: olivine, cation site distribution, analytical electron microscopy, ALCHEMI

Precipitated structures of pyroxene lamellar with various directions

*Shoichi TOH¹

1. Fukuoka University

Keywords: augite, pigeonite, exsolution, texture



Crystal chemistry of bridgmanite with Al/Fe contents expected in MORB component

*Akihiko NAKATSUKA¹, Hiroshi FUKUI², Seiji KAMADA³, Naohisa HIRAO², Makio OHKAWA⁴, Kazumasa SUGIYAMA⁵, Takashi YOSHINO⁶

1. Yamaguchi Univ. Sci. Tech. Innov., 2. JASRI, 3. AD Science Inc., 4. Hiroshima Univ. Adv. Sci. Eng., 5. Tohoku Univ. IMR, 6. Okayama Univ. IPM

Bridgmanite (simplified formula MgSiO_3) is the most abundant constituent in the Earth's lower mantle. The effects of the incorporation of Fe and Al into bridgmanite can have a large effect on the physical properties and rheology of the lower mantle. Bridgmanite formed from a mid-ocean ridge basalt (MORB) component of subducting slabs contains larger amounts of Fe and Al than that formed from a pyrolytic composition. This difference in bridgmanite composition can cause a difference in the incorporation mechanism of Fe and Al into the crystal structure between subducting slabs and their surrounding lower mantle. This should cause heterogeneity in the physical properties and rheology of the lower mantle. Elucidating the crystal chemistry of bridgmanite formed from the MORB composition is a key to resolving this issue. The precise crystal chemistry examined employing a single crystal is, therefore, significant for gaining a detailed understanding of lower-mantle dynamics. In particular, the use of ^{57}Fe -Mössbauer spectroscopy is indispensable for distinguishing the valence and spin states of Fe, which cannot be directly observed by X-ray diffraction. For this purpose, we characterize $\text{Mg}_{0.662}\text{Fe}_{0.338}\text{Si}_{0.662}\text{Al}_{0.338}\text{O}_3$ bridgmanite single-crystal, with the Fe and Al contents expected in MORB, by a combination of single-crystal X-ray diffraction, synchrotron ^{57}Fe -Mössbauer spectroscopy conducted at SPring-8 BL10XU, and electron probe microanalysis.

The present study reveals that the charge-coupled substitution $^{\text{A}}\text{Mg}^{2+} + ^{\text{B}}\text{Si}^{4+} \leftrightarrow ^{\text{A}}\text{Fe}^{3+}(\text{high-spin}) + ^{\text{B}}\text{Al}^{3+}$ is predominant in the incorporation of Fe and Al into the practically eightfold-coordinated A-site and the sixfold-coordinated B-site in bridgmanite structure. The incorporation of both cations via this substitution enhances the structural distortion due to the tilting of BO_6 octahedra, yielding the unusual expansion of mean $\langle\text{A-O}\rangle$ bond-length due to flexibility of A-O bonds for the structural distortion, in contrast to mean $\langle\text{B-O}\rangle$ bond-length depending reasonably on the ionic radius effect. Moreover, we imply the phase transition behavior and the elasticity of bridgmanite in slabs subducting into deeper parts of the lower mantle, in terms of the relative compressibility of AO_{12} (practically AO_8) and BO_6 polyhedra.

Keywords: Bridgmanite, MORB, Single-crystal X-ray diffraction, Synchrotron Mössbauer

Mechanism of the pressure-induced crystallization of amorphous calcium carbonate

*Chiho Morita¹, Hiroki Kobayashi¹, Hiroyuki Kagi¹

1. UTokyo. Sci.

Keywords: Amorphous calcium carbonate, Calcite, High pressure





図. ACCの圧力誘起結晶化の時分割X線回折パターン

Differences in the effects of Li and Cs on the calcitization of vaterite.

*Noboru FURUKAWA¹, Ami Kinoda²

1. Chiba University, 2. NIWAKA

Keywords: vaterite, calcitization, Lithium, Cesium



Poster presentation | R5: Extraterrestrial materials

📅 Sat. Sep 14, 2024 12:30 PM - 2:00 PM JST | Sat. Sep 14, 2024 3:30 AM - 5:00 AM UTC | 🏢 Entrance
Hall Higashiyama Campus

R5: Extraterrestrial materials

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R5-P-01] Dehydration of Newberyite($\text{MgHPO}_4 \cdot 3\text{H}_2\text{O}$) under low pressures

*Yukako Matsumoto¹, Shogo Tachibana¹ (1. UTokyo)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R5-P-02] Experimental simulation of oxygen isotope exchange between chondrule melt and low-pressure water vapor

*Daiki YAMAMOTO¹, Noriyuki Kawasaki² (1. Kyushu University, 2. Hokkaido University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R5-P-03] Early impact events recorded in anomalous eucrites

*Akira YAMAGUCHI¹ (1. National Institute of Polar Research)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R5-P-04] Internal Structure Analysis of Carbonaceous Material in Yamato 8448 Ureilite

*Masahiro YASUTAKE¹, Megumi Matsumoto², Akira Tsuchiyama^{3,4}, Kentaro Uesugi¹, Akihisa Takeuchi¹, Akira Yamaguchi⁵ (1. JASRI, 2. Tohoku Univ., 3. Ritsumeikan Univ., 4. CAS/GIG, 5. NIPR)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R5-P-05] Effects of initial porosity on shock deformation microstructures in rutile

*Yuhei Umeda^{1,2}, Yuma Nagai^{1,2}, Naotaka Tomioka³, Toshimori Sekine⁴, Masashi Miyakawa⁵, Takamichi Kobayashi⁵, Hitoshi Yusa⁵, Takuo Okuchi^{1,2} (1. KURNS, 2. Kyoto Univ. Eng., 3. JAMSTEC, 4. HPSTAR, 5. NIMS)

Dehydration of Newberyite($\text{MgHPO}_4 \cdot 3\text{H}_2\text{O}$) under low pressures

*Yukako Matsumoto¹, Shogo Tachibana¹

1. UTokyo

Keywords: Bennu, phosphate, dehydration, asteroid, Newberyite

Experimental simulation of oxygen isotope exchange between chondrule melt and low-pressure water vapor

*Daiki YAMAMOTO¹, Noriyuki Kawasaki²

1. Kyushu University, 2. Hokkaido University

Keywords: chondrule melt, low-pressure water vapor, oxygen isotope exchange, kinetics, protosolar disk

Early impact events recorded in anomalous eucrites

*Akira YAMAGUCHI¹

1. National Institute of Polar Research

Keywords: meteorites, achondrites, thermal history, impact

Internal Structure Analysis of Carbonaceous Material in Yamato 8448 Ureilite

*Masahiro YASUTAKE¹, Megumi Matsumoto², Akira Tsuchiyama^{3,4}, Kentaro Uesugi¹, Akihisa Takeuchi¹, Akira Yamaguchi⁵

1. JASRI, 2. Tohoku Univ., 3. Ritsumeikan Univ., 4. CAS/GIG, 5. NIPR

Keywords: Ureilite, Diamond, SR-XCT




図1. 炭素質物質のCT断面像。X線エネルギーは7.35keV。左下に表示コントラストを示す。(a) 黒色部からサンプリングした試料の断面像。60度120度で交わる線構造が確認できる。図中白い粒は鉄に富む包有物である。図中右端にはダイヤモンドを区切る鉄に富む膜が分布している。(b) 境界部からサンプリングした試料の断面像。図中右側にグラファイトから成る茶色部が、左側にダイヤモンドを主体とする黒色部が分布する。図中中央に分布する白色部は鉄に富む層である。茶色部は黒色部と比較し鉄に富む物質に乏しい事が分かる。赤矢印は虫食い状になったダイヤモンドを示す。図中略語：D:ダイヤモンド、Gr:グラファイト、Pt:プラチナ保護膜、Fe-film:鉄に富む膜、Resin:樹脂。

Effects of initial porosity on shock deformation microstructures in rutile

*Yuhei Umeda^{1,2}, Yuma Nagai^{1,2}, Naotaka Tomioka³, Toshimori Sekine⁴, Masashi Miyakawa⁵, Takamichi Kobayashi⁵, Hitoshi Yusa⁵, Takuo Okuchi^{1,2}

1. KURNS, 2. Kyoto Univ. Eng., 3. JAMSTEC, 4. HPSTAR, 5. NIMS

Keywords: Rutile, Shock deformation microstructures, Porosity

Poster presentation | R6: Plutonic rocks, volcanic rocks and subduction factory

📅 Sat. Sep 14, 2024 12:30 PM - 2:00 PM JST | Sat. Sep 14, 2024 3:30 AM - 5:00 AM UTC | 🏢 Entrance
Hall Higashiyama Campus

R6: Plutonic rocks, volcanic rocks and subduction factory

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R6-P-01] Orthopyroxene cumulates from a serpentinite mud-volcano in the mariana forearc: constituent rocks of the forearc Moho transition zone

*Pia Ataka¹, Yuji Ichiyama¹, Shun Takamizawa¹, Akihiro Tamura², Tomoaki Motishita² (1. Chiba University, 2. Kanazawa University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R6-P-02] Formation processes of high-Mg andesite associated with ophiolitic rocks in the Setogawa belt

*Tsumugi Kato¹, Yuji Ichiyama¹, Akihiro Tamura², Tomoaki Morishita² (1. Chiba Univ., 2. Kanazawa Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R6-P-03] Petrological and geochemical study of volcanics from Kuju volcanic group, Kyushu Island, Southwest Japan Arc.

「発表賞エントリー」

*Soma Yamanaka¹, Tomoyuki Shibata¹, Ryotaro Fujihara¹, Masako Yoshikawa¹, Tomo Shibata² (1. Hiroshima Univ. Sci. & Tech, 2. Fukuoka Univ. Sci.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R6-P-04] Petrological study of intrusive rocks and their crustal xenoliths in the Suzu area, Atsumi district, Yamagata

*Kazuhiro Tojo¹, Takashi Hoshide¹ (1. Akita Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R6-P-05] Crystallization conditions of zircon and its crystallization relationship with surrounding minerals in the Tono plutonic complex, Kitakami mountains

*Yasuhiro OGITA^{1,2}, Shuhei Sakata³, Takeshi Ohno⁴, Tatsunori Yokoyama², Satoshi Suzuki¹, Kyoka Endo¹, Takashi Yuguchi⁵ (1. Yamagata Univ., 2. JAEA, Tono Geoscience Center, 3. The Univ. Tokyo, Earthquake Research Inst., 4. Gakushuin Univ., 5. Kumamoto Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R6-P-06] Petrography and magmatic processes of a diorite body distributed in the Kawakami region, Yamaguchi Prefecture, Japan

「発表賞エントリー」

*Ryosei Sugiura¹, Keisuke Eshima¹ (1. Yamaguchi University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R6-P-07] Sr–Nd–Hf isotopic systematics for the Habu granodiorite, eastern part of Yamaguchi Prefecture

*Masaaki OWADA¹, Yusaku Akimoto², Mariko Nagashima¹, Keisuke Eshima¹, Atsushi Kamei³, Nobuhiko Nakano⁴, Kenta Kawaguchi⁵, Yasuhito Osanai⁴ (1. Yamaguchi Univ., 2. YON-C, 3. Shimane Uni., 4. Kyushu Univ., 5. Hiroshima Univ.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R6-P-08] Rare earth element compositions and U–Pb zircon age of granodiorite porphyrites in the Noko Island, western part of northern Kyushu

*Masaki YUHARA¹, Yoshinobu KAWANO², Kazuhiro TSUKADA³, Purevdulam SUKHBAATAR⁴ (1. Fukuoka Univ. Sci., 2. Rissyo Univ. Env. Sci., 3. Nagoya Univ. Mus., 4. Nagoya Univ. Env. St.)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R6-P-09] Compositional changes and growth processes of the Asakura granodiorite, Northern Kyushu, Japan.

「発表賞エントリー」

*Yudai HAMANO¹, Keisuke ESHIMA¹ (1. Yamaguchi Uni. Sci)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R6-P-10] Intrusive relationship and lithofacies classification of Granite–Granodiorite in Eastern Fukuoka Prefecture

「発表賞エントリー」

*Ryo Nakamura¹, Keisuke Eshima¹ (1. Yamaguchi University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R6-P-11] Crystallization processes of quartz in a granitic magma: Implications for the magma chamber processes of Okueyama granite, Kyushu, Japan

*Takashi YUGUCHI¹, Takenori Kato⁴, Yasuhiro Ogita², Minori Watanabe⁶, Asuka Kato⁶, Daichi Itoh⁶, Tatsunori Yokoyama², Shuhei Sakata³, Takeshi Ohno⁵ (1. Kumamoto University, 2. JAEA, 3. Uni. Tokyo ERI, 4. Nagoya University, 5. Gakushuin University, 6. Yamagata University)

12:30 PM - 2:00 PM JST | 3:30 AM - 5:00 AM UTC

[R6-P-13] Differentiation Processes of Granitic Magma Chamber Recorded in 'Oshima-ishi' Distributed in Oshima, Geiyo islands, Ehime Prefecture

*Kazuya SHIMOOKA^{1,2}, Soichiro AONO², Takahito ONISHI², Toko FUKUI², Motohiro TSUBOI¹, Toshiro TAKAHASHI³, Satoshi SAITO² (1. Kwansai Gakuin Univ., 2. Ehime Univ., 3. Niigata Univ.)

Orthopyroxene cumulates from a serpentinite mud-volcano in the Mariana forearc: constituent rocks of the forearc Moho transition zone

*Pia Ataka¹, Yuji Ichiyama¹, Shun Takamizawa¹, Akihiro Tamura², Tomoaki Motishita²

1. Chiba University, 2. Kanazawa University


Keywords: Serpentine, Ophiolite, Mariana Forearc, Moho transition zone, cumulate

Formation processes of high-Mg andesite associated with ophiolitic rocks in the Setogawa belt

*Tsumugi Kato¹, Yuji Ichiyama¹, Akihiro Tamura², Tomoaki Morishita²

1. Chiba Univ., 2. Kanazawa Univ.

Keywords: high-Mg andesite, Setogawa belt, felsic crystalline inclusions, magma mixing



Petrological and geochemical study of volcanics from Kuju volcanic group, Kyushu Island, Southwest Japan Arc.

*Soma Yamanaka¹, Tomoyuki Shibata¹, Ryotaro Fujihara¹, Masako Yoshikawa¹, Tomo Shibata²

1. Hiroshima Univ. Sci. & Tech, 2. Fukuoka Univ. Sci.

Keywords: modal composition, Kuju, arc magma




図 九重火山群火山岩類のモード組成。

Petrological study of intrusive rocks and their crustal xenoliths in the Suzu area, Atsumi district, Yamagata

*Kazuhiro Tojo¹, Takashi Hoshide¹

1. Akita Univ.

Keywords: Petrology, Xenoliths

Crystallization conditions of zircon and its crystallization relationship with surrounding minerals in the Tono plutonic complex, Kitakami mountains

*Yasuhiro OGITA^{1,2}, Shuhei Sakata³, Takeshi Ohno⁴, Tatsunori Yokoyama², Satoshi Suzuki¹, Kyoka Endo¹, Takashi Yuguchi⁵

1. Yamagata Univ., 2. JAEA, Tono Geoscience Center, 3. The Univ. Tokyo, Earthquake Research Inst., 4. Gakushuin Univ., 5. Kumamoto Univ.

No abstract in English.

Keywords: Zircon, U-Pb age, Ti concentration, Tono Plutonic Complex

表1 遠野岩体のジルコンの産状と分析数

岩相	サンプル名	分析点数	ジルコン粒子数				
			in Qtz	in Bt	in Afs	粒間	
中心相	025	23	4	1	1	0	2
	029	3	1	1	0	0	0
	102	11	4	1	0	1	2
主岩相	021	5	4	1	1	0	2
	404	6	3	0	0	0	3
	405	31	11	0	3	0	8

in Qtz: 石英に包有されるジルコン, in Bt: 黒雲母に包有されるジルコン,
in Afs: アルカリ長石に包有されるジルコン, 粒間: 粒間に産出するジルコン

Petrography and magmatic processes of a diorite body distributed in the Kawakami region, Yamaguchi Prefecture, Japan

*Ryosei Sugiura¹, Keisuke Eshima¹

1. Yamaguchi University

Keywords: Grandiorite, Volcanic-plutonic complex, Kwanmon Group, Shunan Group

Sr–Nd–Hf isotopic systematics for the Habu granodiorite, eastern part of Yamaguchi Prefecture

*Masaaki OWADA¹, Yusaku Akimoto², Mariko Nagashima¹, Keisuke Eshima¹, Atsushi Kamei³, Nobuhiko Nakano⁴, Kenta Kawaguchi⁵, Yasuhito Osanai⁴

1. Yamaguchi Univ., 2. YON-C, 3. Shimane Uni., 4. Kyushu Univ., 5. Hiroshima Univ.

We performed geological, petrographical, and geochemical investigations including Sr–Nd–Hf isotopic studies for the Habu granodiorite located in the eastern part of Yamaguchi Prefecture, and discussed its magmatic processes and source materials. The laccolithic Habu granodiorite consists mainly of the leucocratic biotite granodiorite (leucocratic facies) surrounded by the thin-skinned hornblende-biotite granodiorite (melanocratic facies). Zircon U–Pb dating gives ages of c. 105 Ma for both lithofacies. Although the whole-rock chemical data of both lithofacies make monotonous trends on some binary plots, the leucocratic facies are slightly affected by crustal contaminations, as evidenced by Sr–Nd isotopic features. Epsilon Hf isotopic model ages revealed that the Habu granodiorite would be derived from the mafic lower crust probably separated from the depleted mantle during the Mesoproterozoic time.

Keywords: Habu granodiorite, Magmatic processes, Source materials, Multi-isotopic systematics, Igneous activities Southwest Japan

Rare earth element compositions and U-Pb zircon age of granodiorite porphyrites in the Noko Island, western part of northern Kyushu

*Masaki YUHARA¹, Yoshinobu KAWANO², Kazuhiro TSUKADA³, Purevdulam SUKHBAATAR⁴

1. Fukuoka Univ. Sci., 2. Rissho Univ. Env. Sci., 3. Nagoya Univ. Mus., 4. Nagoya Univ. Env. St.

Keywords: granodiorite porphyrite, Noko Island, Sr and Nd isotopic composition, Rare earth element compositions, U-Pb zircon age

Compositional changes and growth processes of the Asakura granodiorite, Northern Kyushu, Japan.

*Yudai HAMANO¹, Keisuke ESHIMA¹

1. Yamaguchi Uni. Sci

Keywords: Cretaceous, Northern Kyushu batholith, Granodiorite, Lithofacies change, Growth processes

Intrusive relationship and lithofacies classification of Granite–Granodiorite in Eastern Fukuoka Prefecture

*Ryo Nakamura¹, Keisuke Eshima¹

1. Yamaguchi University

Keywords: plutonic rock, Cretaceous , plutonic complex, intrusive relationship, magma mixing

Crystallization processes of quartz in a granitic magma: Implications for the magma chamber processes of Okueyama granite, Kyushu, Japan

*Takashi YUGUCHI¹, Takenori Kato⁴, Yasuhiro Ogita², Minori Watanabe⁶, Asuka Kato⁶, Daichi Itoh⁶, Tatsunori Yokoyama², Shuhei Sakata³, Takeshi Ohno⁵

1. Kumamoto University, 2. JAEA, 3. Uni. Tokyo ERI, 4. Nagoya University, 5. Gakushuin University, 6. Yamagata University

Keywords: Quartz, Cathodoluminescence, TitaniQ thermometer, Okueyama granite, Ti-in-zircon thermometer

Differentiation Processes of Granitic Magma Chamber Recorded in 'Oshima-ishi' Distributed in Oshima, Geiyo islands, Ehime Prefecture

*Kazuya SHIMOOKA^{1,2}, Soichiro AONO², Takahito ONISHI², Toko FUKUI², Motohiro TSUBOI¹, Toshiro TAKAHASHI³, Satoshi SAITO²

1. Kwansei Gakuin Univ., 2. Ehime Univ., 3. Niigata Univ.

Keywords: Oshima-ishi, Cretaceous granitoids, Differentiation Processes, Mafic magma, Oshima, Geiyo islands

"Mineralogy 2022"General Outreach lecture

📅 Sat. Sep 14, 2024 3:30 PM - 5:00 PM JST | Sat. Sep 14, 2024 6:30 AM - 8:00 AM UTC | 🏢 ES Hall
Higashiyama Campus

General Outreach lecture

3:30 PM - 5:00 PM JST | 6:30 AM - 8:00 AM UTC

[3Lecture-1-1530-1add] 一般普及講演

"Mineralogy 2022" General Outreach lecture

General Outreach lecture

Sat. Sep 14, 2024 3:30 PM - 5:00 PM ES Hall (Higashiyama Campus)

3:30 PM - 5:00 PM

[3Lecture-1-1530-1add]一般普及講演