
インターナショナルセッション（口頭発表） | セッション記号 A（大気海洋・環境科学） | A-AS 大気科学・気象学・大気環境

[A-AS01_30PM1] Extreme Weather in Cities

コンビーナ:*真木 雅之(鹿児島大学地域防災教育研究センター)、松本 淳(首都大学東京大学院都市環境科学研究科地理環境科学専攻)、小司 禎教(気象研究所気象衛星・観測システム研究部第2研究室)、中谷 剛(独立行政法人防災科学技術研究所)、座長:真木 雅之(鹿児島大学地域防災教育研究センター)

2014年4月30日(水) 14:15 ~ 15:45 423 (4F)

It is recognized that large cities with a population of several million people are inherently vulnerable to extreme weathers such as torrential rain, lightning, strong wind, giant typhoon, and heat wave. It is argued that the occurrence of extreme weather phenomena tends to increase due to the climatic change. Cooperating with domestic and international academic scientists, the session will focus on the mechanism of extreme weather, its monitoring and prediction methods, effects of urbanization on hazards, and social experiments on resilient cities.

15:00 ~ 15:15

[AAS01-P03_PG]An Ensemble Nowcasting of Rainfall over the Kanto Region, Japan

ポスター講演3分口頭発表枠

*P.C. Shakti¹、MISUMI Ryohei¹、NAKATANI Tsuyoshi¹、MAKI Masayuki²、SEED Alan³ (1.National Research Institute for Earth Science and Disaster Prevention(NIED), Tsukuba, Japan、2.Kagoshima University, Kagoshima, Japan、3.Bureau of Meteorology, Melbourne, Australia)

キーワード : nowcasting, ensemble, weather radar, extreme rainfall, STEPS, hydrological model

Every year weather-related disasters: extreme rainfall, landslides and flooding destroy livelihoods and damage economics somewhere on the planet. Recently, number of flash flooding is believed to be increasing specially in urban areas. It has being a great challenge to forecast flood warning and urban drainage management. Nowcasting of rainfall (very short-range forecasting) is an important tool to minimize or manage all these weather-related disasters since precipitation is the main input. Common practice to forecast heavy precipitation for hydrological application varies from 0-6 hr and there are different kinds of nowcasting based on different method.

Nowcasting of rainfall comprises the detailed description of the current weather along with forecasts obtained by extrapolation for a different time period ahead. In this study, we focus on ensemble nowcasting of rainfall. It refers to the fact that many forecasts are produced, with the rainfall areas moving at slightly different speeds, and with the small rainfall features represented by slightly different random statistics. By comparing these different nowcasting of rainfall, the forecaster can decide how likely a particular weather event will be. It gives a much better idea of what weather events may occur at a particular time. Short Term Ensemble Prediction System (STEPS), one of the most advanced Quantitative Precipitation Forecast (QPF) systems currently available is considered for nowcasting of rainfall. Japan Meteorological Agency (JMA) and X-band multi-parameter (MP) radar data were considered to produce an ensemble nowcasting of rainfall. First, JMA radar rainfall data of Kanto region was fixed to check the performance of STEPS. Skill scores showed that STEPS can give a good forecast for less than one hour. However, more uncertainties can be seen during the starting and ending of rain event. High resolution of data (MP data) also used in the STEPS under the default condition. Overall, an ensemble nowcasting of rainfall seems close with real time data, which could be interesting to use them in hydrological model.