

[EE] Eveningポスター発表 | セッション記号 P (宇宙惑星科学) | P-PS 惑星科学

[P-PS01] Outer Solar System Exploration Today, and Tomorrow

コンビーナ:木村 淳(大阪大学)、笠羽 康正(東北大学大学院 理学研究科 地球物理学専攻)、Vance Steven(Jet Propulsion Laboratory, Caltech、共同)、Kunio M. Sayanagi (Hampton University)

2018年5月21日(月) 17:15 ~ 18:30 ポスター会場 (幕張メッセ国際展示場 7ホール)

The giant planets provide many keys to understanding planetary processes. They play an important role in shaping our solar system, and the physical and chemical processes they harbor also provide a unique opportunity to study the phenomena relevant for studying

Earth and other planets, including exoplanetary systems. In this session, we discuss a wide range of topics encompassing the giant planets and their moons, including their origins, interiors, atmospheres, compositions, surface features, and electromagnetic fields. To advocate for current and future outer planets exploration (Cassini, Juno, New Horizons, JUICE, and beyond), we also call for discussions on future missions to explore giant planet systems, including how to develop better international cooperation. Discussion in this latter category will include progress in developing a solar sail mission concept for observing the Jupiter system and its trojan asteroids.

[PPS01-P12] Enhancement of the Jovian Magnetospheric Plasma Circulation Caused by the Change in Plasma Supply From the Satellite Io

*土屋 史紀¹、吉岡 和夫²、木村 智樹³、古賀 亮一¹、村上 豪⁴、山崎 敦⁴、埜 千尋⁵、鍵谷 将人¹、鈴木 文晴²、疋田 伶奈²、吉川 一朗²、笠羽 康正⁶、北 元⁶、三澤 浩昭¹、坂野井 健¹ (1.東北大学大学院理学研究科惑星プラズマ・大気研究センター、2.東京大学新領域創成科学研究科、3.理化学研究所、4.宇宙科学研究所、5.情報通信研究機構、6.東北大学大学院理学研究科)

キーワード：木星、イオ、磁気圏

The innermost Galilean satellite, Io, supplies a large amount of volcanic gases to the Jovian magnetosphere, and the outward transport of ionized gasses is responsible for forming a huge and rotation-dominant magnetosphere. The plasma supply from the satellite has a key role in the characterization of the Jovian magnetosphere. In fact, significant variations of the plasma population in the inner magnetosphere caused by the volcanic eruptions in Io were found in the early 2015, using a continuous data set of the Io plasma torus obtained from an extreme ultraviolet spectroscopy onboard the HISAKI satellite. From the time evolution of the Io plasma torus radial distribution, it was found that enhanced plasma supply from the inner to the middle magnetospheres occurred during approximately 50 days and the average outward speed is estimated to be 140 m/s. Intense short-lived auroral brightenings—which represent transient energy releases in the outer part of the magnetosphere—occurred frequently during this period. The short-lived auroral brightenings accompanied well-defined sporadic enhancements of the ion brightness in the plasma torus, indicating a rapid inward transport of energy from the outer part of the magnetosphere and the resultant enhancement of hot electron population in the inner magnetosphere. This is the first observational evidence showing that the strength of a plasma source in the inner magnetosphere can control a large-scale radial circulation of mass and energy in a rotation-dominant magnetosphere.