

Flux decrease of outer radiation belt electrons associated with solar wind pressure pulse: A Code coupling simulation of GEMSIS-RB and GEMSIS-GM

*伊藤 大輝¹、三好 由純¹、齊藤 慎司⁴、松本 洋介²、天野 孝伸³、林 昌広¹

*Ito Hiroki¹, Yoshizumi Miyoshi¹, Shinji Saito⁴, Yosuke Matsumoto², Takanobu Amano³, Masahiro Hayashi¹

1. 名古屋大学宇宙地球環境研究所、2. 千葉大学大学院理学研究科、3. 東京大学 地球惑星科学専攻、4. 名古屋大学 大学院理学研究科

1. Institute for Space-Earth Environmental Research, Nagoya University, 2. Graduate School of Science, Chiba University, 3. Department of Earth and Planetary Science, University of Tokyo, 4. Graduate School of Science, Nagoya University

Relativistic electron flux of the outer radiation belt dynamically changes in response to solar wind variations. There exist several conditions to cause the flux drop-out of outer belt electrons. The magnetopause shadowing (MPS) is one of the process to cause the rapid loss of outer belt electrons (e.g., Kim et al., 2008). In this study, we performed a code-coupling simulation using GEMSIS-RB test particle simulation code (Saito et al., 2010) and GEMSIS-GM global MHD magnetosphere simulation code (Matsumoto et al., 2010) to specify which electrons of the outer belt will be lost due to the MPS, by investigating pitch angle and MLT dependences. We calculated the guiding center equation of each particle with initial L-shells 7.5 Re to 9.0 Re, energies 300 keV to 3 MeV and pitch angles 60 degrees to 120 degrees. The initial dynamic pressure of the solar wind is 1.0 nPa. The standoff distance of magnetopause at the subsolar point was 12 Re as an initial state. The magnetopause moved to 9 Re due to the dynamic pressure enhancement when the solar wind pressure became 2.5 nPa. After the enhancement of the dynamic pressure for 40 sec, the L-values of the magnetopause moved back to 10 Re due to the dynamic pressure decrease. We found that electron flux decreased during the period, which strongly depended on MLT and pitch angle of electrons. The simulation showed that more than 80% of lost electrons lost from the magnetopause at 8 ~ 11 MLT. The pitch angles of 90% of lost electrons are 85 ~ 95 degrees. Because the MPS process causes loss of electrons at L values higher than 6.75, it is suggested that other loss processes are necessary for the loss at lower L-values. After the MPS, the electron fluxes at lower L-shells decreased through outward motion of trapped electrons due to $E \times B$ drift. Our simulation results suggest that the dominant loss processes of electron flux in the MPS at $L > 6.75$, while the outward motion at $L < 6.75$.

キーワード：マグネットポーズシャドウィング

Keywords: magnetopause shadowing