

Computer simulations of electron scatterings through chorus-wave particle interactions

伊藤 義起¹、*三好 由純¹、齊藤 慎司²

Yoshiki Ito¹, *Yoshizumi Miyoshi¹, Shinji Saito²

1. 名古屋大学宇宙地球環境研究所、2. 情報通信研究機構

1. Institute for Space-Earth Environmental Research, Nagoya University, 2. NICT

Whistler mode chorus waves cause scattering and acceleration of energetic electrons in the inner magnetosphere, and recent studies identified that chorus waves cause the pulsating aurora. The interaction processes have been modeled as diffusions in the velocity space, and the scattering rate increases with increasing the wave amplitude. However, actual wave-particle interactions with chorus waves are non-linear process, so that it is expected that the scattering rate does not have a simple correlation with wave amplitude. In this study, we conduct a computer simulation to investigate electron scattering process through chorus-wave particle interactions with the GEMSIS-RBW code. The GEMSIS-RBW simulation calculates variations of local pitch angle and energy of each test particle by the imposed chorus waves by calculating the Maxwell equations.

We calculate the trajectory of a number of electrons with various energy and pitch angles. We analyzed the electron motion in the phase space and try to discriminate the motion with the parameter ρ [Bortnick et al., 2008] that is a proxy of the ratio of the wave-induced and the background inhomogeneity effects for the momentum change of the resonant electron. We identified that the phase trapping effect decreases the precipitating flux and contribute to generation of the butterfly type pitch angle distributions. Moreover, we identified diffusion and dislocation contribute to the precipitations into the atmosphere. Multiple resonance changes the ρ of electrons from the phase trapping to dislocation, which causes rapid precipitations of electrons.

キーワード：波動粒子相互作用、コンピューターシミュレーション

Keywords: wave-particle interactions, Computer simulation