

小惑星リュウグウとベヌーの外来物質から示唆される衝突史

Impact histories inferred from exogenic boulders on asteroids Ryugu and Bennu

*巽 瑛理^{1,2}、杉本 知穂²、杉田 精司²、Popescu Marcel⁴、Campins Humberto¹¹、de León Julia³、Amy Simon⁵、Kaplan Hannah⁶、長 勇一郎²、諸田 智克²、本田 理恵⁷、亀田 真吾⁸、横田 康弘³、湯本 航生²、青木 美波²、DellaGiustina Daniella¹⁰、Golish Dathon¹⁰、Licandro Javier¹、Rizos Juan Luis¹、廣井 孝弘¹²、Domingue Deborah¹³、Michel Patrick¹⁴、Lauretta Dante¹⁰、山田 学¹⁵、坂谷 尚哉⁸、神山 徹⁹、本田 親寿¹⁶、早川 雅彦³、松岡 萌³、鈴木 秀彦¹⁷、小川 和律³、澤田 弘崇³、*ERI TATSUMI^{1,2}、Chiho Sugimoto²、Seiji Sugita²、Marcel Popescu⁴、Humberto Campins¹¹、Julia de León³、Amy A. Simon⁵、Hannah H. Kaplan⁶、Yuichiro Cho²、Tomokatsu Morota²、Rie Honda⁷、Shingo Kameda⁸、Yasuhiro Yokota³、Koki Yumoto²、Minami Aoki²、Daniella N. DellaGiustina¹⁰、Dathon R. Golish¹⁰、Javier Licandro¹、Juan Luis Rizos¹、Takahiro Hiroi¹²、Deborah Domingue¹³、Michel Patrick¹⁴、Dante S. Lauretta¹⁰、Manabu Yamada¹⁵、Naoya Sakatani⁸、Toru Kouyama⁹、Chikatoshi Honda¹⁶、Masahiko Hayakawa³、Moe Matsuoka³、Hidehiko Suzuki¹⁷、Kazunori Ogawa³、Hirotaka Sawada³

1. カナリア天文物理学研究所、2. 東京大学、3. 宇宙科学研究所、4. ルーマニア天文研究所、5. NASAゴダード宇宙飛行センター、6. サウスウエスト研究所、7. 高知大学、8. 立教大学、9. 産業総合研究所、10. アリゾナ大学、11. セントラルフロリダ大学、12. ブラウン大学、13. 惑星科学研究所、14. コートダジュール天文台、15. 千葉工業大学、16. 会津大学、17. 明治大学

1. Instituto de Astrofísica de Canarias, 2. Univ. of Tokyo, 3. ISAS/JAXA, 4. Astronomical Institute of the Romanian Academy, 5. NASA Goddard Space Flight Center, 6. Southwest Research Institute, 7. Kochi Univ., 8. Rikkyo Univ., 9. AIST, 10. Univ. of Arizona, 11. Univ. of Central Florida, 12. Brown Univ., 13. Planetary Science Institute, 14. Observatoire de la Côte d'Azur, 15. Chiba Institute of Tech., 16. Univ. of Aizu, 17. Meiji Uni.

Rubble-pile asteroids, such as (162173) Ryugu and (101955) Bennu, are formed as a result of catastrophic disruption of a parent body and re-accumulation of the fragments by self-gravity (Michel and Ballouz et al. 2020). Therefore, reaccumulated rubble piles could include mixtures of materials from both the parent body and its catastrophic impactor, as they did in the case of 2008 TC₃ and the Almahata Sitta meteorites (Jenniskens et al. 2009). Indeed, six unusually bright, basaltic, meter-scale boulders were recently identified on Bennu's dark surface (DellaGiustina et al. 2021), and their close spectral matches to the Howardite-Eucrite-Diogenite (HED) meteorites and Vesta family members indicate that they originated from asteroid (4) Vesta. In parallel, bright exogenic anhydrous-silicate-rich materials were found on Ryugu (Tatsumi et al. 2021). The bright boulders on Ryugu are consistent with ordinary chondrite meteorites, based on their albedo and the weak or even absent absorption band at 2 μm. These exogenous materials on the surfaces of rubble-pile asteroids could be a key to constraining their specific impact conditions and collisional evolution.

In this study, we report further study of the visible spectrophotometry and morphology of exogenic boulders on Ryugu and Bennu using two multi-band cameras: the telescopic Optical Navigation Camera (ONC-T) onboard the Hayabusa2 spacecraft, and MapCam onboard the Origins, Spectral Interpretation, Resource Identification, and Security-Regolith Explorer (OSIRIS-REx) spacecraft, respectively.

For Ryugu, we first detected objects with peak reflectance brighter by a factor of >1.5 than the surrounding area. Tatsumi et al. (2021) found two spectral groups among the bright boulders: S-type with an absorption near $1\ \mu\text{m}$ and C/X-type with a flat spectrum. That study found 6 S-types and 15 C/X-types, and here we report the additional finding of two S-types and ~ 70 C/X-types. We found that S-type bright boulders follow two parallel trends consistent with space weathering of ordinary chondrites. Fragments from projectiles suggest that Ryugu may not have formed directly from the original parent body, i.e., it is from a second or higher generation of collisional descendants (Sugimoto et al. in revision).

For Bennu, we directly surveyed the absorption near $1\ \mu\text{m}$, which is indicative of anhydrous silicates. We propose 77 objects of exogenic origin including 6 boulders previously reported by DellaGiustina et al. (2021). We find that proposed exogenic objects follow two mixing trends with respect to the average spectrum of Bennu (Tatsumi and Popescu et al. submitted). One trend can be explained by mixing with HEDs, as discussed in DellaGiustina et al. (2021), and the other trend can be explained by mixing with another, of olivine-rich composition. Near-infrared spectra also support the possibility of multiple compositions among exogenic objects on Bennu. Le Corre et al. (submitted) also reached to the similar conclusion with a different methodology. Thus, similar to Ryugu, Bennu could represent a second or higher generation collisional fragment.

Using high-resolution images, we found that more than half of the exogenic objects that we identified on Bennu have breccia-like or inclusion-like morphologies. We do not observe a correlation between morphology and spectral shape.

Direct comparison of reflectance spectra of exogenic materials between Ryugu and Bennu suggests a compositional difference. Furthermore, the number density of exogenic objects $> 0.5\ \text{m}$ on Bennu is ~ 40 times that on Ryugu (Tatsumi and Popescu et al. submitted). Bennu exhibits much more contamination from exogenic materials than Ryugu, which likely reflects differences in impact conditions, such as impact velocity and impactor size.

Michel and Ballouz et al. (2020) Nat. Commun. 11, 2655

Jenniskins et al. (2009) Nature 458, 485-488.

DellaGiustina et al. (2021) Nat. Astron. 5, 31-38.

Tatsumi et al. (2021) Nat. Astron. 5, 39-45.

Sugimoto et al. (in revision) Icarus.

Tatsumi and Popescu et al. (submitted) MNRAS.

Le Corre et al. (submitted) PSJ.

キーワード：小惑星、はやぶさ2、オシリスレックス、リュウグウ、ベヌー

Keywords: Asteroids, Hayabusa2, OSIRIS-REx, Ryugu, Bennu