

Investigation of the formation process of the source plasma driving SAPS fine structures found by Arase and SuperDARN

深見 岳弘¹、*熊本 篤志¹、加藤 雄人¹、西谷 望²、堀 智昭²、笠羽 康正¹、土屋 史紀¹、寺本 万里子³、木村 智樹⁴、笠原 禎也⁵、小路 真史²、中村 紗都子²、北原 理弘²、松岡 彩子⁶、今城 峻⁶、笠原 慧⁷、横田 勝一郎⁸、桂華 邦裕⁷、風間 洋一⁹、Wang Shiang-Yu⁹、Jun Chae-Woo²、浅村 和史¹⁰、三好 由純²、篠原 育¹⁰、Shepherd Simon¹¹

Takehiro Fukami¹, *Atsushi Kumamoto¹, Yuto Katoh¹, Nozomu Nishitani², Tomoaki Hori², Yasumasa Kasaba¹, Fuminori Tsuchiya¹, Mariko Teramoto³, Tomoki Kimura⁴, Yoshiya Kasahara⁵, Masafumi Shoji², Satoko Nakamura², Masahiro Kitahara², Ayako Matsuoka⁶, Shun Imajo⁶, Satoshi Kasahara⁷, Shoichiro Yokota⁸, Kunihiro Keika⁷, Yoichi Kazama⁹, Shiang-Yu Wang⁹, Chae-Woo Jun², Kazushi Asamura¹⁰, Yoshizumi Miyoshi², Iku Shinohara¹⁰, Simon G. Shepherd¹¹

1. 東北大学大学院理学研究科、2. 名古屋大学宇宙地球環境研究所、3. 九州工業大学、4. 東京理科大学、5. 金沢大学、6. 京都大学、7. 東京大学、8. 大阪大学、9. 中央研究院天文及天文物理研究所、10. JAXA ISAS、11. Dartmouth College
1. Graduate School of Science, Tohoku University, 2. Institute for Space-Earth Environmental Research, Nagoya University, 3. Kyushu Institute of Technology, 4. Tokyo University of Science, 5. Kanazawa University, 6. Kyoto University, 7. The University of Tokyo, 8. Osaka University, 9. Institute of Astronomy and Astrophysics, Academia Sinica, 10. JAXA ISAS, 11. Dartmouth College

Contribution of magnetospheric pressure inhomogeneities to subauroral polarization stream fine structures (SAPS-FS) is investigated on the basis of an event analysis of conjugated Arase satellite and SuperDARN observations.

SAPS is a strong westward flow in the subauroral region and an important magnetosphere-ionosphere coupling signature [Foster and Burke, 2002]. Previous studies reported various fine-scale and short-time variations within SAPS [e.g. Erickson et al., 2002; Mishin et al., 2003; Oksavik et al., 2006; Makarevich and Bristow, 2014]. There still have been many open questions for SAPS-FS, such as which spatial or temporal variations they are, and what kind of physical processes and source plasma in the magnetosphere generate SAPS-FS as observed in the ionosphere.

In order to address those questions, we analyzed SuperDARN-Arase conjugated observations on July 9th, 2017, which was carried out as a special campaign observation. SuperDARN Christmas Valley East radar observed the two-dimensional structures of SAPS-FS in the ionosphere. The fine structures moved equatorward and simultaneously extended westward. They merged and formed a latitudinally-broad SAPS 10-20 minutes after they started to be observed by SuperDARN. On the other hand, Arase traveled radially outward on the duskside in the inner magnetosphere and encountered four positive peaks of the outward electric field. Upon each passage of three of the four peaks of electric field, the satellite also detected eastward perturbation in the magnetic field. In addition, the three peaks were accompanied by enhancement of the hot ion pressure with a negative peak of the magnetic field intensity, which can be interpreted in terms of the diamagnetic effect. The first and second pressure peaks were formed by rapid enhancements of proton flux with small energy-time dispersion. It is suggested that the earthward and anti-earthward field aligned currents (FACs) were respectively generated at the earthside and anti-earthside edges of the ion pressure lumps. Since we could confirm that SAPS-FS in the event formed spatial structures on the basis of the continuous and two-dimensional SuperDARN observation, we can expect SAPS-FS sources also forms spatial structures in the inner magnetosphere. We examined the correspondence between SAPS-FS and the magnetospheric electric fields based on T04s magnetic field model [Tsyganenko and Sitnov, 2005]. While the one-to-one correspondence could not be identified, it was supported that the corresponding uniform structures were simultaneously formed in the ionosphere

and magnetosphere in the event by coincidence of the estimations of the structure's size and speed in both regions. Our results support that the current generator theory can be applied not only to SAPS but also to SAPS-FS [e.g., Wang et al., 2019].

For this event, we performed a back-tracing of protons along their drift paths and also compared it with geosynchronous satellite data to discuss whether the particle injection into the inner magnetosphere could be the origin of SAPS-FS. The resultant drift calculation indicate that high-energy protons did not come from the nightside plasma sheet to Arase under simple convection electric field as expressed in the Volland-Stern model [Volland 1973; Maynard and Chen 1975]. If an azimuthally-localized electric field, modeling those of substorm injection, added in the night side, the high-energy protons accelerated earthward and reach Arase. On the basis of the geosynchronous satellite observations, a signature of the single injection was found by GOES-13 prior to the SAPS-FS observation. The travel time between from GOES-13 to Arase is roughly consistent with the typical earthward speed of substorm injection reported by previous studies. These results suggest that the hot plasma was injected multiple times into the inner magnetosphere and formed fine-scale, enhanced pressure regions [e.g., Gkioulidou et al., 2014] during this SAPS-FS event.

キーワード : Subauroral polarization streams (SAPS)、SAPS微細構造 (SAPS-FS)、あらせ探査機 (ERG)、SuperDARN、粒子インジェクション

Keywords: Subauroral polarization streams (SAPS), SAPS fine structures (SAPS-FS), Arase (ERG) spacecraft, SuperDARN, Particle injection