

Interannual variations in $\delta(\text{APO})$ and $\delta(\text{Ar}/\text{N}_2)$ observed at four Japanese stations for the period 2012-2022

*Shigeyuki Ishidoya¹, Yasunori Tohjima², Shohei Murayama¹, Yosuke Niwa², Kentaro Ishijima³, Satoshi Sugawara⁴, Tsuboi Kazuhiro³, Nobuyuki Aoki¹

1. National Institute of Advanced Industrial Science and Technology (AIST), 2. National Institute for Environmental Studies, 3. Meteorological Research Institute, 4. Miyagi University of Education

Atmospheric Potential Oxygen ($\text{APO} = \text{O}_2 + 1.1 \times \text{CO}_2$) ($\delta(\text{APO})$) varies due to air-sea O_2 , N_2 and CO_2 fluxes and fossil fuel O_2 and CO_2 fluxes, while atmospheric Ar/N_2 ratio ($\delta(\text{Ar}/\text{N}_2)$) varies due only to air-sea Ar and N_2 fluxes. It has been reported by past studies that seasonal and interannual variations in $\delta(\text{APO})$ are driven mainly by the air-sea O_2 and N_2 fluxes, although the air-sea CO_2 and fossil fuel fluxes cause a secular $\delta(\text{APO})$ trend (e.g. Tohjima et al., 2019; Ishidoya et al., 2021). As to the air-sea exchange, Ar and N_2 fluxes are driven by solubility change, and O_2 flux is driven by both solubility and biospheric changes. Therefore, it is expected that we can separate an interannual variation in $\delta(\text{APO})$ due to the solubility change ($\delta(\text{APO})_{\text{therm}}$) from that to the net marine biological activities ($\delta(\text{APO})_{\text{netbio}}$) by a combined analysis of $\delta(\text{APO})$ and $\delta(\text{Ar}/\text{N}_2)$. We have conducted simultaneous observations of $\delta(\text{APO})$ and $\delta(\text{Ar}/\text{N}_2)$ at various observational sites, and the data longer periods than 10 years have been obtained at Tsukuba (36°N, 140°E), Hateruma Island (24°N, 124°E), Cape Ochiishi (43°N, 146°E), and Takayama (36°N, 137°E), Japan (updated from Ishidoya et al., 2021). The annual change rate of the average $\delta(\text{APO})_{\text{therm}}$ at the four sites, obtained by multiplying a coefficient of 0.9 derived from differences in the solubility in O_2 and Ar (Weiss, 1970), was found to vary in phase with the Southern Oscillation Index (SOI) and the annual change rate of the global ocean heat content. On the other hand, the corresponding annual change rate of the average $\delta(\text{APO})_{\text{netbio}}$ obtained by subtracting the rate of $\delta(\text{APO})_{\text{therm}}$ from $\delta(\text{APO})$ (small contributions of the air-sea CO_2 and fossil fuel fluxes are also subtracted based on a simulation using an atmospheric transport model), varied in opposite phase with SOI. These responses of $\delta(\text{APO})_{\text{therm}}$ and $\delta(\text{APO})_{\text{netbio}}$ to El Niño / La Niña events are qualitatively consistent with those expected from the simulations based on a community earth system model by Eddebbbar et al. (2017).

Acknowledgements

We thank staff of Global Environmental Forum (GEF) for their works to collect the air samples at Hateruma and Ochiishi stations. This study was partly supported by the JSPS KAKENHI (grant nos. 22H05006, 19H01975 and 15H02814) and the Global Environment Research Coordination System from the Ministry of the Environment, Japan (grant nos. METI1454 and METI1953).

References

- Eddebbbar, Y. A., et al., Impacts of ENSO on air-sea oxygen exchange: Observations and mechanisms, *Global Biogeochem. Cy.*, 31, 901–921, 2017.
- Ishidoya, S., et al., Secular change in atmospheric Ar/N_2 and its implications for ocean heat uptake and Brewer-Dobson circulation, *Atmos. Chem. Phys.*, 21, 1357–1373, 2021.
- Ishidoya, S., et al., Spatiotemporal variations of the $\delta(\text{O}_2/\text{N}_2)$, CO_2 and $\delta(\text{APO})$ in the troposphere over the Western North Pacific, *Atmos. Chem. Phys.*, 22, 6953–6970, 2022.
- Tohjima, Y., et al., Global carbon budgets estimated from atmospheric O_2/N_2 and CO_2 observations in the western Pacific region over a 15-year period, *Atmos. Chem. Phys.*, 19, 9269–9285, 2019.
- Weiss, R. F., The solubility of nitrogen, oxygen and argon in water and seawater, *Deep-Sea Res.*, 17,

721–735, 1970.

Keywords: Atmospheric Potential Oxygen (APO), Atmospheric Ar/N₂ ratio, Interannual variations