

## Large-scale FAC pattern and SW-M-I coupling 2

\*中溝 葵<sup>1</sup>、吉川 顕正<sup>2,5</sup>、中田 裕之<sup>3</sup>、深沢 圭一郎<sup>4</sup>、田中 高史<sup>5</sup>

\*Aoi Nakamizo<sup>1</sup>, Akimasa Yoshikawa<sup>2,5</sup>, Hiroyuki Nakata<sup>3</sup>, Keiichiro Fukazawa<sup>4</sup>, Takashi Tanaka<sup>5</sup>

1. 情報通信研究機構 電磁波研究所、2. 九州大学大学院理学研究院、3. 千葉大学大学院工学研究院、4. 総合地球環境研究所、5. 九州大学国際宇宙惑星環境研究所

1. Applied Electromagnetic Research Institute, National Institute of Information and Communications Technology, 2. Faculty of Science, Kyushu University, 3. Graduate School of Engineering, Chiba University, 4. Research Institute for Humanity and Nature, 5. ICSWSE

The motivation for this study arises from a question: While it is observationally known that the Field-Aligned Current (FAC) patterns change depending on factors such as the IMF orientation, can they also be influenced by ionospheric conductance? Here note that we are targeting the FAC pattern, not the electric potential pattern for which we have reported on various types of deformation and its causes in our past research.

Since FAC is driven by the SW-M-I dynamics, we investigate this issue using a global MHD model. We performed the simulations with various combinations of conductance distributions and IMF orientations:

- (1) Uniform conductance distribution
- (2) Distribution including only the increase due to solar illumination
- (3) Distribution including an increase in the auroral zone

IMF-By values were considered as zero, positive, and negative.

As an initial result, we found that when there is auroral enhancement in the conductance distribution, the FAC patterns exhibit dawn-dusk asymmetry.

We will show further details, including other results and discussions related to SW-M-I coupling.

キーワード：磁気圏電離圏結合、沿磁力線電流、磁気圏シミュレーション

Keywords: M-I coupling, Field aligned current, Magnetosphere simulation