

サンマ (*Cololabis saira*) の加入量変動に対して成長と水温が与える影響に関するモデリング

Modeling effects of growth and temperature on the recruitment variability of Pacific saury (*Cololabis saira*)

*小柳津 瞳¹、伊藤 進一¹、安倍 大介²、伊藤 幸彦¹

*Hitomi Oyaizu¹, Shin-ichi Ito¹, Daisuke Ambe², Sachihiko Itoh¹

1. 東京大学大気海洋研究所、2. 水産研究・教育機構 中央水産研究所

1. Atmosphere and Ocean Research Institute, The University of Tokyo, 2. National Research Institute of Fisheries Science, Japan Fisheries Research and Education Agency

Pacific saury (*Cololabis saira*) is a commercially and ecologically important pelagic fish in the North Pacific. The variability in stock abundance cannot be explained solely by fisheries catch but also related to the reproductive success. In this study, we examine the recruitment variability of Pacific saury using an individual-based model combining a bioenergetics, migration and mortality models. We parameterize the mortality rate with the weight, growth rate and temperature. The annual survival rates (recruitment per spawning biomass: RPS) from the model (mRPS) are calculated from the number of survived fish at age-1, and compared with RPS derived from the stock assessment for 2003–2012. The interannual variability in RPS is well reproduced in the model, especially in cases parameterizing the mortality using the weight and temperature, and weighting the spring-spawned cohort. The importance of the spring-spawned cohort is consistent with the hypothesis derived from observations in 1990–1998.

キーワード：サンマ、Individual-based model、加入量変動、成長、水温

Keywords: Pacific saury, Individual-based model, recruitment variability, growth, temperature