

## Numerical simulation on the albedo of rough surfaces

\*千秋 博紀<sup>1</sup>、諸田 智克<sup>2</sup>、横田 康弘<sup>3</sup>、坂谷 尚哉<sup>4</sup>

\*Hiroki Senshu<sup>1</sup>, Tomokatsu Morota<sup>2</sup>, Yasuhiro Yokota<sup>3</sup>, Naoya Sakatani<sup>4</sup>

1. 千葉工業大学、2. 名古屋大学、3. 高知大学、4. 明治大学

1. Chiba Institute of Technology, 2. Nagoya University, 3. Kochi University, 4. Meiji university

The bond albedo, or energy reflectance, is one of the most important physical parameters because it decides the thermal evolution of the surface. However the bond albedo could change with the roughness of the surface. Although the Hapke's parameter is often introduced to represent the effect of roughness on the phase function, the parameter is not straightforward.

We developed a new numerical model to simulate the image of a rough surface and by using the model calculated the bond albedo of the rough surface.

According to our model, even for the case with Lambertian polygons the bulk bond albedo depends not only on the roughness of the surface but also on the solar incident angle, because of heterogeneous distribution of irradiated area. In this presentation we will propose a new equation to calculate the bond albedo of a rough surface.

キーワード：アルベド、凹凸表面、熱進化

Keywords: albedo, rough surface, thermal evolution